

1 **Reconstructing North Atlantic deglacial surface**
2 **hydrography and its link to the Atlantic overturning**
3 **circulation**

4 David J.R. Thornalley*, Harry Elderfield & I. Nick McCave.

5

6 The Godwin Laboratory for Palaeoclimate Research, Department of Earth Sciences, University of
7 Cambridge, Downing Street, Cambridge, CB2 3EQ, UK.

8 *Present address: School of Earth and Ocean Sciences, Cardiff University, Main Building, Park Place,
9 Cardiff, CF10 3YE, UK.

10 Correspondence should be addressed to DJRT. Email: d.thornalley@cantab.net

1 **ABSTRACT**

2 Paired Mg/Ca- $\delta^{18}\text{O}$ measurements on multiple species of planktic foraminifera are
3 combined with published benthic isotope records from south of Iceland in order to
4 assess the role North Atlantic freshwater input played in determining the evolution of
5 hydrography and climate during the last deglaciation. We demonstrate that
6 *Globigerina bulloides* and *Globorotalia inflata* are restricted to intervals when warm
7 Atlantic waters reached the area south of Iceland, and therefore Mg/Ca- $\delta^{18}\text{O}$ data
8 from these species monitor changes in the temperature and seawater $\delta^{18}\text{O}$ signature of
9 the northward inflow of Atlantic water to the area. In contrast, *Neogloboquadrina*
10 *pachyderma* (sinistral) calcifies within local subpolar/polar waters and new Mg/Ca-
11 $\delta^{18}\text{O}$ analyses on this species document changes in this water mass. We observe two
12 major surface ocean events during Heinrich Stadial 1 (~17-14.7 ka): an early
13 freshening of the Atlantic Inflow (~17-16 ka), and a later interval (16-14.7 ka) of local
14 surface freshening, sea-ice formation and brine rejection that was associated with a
15 further reduction in deep ocean ventilation. Centennial-scale cold intervals during the
16 Bølling-Allerød (BA, 14.7-12.9 ka) were likely triggered by the rerouting of North
17 American continental run-off during ice-sheet retreat. However, the relative effects of
18 these freshwater events on deep ventilation and climate south of Iceland appears to
19 have been modulated by the background climate deterioration. Two freshwater events
20 occurred during the Younger Dryas cold interval (YD, 12.9-11.7 ka), both
21 accompanied by a reduction in deep ventilation south of Iceland: an early YD
22 freshening of the Atlantic Inflow and local subpolar/polar waters, and a late YD ice-
23 rafted detritus event that was possibly related to brine formation south of Iceland.
24 Based on our reconstructions, the strengthening of the Atlantic Meridional
25 Overturning Circulation at the onset of BA and Holocene may have been promoted by

1 the subsurface warming of subpolar/polar water, brine formation that drew warm
2 saline Atlantic water northwards, and the high background salinity of the Atlantic
3 Inflow.

4 **Keywords: Deglaciation, North Atlantic, melt water, climate, temperature,**
5 **salinity**

1 **1. INTRODUCTION**

2 Earth's climate during the Middle and Late Pleistocene has been dominated by ~100
3 000 year timescale glacial-interglacial cycles, associated with ice-sheet growth and
4 decay and sea-level change (e.g. Clark et al., 1999; Waelbroeck et al., 2002; Siddall et
5 al., 2003; Clark et al., 2009), fluctuating global temperatures (NGRIP members, 2004;
6 Jouzel et al., 2007) and atmospheric carbon dioxide concentrations (Lüthi et al., 2008
7 and references therein). A long-standing goal of palaeo-climate research has been to
8 understand the mechanism(s) responsible for governing glacial terminations, which
9 are typified by the transition from the Last Glacial Maximum (LGM, ~26-19 ka, Clark
10 et al., 2009) to the present warm interglacial – the Holocene (~11-0 ka). (In this paper
11 all ages are in ka before 2000 (b2k).)

12 It is widely accepted that changes in solar insolation (indexed by summer
13 insolation at 65 °N), due to orbital variability, are the likely driver for glacial
14 terminations. However, at least two observations can be called upon to demonstrate
15 the complexity of glacial terminations, thereby implicating the involvement of other
16 components of the climate system: (1) The last termination (T-I, ~19-7 ka) involved a
17 number of abrupt centennial to millennial timescale climate jumps and reversals (e.g.,
18 NGRIP members, 2004), such as the Younger Dryas cold interval (YD, 12.9-11.7 ka),
19 which cannot solely be attributed to a gradual change in solar insolation. (2) Recent
20 work by Cheng et al. (2009) has demonstrated that earlier terminations do not all
21 follow the same sequence of events, e.g. Terminations I and III (T-I, T-III) both
22 contain abrupt warm intervals (the Bølling-Allerød, BA, and its T-III equivalent)
23 bracketed by two cold intervals (the Younger Dryas and the Heinrich Stadial/'Mystery
24 Interval' and their equivalents) whilst Termination II and IV only consist of a single
25 Heinrich Stadial/'Mystery Interval' cold event. A likely cause for this complexity is

1 the interplay between freshwater discharge from decaying ice-sheets and changes in
2 the mode of circulation of the Atlantic meridional overturning circulation (AMOC)
3 (Keigwin et al., 1991; Bjorck et al., 1996; Sarnthein et al., 2000; Clark et al., 2001;
4 Carlson, 2008; Thornalley et al., 2010a; Obbink et al., 2010); although the possible
5 involvement of changing atmospheric circulation patterns (e.g., Seager & Battisti,
6 2007; Eisenman et al., 2009) and low-latitude processes (e.g., Clement & Peterson,
7 2008) should not be neglected.

8 In their examination of the mechanisms of the past four terminations, Cheng et
9 al. (2009), building upon previous studies (e.g., Clark et al., 2004; 2009), propose that
10 rising Northern Hemisphere summer insolation triggered the initial disintegration of a
11 massive, isostatically compensated Laurentide ice-sheet. The discharge of ice and
12 freshwater, occurring as a Heinrich Event (Hemming, 2004), caused a reduction in the
13 AMOC, leading to the ensuing cold Heinrich Stadial interval over the North Atlantic
14 (note, in this study, the term ‘Heinrich Event’ refers to an ice-sheet collapse and
15 discharge of ice, whereas ‘Heinrich Stadial’ refers to the entire cold interval – this was
16 not continually occupied by IRD). This North Atlantic cold anomaly caused a
17 southward shift in atmospheric circulation patterns and a build up of heat in the South
18 Atlantic via the bipolar see-saw effect (Stocker & Johnsen, 2003; Barker et al., 2009).
19 Consequently, ventilation and upwelling of CO₂ rich deep waters was enhanced in the
20 Southern Ocean (Toggweiler et al. 2006; Anderson et al., 2009; Barker et al; 2009)
21 and CO₂ was released to the atmosphere, promoting further global warming.
22 Following these initial events, further feedbacks, such as albedo changes and
23 alkalinity-based feedbacks between sea-level and atmospheric CO₂, likely contributed
24 to the full glacial-interglacial transition (Cheng et al., 2009).

1 The complexity of T-I and T-III, in contrast to the simpler one-step
2 termination observed for T-II and T-IV, has been related to a slower rate of insolation
3 rise (Ruddiman et al., 1980; Oppo et al., 1997; Carlson, 2008; Cheng et al., 2009).
4 During T-I and T-III, the disintegration of circum-North Atlantic ice-sheets and
5 freshwater flux to the North Atlantic was sufficiently slow to allow a transient
6 reinvigoration of the AMOC. However, with climate in an intermediate state and
7 remnant large circum-Atlantic ice-sheets, strong NADW formation was unstable and
8 continued melt-water input eventually caused weakening of the AMOC. The
9 subsequent North Atlantic cooling (e.g., the YD) facilitated the final release of CO₂ to
10 the atmosphere. A further factor determining the finer scale structure of terminations
11 was probably the precise geometry of ice-sheet retreat, which will have controlled the
12 rerouting of freshwater input into the North Atlantic (Clark et al., 2001; Obbink et al.,
13 2010; Thornalley et al., 2010a).

14 Southern Hemisphere processes may also play an active role in determining
15 the sequence of events during a termination, with variants on a common theme being
16 proposed (e.g. Keeling & Stephens, 2001; Weaver et al., 2003; Rickaby & Elderfield,
17 2005): Depending upon their respective densities, Antarctic Intermediate Water
18 (AAIW) may compete with North Atlantic Deep Water (NADW) or Glacial North
19 Atlantic Intermediate Water (GNAIW) for occupation of the zone between ~1-2.5 km.
20 When surface water at AAIW formation sites attains a density greater than that found
21 at the deep convection sites in the North Atlantic, NADW formation will decrease,
22 AAIW formation will increase, and vice versa. For example, Weaver et al. (2003)
23 have proposed that an Antarctic origin for meltwater pulse-1a (mwp-1a) may have
24 triggered a reduction in the density of AAIW causing a renewal of NADW formation
25 leading to the onset of the BA.

1 To further investigate the role that freshwater input to the North Atlantic
2 played in determining the centennial to millennial timescale structure of T-I, we
3 examine material from sediment cores located in the northern North Atlantic on the
4 South Iceland Rise. The northern North Atlantic is a sensitive location for
5 ascertaining past changes in the AMOC and freshwater inputs to the high latitude
6 deep convection sites. In this study we provide a synthesis of the palaeoceanographic
7 changes south of Iceland during T-I, drawing upon previously published data and new
8 paired $\delta^{18}\text{O}$ -Mg/Ca measurements. We utilise the habitat preferences of different
9 planktic foraminifera species to reconstruct the changing properties of local
10 subpolar/polar water and inflowing warm Atlantic water. Published paired $\delta^{18}\text{O}$ -
11 Mg/Ca measurements for *Globigerina bulloides* and *Globorotalia inflata* record
12 changes in the warm Atlantic water (Thornalley et al., 2009, 2010a), whilst new
13 paired $\delta^{18}\text{O}$ -Mg/Ca measurements on *Neogloboquadrina pachyderma* (sinistral) from
14 two cores (located ~100 and 200 km south of Iceland) are used to examine the
15 deglacial changes in local subpolar/polar water. These results are compared to
16 published benthic isotope data from the South Iceland Rise depth transect (Thornalley
17 et al., 2010b).

18

19 **2. REGIONAL SETTING**

20 The four cores used in this study are located on the South Iceland Rise beneath the
21 surface flow of a major branch of the North Atlantic Current and the deep flow of
22 Iceland-Scotland Overflow Water (Figure 1).

23 The surface circulation of the northern North Atlantic consists of a warm
24 saline northeastward flow from Newfoundland to the Nordics Seas (the North Atlantic
25 Current, NAC) that draws water from the cold, fresh, subpolar gyre (SPG) and warm,

1 saline, subtropical gyre (STG). The major cold surface current of the Northeast
2 Atlantic is the East Greenland Current (EGC) which flows south around Greenland
3 and into the Labrador Sea. The warm NAC that enters the Nordic Seas (the Atlantic
4 Inflow) cools, sinks and returns southwards as Iceland-Scotland and Denmark Straits
5 Overflow Water (ISOW and DSOW). These overflows entrain adjacent subsurface
6 water masses and after passing around the southern tip of Greenland combine with
7 Labrador Sea Water (LSW, formed by deep convection in the Labrador Sea) to form
8 North Atlantic Deep Water (NADW).

9 Modern inter-annual variability in the temperature and salinity of the Atlantic
10 Inflow is controlled by subpolar gyre dynamics: enhanced freshwater input to the
11 Labrador Sea causes the SPG to retract westwards allowing warm saline STG water to
12 dominate the Inflow (Hátún et al., 2005). Episodically, the ocean south of Iceland is
13 also influenced by sea-ice and near surface subpolar water that is advected, via the
14 East Icelandic Current (EIC), southwards out of the Nordic Seas, typically when
15 atmospheric circulation is in the negative phase of the North Atlantic Oscillation, such
16 as during the late 1960s (Blindheim & Østerhus, 2005).

17 During the LGM and cold stadial intervals, sea-ice and subpolar/polar waters
18 extended south of Iceland (to at least 60 °N (Pflaumann et al., 2003)). However, there
19 were episodic, likely subsurface, incursions of warm Atlantic water (e.g. Rasmussen
20 & Thomsen, 2004). Freshwater input from surrounding ice sheets and extensive sea
21 ice cover severely limited deep open ocean convection in the Nordic Seas (Labeyrie et
22 al., 1992). Instead, there was possible brine formation in the Nordic Seas (Meland et
23 al., 2008) and convection south of Iceland forming Glacial North Atlantic
24 Intermediate Water (GNAIW) ventilating the North Atlantic to ~2 km depth, below

1 which water of southern source filled the ocean basin (Oppo & Lehman 1993; Lynch
2 Stieglitz et al., 2007).

3

4 **3. MATERIALS AND METHODS**

5 **3.1 Age models**

6 We use previously published age models, details of which can be found in Thornalley
7 et al. (2009, 2010a, and 2010b). Because of the variable surface reservoir age of the
8 deglacial northern North Atlantic, deglacial age models are based on tying sharp
9 changes in the percent abundance of the subpolar/polar species, *N. pachyderma*
10 sinistral (*Nps*), with abrupt climate transitions in the NGRIP ice core. Additional
11 robust tie points are provided by Icelandic tephra layers found in both the marine
12 cores and NGRIP. The combination of Holocene radiocarbon dates, and tephra and %
13 *Nps* correlation after ~14.7 ka (the onset of the BA), provides tight age control for the
14 cores. Yet prior to the BA, considerable uncertainty exists. LGM ¹⁴C dates use an
15 assumed reservoir age of 800 years, and an additional tie-point is provided at 17.2 ka
16 by correlating benthic $\delta^{18}\text{O}$ between the RAPiD cores and the well-dated core SU81-
17 18 (38°N, 10°W) (Waelbroeck et al., 2001), located at a latitude where changes in the
18 surface radiocarbon reservoir age are thought to have been relatively small
19 (Waelbroeck et al., 2001). This assumes that the decrease in benthic $\delta^{18}\text{O}$ at the onset
20 of Heinrich Stadial 1 (HS-1, ~14.7 – 17 ka) was approximately synchronous across
21 the Northeast Atlantic. Caution should therefore be used when comparing the timing
22 of climate events recorded by the RAPiD cores between ~15 ka and ~19 ka with
23 existing datasets.

24

25 **3.2 Core quality**

1 All cores were acquired on Charles Darwin cruise 159 for the RAPID programme of
2 NERC (McCave, 2005). P denotes a piston core and K a kasten (15 cm square
3 gravity) core.

4 **RAPiD-10-1P** (62°58.53' N, 17°35.37' W, 1237 m water depth): The
5 proximal location to Iceland (~100 km) provides high sedimentation rates (50-60
6 cm.ka⁻¹) during the LGM, HS-1 and the YD. Lower sedimentation rates occur during
7 the BA and Holocene (12 and 4 cm.ka⁻¹ respectively) (Thornalley et al., 2010b).

8 **RAPiD-12-1K** (62°05.43' N, 17°49.18' W, 1938 m water depth): High
9 sedimentation rates (~125 cm.ka⁻¹) during the early Holocene (~ 8 to 11.7 ka) were
10 caused by the input of tephra and the liberation of sediment during the deglaciation of
11 Iceland. The onset of strong overflows at ~ 8 ka produce a foraminiferal sand deposit
12 (~22 cm.ka⁻¹). The core is well dated, with 13 radiocarbon dates on *G. bulloides*
13 during the Holocene (Thornalley et al., 2009).

14 **RAPiD-15-4P** (62°17.58' N, 17°08.04' W, 2133 m water depth): The core
15 contains an excellent deglacial sequence after ~15.8 ka with sedimentation rates of
16 12-42 cm.ka⁻¹. The effects of bioturbation are minor, with an estimated mixing
17 interval of ~2 cm (Thornalley et al., 2010a). The core contains similar Holocene
18 deposits to RAPiD-12-1K. Recently obtained, unpublished radiocarbon dates on *Nps*
19 suggest that between core depths of 526 and 543 cm (~15.8 to 17.2 ka) the core
20 contains some reworked older material. Because only the most pristine foraminifera
21 were picked for isotope and trace metal analyses this reworked material was
22 successfully avoided, as confirmed by the similar benthic and planktic stable isotope
23 data and planktic Mg/Ca data from nearby cores, which show no signs of reworking
24 (Figure 2 and 3).

1 **RAPiD-17-5P** (61°28.90' N, 19°32.16' W, 2303 m water depth): The core
2 contains an excellent Holocene section with high sedimentation rates (~80 cm.ka⁻¹).
3 Lower sedimentation rates occur during the deglaciation: HS-1, 7 cm.ka⁻¹; BA, 9-15
4 cm.ka⁻¹; YD, 15 cm.ka⁻¹ (Thornalley et al., 2010b).

5

6 **3.3 *N. pachyderma* sinistral stable isotope and Mg/Ca measurements**

7 Paired δ¹⁸O-Mg/Ca analysis of planktic foraminifera can be used to reconstruct past
8 ocean temperatures and seawater δ¹⁸O (δ¹⁸O_{sw} – a property that shows a near linear
9 relationship with salinity in the modern surface ocean) (Elderfield & Ganssen, 2000).

10

11 100-200 tests of the planktic foraminifera *Neogloboquadrina pachyderma* sinistral
12 (*Nps*) were picked from the 150-250 μm fraction. Samples were crushed,
13 homogenised and split to produce ~100-150 μg for stable isotopic analysis, the
14 remainder being used for Mg/Ca analysis. Stable isotope measurements were
15 performed using the Godwin Laboratory VG Prism mass spectrometer attached to a
16 Micromass Multicarb Sample Preparation System. Measurements of δ¹⁸O and δ¹³C
17 were determined relative to the Vienna Peedee belemnite (VPDB) standard, with an
18 analytical precision better than 0.08 ‰ and 0.06 ‰ respectively. Mg/Ca cleaning was
19 conducted following the methods of Barker et al. (2003), screening for contaminating
20 ferromanganese growths, clay minerals and silicate particles. Samples were dissolved
21 and diluted to a [Ca] of 100 ppm and analysed using a Varian Vista inductively
22 coupled plasma optical emission spectrometer (ICP-OES). Analytical precision of
23 Mg/Ca ratios based on replicates of foraminiferal standards is 3%.

24

25 **3.4 *N. pachyderma* sinistral calibrations**

1 To reconstruct past temperature and $\delta^{18}\text{O}_{\text{sw}}$ we assume a 9-10 % sensitivity for the
2 Mg/Ca ratio of planktic foraminifera per degree Celsius temperature change
3 (Elderfield & Ganssen, 2000; Anand et al., 2003; Lea, 2003), and equilibrium
4 precipitation of inorganic calcite (Kim & O'Neil, 1997). The temperature effect was
5 then removed from measured calcite $\delta^{18}\text{O}$ to yield $\delta^{18}\text{O}_{\text{sw}}$. These values were then
6 corrected for the whole ocean change in $\delta^{18}\text{O}$ of ~ 1 ‰ that occurred during T-I by
7 scaling to the sea-level curve of Fairbanks et al. (1989) using an LGM to late
8 Holocene sea-level change of 120 m. This produces 'ice-volume corrected' seawater
9 $\delta^{18}\text{O}$ ($\delta^{18}\text{O}_{\text{sw-ivc}}$).

10 The Mg/Ca calibrations for the published *G. bulloides* and *G. inflata* data
11 (Thornalley et al., 2009; Thornalley et al., 2010a) use an equation of the form: Mg/Ca
12 = $B \exp(0.1T)$ where T is temperature and the constant B is calibrated to the core tops
13 yielding values of 0.794 and 0.675 for *G. bulloides* and *G. inflata* respectively.
14 Alternative Mg/Ca calibrations for *G. inflata* have been developed, and differences
15 between the calibrations may be caused by a preference for different studies to use a
16 different morphotype (Hathorne et al., 2009). Consistent with the use of a 9-10 %
17 sensitivity of Mg/Ca per degree Celsius temperature, we employ the Mg/Ca
18 calibration of Elderfield & Ganssen (2000) for our *Nps* data: Mg/Ca = $0.52 \exp(0.1T)$.
19 The absence of *Nps* within late Holocene sediments on the South Iceland Rise
20 prevented confirmation of the value of B . The calibration of Elderfield & Ganssen
21 (2000) is utilised rather than the recent calibration by Kozdon et al. (2009a) because
22 the Elderfield & Ganssen (2000) calibration has successfully been applied to a study
23 of late Holocene sediments in the nearby Nordic Seas, with these samples being run in
24 the same laboratory and following the same methods as this study (Nyland et al.,

1 2006). Use of the Kozdon et al. (2009a) calibration would lower the reconstructed
2 temperatures by approximately 0.3-1.5 °C.

3 It has been widely demonstrated that planktic species do not always precipitate
4 calcite in equilibrium. In accordance with the reconstructed temperatures south of
5 Iceland (7-12 °C), and the culture study work of Bemis et al (1998), we use a $\delta^{18}\text{O}$
6 offset of -0.6‰ for *G. bulloides*. Numerous authors have also observed that post-
7 gametogenetic processes cause *Nps* not to calcify in equilibrium with $\delta^{18}\text{O}_{\text{sw}}$ and a
8 disequilibrium factor of ~0.6‰ has been widely used (Stangeew, 2001; Simstich et
9 al., 2003; Nyland et al., 2006). Studies from Arctic waters suggest a higher
10 disequilibrium of ~1‰ (Nyland et al., 2006 and references therein) whilst a study
11 from the Icelandic continental shelf suggests a 0.25‰ offset (Smith et al., 2005) and
12 sediment trap data from the Irminger Sea indicate no offset (Jonkers et al., 2010).
13 Kozdon et al. (2009b) have demonstrated that this offset factor can be highly variable
14 due to a ~ 2.1‰ difference in isotopic composition of the ontogenetic calcite and crust
15 calcite. This study therefore urges caution when interpreting down-core $\delta^{18}\text{O}_{\text{sw}}$
16 records which may vary independently of temperature or salinity. The replication of
17 down-core trends in $\delta^{18}\text{O}_{\text{sw}}$ between two cores provides some additional confidence
18 that our reconstructed $\delta^{18}\text{O}_{\text{sw}}$ data reflect real changes in $\delta^{18}\text{O}_{\text{sw}}$. In this study we use
19 an offset of 0.6‰, which was successfully used in the Nordic Seas study of Nyland et
20 al. (2006).

21

22 **4. RESULTS**

23 **4.1 Planktic foraminifera habitat preferences**

24 Modern global core top studies show that *G. bulloides* and *G. inflata* become a
25 significant component of the planktic foraminiferal assemblage when summer sea

1 surface temperatures (SSTs) exceed $\sim 7^{\circ}\text{C}$ (Hilbrecht, 1996). This limitation implies
2 that during the deglaciation, the presence of *G. bulloides* and *G. inflata* will largely be
3 restricted to intervals when relatively warm Atlantic water is advected to the south of
4 Iceland. In contrast, *Nps* abundance increases dramatically when summer SSTs fall
5 below $\sim 7^{\circ}\text{C}$ (Hilbrecht, 1996), when subpolar/polar waters dominate south of Iceland.
6 Reconstructed Mg/Ca-temperatures for the three species are shown in **figure 4** and the
7 data confirm their expected habitat preferences.

8 The transitional species, *G. inflata* (and to a lesser extent, the warm subpolar
9 species, *G. bulloides*), will only be present in the sediment cores when relatively
10 warm Atlantic water was located over the South Iceland Rise, and paired Mg/Ca- $\delta^{18}\text{O}$
11 measurements on these species will reconstruct the changing properties of the Atlantic
12 Inflow. *G. inflata* $\delta^{18}\text{O}_{\text{sw-ivc}}$ will be particularly sensitive at detecting the advection of
13 Laurentide Ice Sheet (LIS) freshwater input to the NE Atlantic via the NAC
14 (Thornalley et al., 2010a). In contrast, measurements on *Nps* will record the changing
15 properties of local subpolar/polar waters south of Iceland. A further consideration is
16 the depth habitats of the species: *G. bulloides* lives in the near-surface layer ($\sim 0\text{-}50$ m,
17 Ganssen & Kroon, 2000), where it may be influenced by any thin overlying
18 freshwater layer; *G. inflata* lives at the base of the seasonal thermocline ($\sim 100\text{-}200$ m
19 south of Iceland at present)(Ganssen & Kroon, 2000; Cleroux et al., 2007); *Nps*
20 calcifies in the mixed layer and then sinks below the pycnocline to reproduce, where it
21 gains its secondary calcite, resulting in an apparent calcification depth of $\sim 100\text{-}200$ m
22 (Simstich et al., 2003). Because none of these species are true sea-surface dwellers,
23 surface freshening will be underestimated, a phenomenon that is well documented for
24 *Nps* (Simstich et al., 2003). Furthermore, it has been suggested that *Nps* cannot
25 tolerate salinities $< 34\text{-}34.5$ p.s.u and rather than reflecting strong surface freshening,

1 extremely low *Nps* $\delta^{18}\text{O}_{\text{sw}}$ is a likely indicator of local brine formation (Hillaire-
2 Marcel & deVernal, 2008).

3 The habitat preference of different planktic foraminifera species underpins
4 modern analogue techniques, which use changes in faunal assemblage to reconstruct
5 past SST change (e.g. Waelbroeck et al., 1998). In the North Atlantic the percent
6 abundance of *Nps* can be used to estimate past mean SSTs: 95% abundance of *Nps*
7 indicates an SST of $<4^\circ\text{C}$ and when *Nps* abundance drops to $<20\%$ an SST of $\sim 9^\circ\text{C}$ is
8 indicated (Bé & Tolderlund, 1971). Down-core % *Nps* data is presented in figure 3.

9 Combining faunal assemblage data and paired Mg/Ca- $\delta^{18}\text{O}$ on different
10 species allows a comprehensive reconstruction of the hydrography of the upper ocean.
11 Down-core variations in % *Nps* reflect mean SST changes averaged over the
12 bioturbated sampling intervals (typically 50-200 years), whereas paired Mg/Ca- $\delta^{18}\text{O}$
13 measurements record the changing properties of different water masses (the Atlantic
14 Inflow and subpolar/polar water). The two datasets are generally consistent with one
15 another, as can be demonstrated using simple assumptions (i.e., *G. bulloides*/*G. inflata*
16 Mg/Ca-temperatures record Atlantic Inflow temperatures and *Nps* Mg/Ca-
17 temperatures record subpolar/polar water temperatures, and the proportionate time
18 each water mass is present at the core site is approximated by the % *Nps*). For
19 example, during the early YD, the abundance of *Nps* is $\sim 85\%$, indicating an SST of
20 $\sim 4.5\text{-}5^\circ\text{C}$, whilst the weighted mean Mg/Ca temperature produces a temperature of
21 4.6°C ($85\% \times 4^\circ\text{C} + 15\% \times 8^\circ\text{C}$). Similar calculations can be made for other time
22 intervals, e.g. the early Holocene ($>9^\circ\text{C}$; 10.2°C) or the late Bølling ($\sim 6^\circ\text{C}$; 6.5°C).

23

24 **4.2 Down-core *N. pachyderma* (sinistral) Mg/Ca-d ^{18}O**

1 Paired Mg/Ca- $\delta^{18}\text{O}$ measurements for *Nps* from RAPiD-10-1P and RAPiD-15-4P are
2 presented in figure 3. Mg/Ca ratios indicate an overall warming from the LGM to the
3 Holocene. Mg/Ca ratios increase by $0.1 \text{ mmol.mol}^{-1}$ (equivalent to a 1.2°C warming)
4 between 17 and 16 ka, and fluctuations in Mg/Ca and $\delta^{18}\text{O}$ during the BA from the
5 high resolution core RAPiD-15-4P are in phase with changes in % *Nps*, which
6 correlate with abrupt multi-centennial timescale cooling and warming of Greenland
7 air temperatures (Thornalley et al., 2010a). RAPiD-15-4P *Nps* Mg/Ca ratios then
8 suggest cooling during the early YD and warming during the late YD, continuing into
9 the Holocene. $\delta^{18}\text{O}_{\text{sw-ivc}}$ displays low values between 16 ka and 14.8 ka and
10 subsequent fluctuations occur throughout the BA.

11 The results from the two cores are generally consistent with one another
12 (figure 3). A notable difference occurs during Heinrich Stadial 1 (HS-1, and to a
13 lesser extent, the BA), when RAPiD-10-1P records significantly lower $\delta^{18}\text{O}_{\text{sw-ivc}}$.
14 Based on the low *Nps* $\delta^{18}\text{O}_{\text{sw-ivc}}$ in RAPiD-10-1P between 14.8-16 ka, there was
15 significant surface freshening south of Iceland and it is likely that active sea-ice
16 formation and brine rejection occurred. The more extreme $\delta^{18}\text{O}_{\text{sw-ivc}}$ in RAPiD-10-1P
17 presumably reflects the core's proximal location to Iceland where it will be influenced
18 by freshwater from the nearby Icelandic ice-sheet and active coastal sea-ice
19 formation. The suggested brine formation during this interval is consistent with low
20 benthic $\delta^{18}\text{O}_{\text{ivc}}$ (Figure 5 and Thornalley et al., 2010b).

21

22 **5. DISCUSSION**

23 **5.1 The phasing of surface and deep ocean changes**

24 In **figure 5** the phasing between surface and deep ocean records is examined, with the
25 primary aim of assessing the relationship between freshwater input and deep ocean

1 circulation changes. Accurate phase relationships can be inferred throughout the
2 deglacial because the derived surface temperatures and $\delta^{18}\text{O}_{\text{sw-ivc}}$ and benthic isotope
3 data have all been obtained from RAPiD-15-4P.

4

5 **5.1.1 The Last Glacial Maximum and Early Deglacial (17-21 ka)**

6 Benthic $\delta^{18}\text{O}_{\text{ivc}}$ from the depth transect south of Iceland support the traditional view
7 that GNAIW ventilated the North Atlantic above ~2 km, with southern source water
8 below (Boyle and Keigwin, 1987; Oppo et al., 1993, Lynch-Stieglitz et al., 2007;
9 Thornalley et al., 2010b).

10 Previous LGM reconstructions (Glacial Atlantic Ocean Mapping Project,
11 GLAMAP, Pflaumann et al., 2003) indicate extensive winter sea-ice cover over the
12 South Iceland Rise, with winter SSTs ~0.5-2°C, and summer SSTs ~3.5-5°C,
13 consistent with our *Nps* Mg/Ca-temperatures of 3-4.5°C and high % *Nps* abundance
14 (figure 3). The relatively saline conditions indicated by our *Nps* $\delta^{18}\text{O}_{\text{sw-ivc}}$ data imply
15 that *Nps* was likely calcifying below any near surface freshwater layer that had
16 formed by melting winter sea-ice, analogous to their modern habitat preferences in the
17 Nordic Seas (Simstich et al., 2003). Winter sea-ice coverage likely limited deep
18 convection south of Iceland, although the high inferred *Nps* salinity suggests that in
19 polynyas deep mixing forming GNAIW may have been possible.

20 *Nps* $\delta^{18}\text{O}_{\text{sw-ivc}}$ does not show any significant freshwater event between 21 ka
21 and 17 ka that can be related to the sea-level rise and melt-water pulse that occurred at
22 ~19 ka, likely sourced from Northern Hemisphere ice-sheets (Clark et al., 2004). An
23 explanation for this discrepancy may either be the low sensitivity of *Nps* $\delta^{18}\text{O}_{\text{sw-ivc}}$ at
24 detecting surface freshening, and/or, the freshwater was not advected within
25 subpolar/polar waters to the South Iceland Rise. (Note - *G. inflata* and *G. bulloides*

1 $\delta^{18}\text{O}_{\text{sw-ivc}}$, that would monitor changes in Atlantic water, have not been measured due
2 to the low abundance of these species during this interval.)

3

4 **5.1.2 Heinrich Stadial 1 (14.7-17 ka)**

5 IRD concentration increase in the Nordic Seas and south of Iceland between 17.6 ka
6 and 16 ka, remaining high until the onset of the BA at 14.7 ka, suggests a significant
7 influence of ice-bergs and associated freshwater (figure 5). It is likely that the $\sim 1.5^\circ\text{C}$
8 warming recorded by *Nps* at ~ 16 -17 ka was associated with this surface freshening,
9 since upper water column stratification will have increased, thereby preventing
10 atmospheric cooling of the subsurface water within which *Nps* gained its secondary
11 calcite. Other studies also suggest the North Atlantic subsurface warmed during HS-1
12 (e.g. Shaffer et al., 2004; Clark et al., 2007; Knorr & Lohman, 2007; Liu et al., 2009;
13 Alvarez-Solas et al., 2010), although the modelling study by Knorr & Lohman (2007)
14 suggests the cause was increased horizontal mixing with intermediate and upper ocean
15 subtropical water masses that warmed in response to the reduced AMOC – consistent
16 with the warm Atlantic Inflow temperatures recorded during HS-1 south of Iceland.

17 Northeast (NE) Atlantic IRD data record two distinct peaks during HS-1: a
18 detrital carbonate IRD peak (Heinrich Event 1B, H1B) at ~ 17 ka (characteristic of the
19 LIS) and a later peak (Heinrich Event 1A, H1A) containing haematite-coated and
20 volcanic grains at 15-16 ka, more characteristic of the ice-sheets surrounding the NE
21 Atlantic (Bard et al., 2000; Peck et al., 2007). (It should be noted that NE Atlantic
22 precursor IRD events prior to the detrital carbonate peak have also been identified
23 (e.g. Bond & Lotti, 1995; Elliot et al., 1998; Peck et al., 2007; Hall et al., 2006).) The
24 detrital carbonate IRD peak is accompanied by low *G. bulloides* and *G. inflata* $\delta^{18}\text{O}_{\text{sw-ivc}}$
25 which have been attributed to the advection of LIS-sourced freshwater to south of

1 Iceland during H1B (Thornalley et al., 2010a). The absence of a strong freshwater
2 signal in the Atlantic Inflow (*G. inflata* $\delta^{18}\text{O}_{\text{sw-ivc}}$) during the second IRD peak (H1A)
3 is consistent with a NE Atlantic source (rather than LIS source). Further support for a
4 proximal source for H1A is provided by the timing of freshening and inferred sea-ice
5 and brine formation south of Iceland (14.8-16 ka, figure 3).

6 Benthic isotope data (figure 5) can be used to examine the effect of these
7 freshwater events on the deep ocean south of Iceland. Decreasing $\delta^{13}\text{C}$ suggests a
8 reduction in deep ocean ventilation. Decreasing $\delta^{18}\text{O}_{\text{sw-ivc}}$ can be interpreted as an
9 increasing influence of brine during HS-1 (Thornalley et al., 2010b). However, the
10 inference of brine based on low benthic $\delta^{18}\text{O}$ is a matter of ongoing debate (Dokken
11 and Jansen, 1999; Bauch and Bauch, 2001; Meland et al., 2008; Rasmussen and
12 Thomsen, 2009a and 2009b; Thornalley et al., 2010b). Rasmussen and Thomsen
13 (2009a) suggest that in the modern Nordic Seas, brines with a low $\delta^{18}\text{O}$ cannot reach
14 intermediate depths because they do not attain a density great enough to sink below
15 the subsurface saline Atlantic layer. Yet our knowledge of the thickness and salinity
16 of this layer during HS-1 is limited. During HS-1, when deep convection decreased
17 due to enhanced surface freshwater input, the Atlantic Inflow was almost certainly
18 reduced and via diffusion the subsurface may have freshened. Combined with intense
19 sea-ice and hence brine formation, low $\delta^{18}\text{O}$ brines plausibly reached greater depths
20 than in today's Nordic Seas.

21 The initial decrease in ventilation south of Iceland at ~17 ka was likely
22 triggered by freshwater discharge from the LIS during H1B and its advection to the
23 NE Atlantic, although earlier European precursor events also began to weaken the
24 AMOC prior to H1B (Hall et al., 2006; Menot et al., 2006; Rinterknecht et al., 2006;
25 Clark et al., 2009). By 15.8 ka, it appears that the source of freshwater had switched

1 from the LIS to the proximal NE Atlantic ice-sheets and deep ventilation was further
2 reduced and intensive brine formation occurred south of Iceland.

3

4 **5.1.3 The Bølling-Allerød**

5 The onset of the BA was accompanied by an increase in deep ocean ventilation, a
6 decrease in IRD south of Iceland and in the Nordic Seas, and a decrease in the %
7 abundance of *Nps*, all of which suggest an enhanced inflow of warm Atlantic water to
8 south of Iceland and the Nordic Seas, and the resumption of open ocean convection in
9 the northern North Atlantic and strong AMOC, consistent with earlier studies (e.g.
10 Dokken & Jansen, 1999; Sarnthein et al., 2000; Robinson et al., 2005)

11 Following the initial warming and resumption of strong AMOC at ~14.7 ka,
12 the BA was characterised by a gradual deterioration in climate, with several multi-
13 centennial cold intervals superimposed upon the longer term cooling trend. These
14 cold intervals have been linked to rerouting of freshwater from the LIS as the ice-
15 margin fluctuated during the deglaciation (Clark et al., 2001; Obbink et al., 2010;
16 Thornalley et al., 2010a;). The freshwater supply to the south of Iceland increased
17 during warm intervals as the LIS retreated and freshwater was diverted eastwards,
18 closer to the high latitude deep convection sites. Peak freshening of the Atlantic
19 Inflow triggered a reduction in AMOC and/or a growth in sea-ice that led to the
20 subsequent cold interval. During this ensuing cold interval, the LIS re-advanced and
21 freshwater was diverted southwards, allowing climate amelioration.

22 Low *Nps* $\delta^{18}\text{O}_{\text{sw-ivc}}$ during the warm intervals of the BA suggest enhanced
23 melting of circum-NE Atlantic ice-sheets which caused freshening of local
24 subpolar/polar water. Local maxima (saline) in *Nps* $\delta^{18}\text{O}_{\text{sw-ivc}}$ occur during cold

1 intervals - the Older Dryas (OD, ~14.0 ka) and the Intra-Allerod Cold Interval (IACP,
2 ~13.0-13.4 ka).

3 The impact of the surface freshwater events can again be assessed by
4 examining benthic isotope data. Benthic $\delta^{13}\text{C}$ from south of Iceland indicate a
5 reduction in deep ocean ventilation during the IACP, although a similar signal cannot
6 be clearly detected during the OD: it is possible that any benthic $\delta^{13}\text{C}$ shift was
7 masked by other effects, such as air-sea-exchange or local productivity changes.

8 The end of the BA, just prior to the onset of the YD, is characterised by a brief
9 (200-300 years) recovery out of the IACP: IRD in the Nordic Seas decreased and
10 there was better ventilation of the deep ocean south of Iceland. It should also be noted
11 that the inferred salinity of the Atlantic Inflow had increased to its high, deglacial
12 background level (Thornalley et al., 2010a)

13

14 **5.1.4 The Younger Dryas**

15 The involvement of freshwater in triggering the Younger Dryas has long been debated
16 and remains unresolved. An early proposal for the cause of the Younger Dryas was
17 the rerouting of LIS meltwater from southern to eastern outlets (e.g. Broecker et al.,
18 1989), but conclusive palaeoceanographic evidence to support this rerouting has
19 remained elusive, with deVernal et al. (1996) finding no evidence for increased
20 freshwater flux through the St. Lawrence. This contrasts with the findings of a later
21 study (Carlson et al., 2007a), although the validity of their freshwater reconstruction
22 methods can be questioned (Peltier et al., 2008). This has led to the suggestion of an
23 Arctic freshwater trigger for the YD (e.g. Tarasov and Peltier, 2005; Murton et al.,
24 2010), but once again, convincing palaeoceanographic evidence of ocean freshening
25 to support the hypothesis is so far absent (Carlson and Clark, 2008). Regardless of the

1 initial trigger, the abrupt increase in IRD in the Nordic Seas suggests that discharge of
2 ice-bergs there played an important role in suppressing AMOC throughout the YD.

3 The onset of the YD (~12.8 ka) was accompanied by a brief (~100 year)
4 freshwater event south of Iceland recorded by *G. inflata* and *G. bulloides* $\delta^{18}\text{O}$ (Event
5 II in Thornalley et al., 2010a), although the event was not captured by our slightly
6 lower resolution $\delta^{18}\text{O}_{\text{sw-ivc}}$ data. These data therefore suggest there was a small
7 freshwater flux to the NE Atlantic via the NAC. However, low *Nps* $\delta^{18}\text{O}_{\text{sw-ivc}}$ persist
8 from the final warm interval of the BA into the early YD, indicating relatively fresh
9 local subpolar/polar water. Since the supply of subpolar/polar waters to south of
10 Iceland largely comes from the Nordic Seas, an Arctic freshwater event cannot be
11 excluded. (Although this reasoning is somewhat tenuous, and the brief freshening of
12 subpolar waters at the onset of the YD may indeed have been cause by the freshening
13 of the Atlantic Inflow.)

14 Examination of the benthic isotope data elucidates the changing
15 palaeoceanography of the South Iceland Rise during the YD. Deep ventilation
16 reduced from 12.8 ka to 12.6 ka, possibly caused by the YD-onset freshwater event.
17 Between 12.6 ka and ~12.1 ka there appears to have been moderate ventilation of the
18 deep ocean south of Iceland, before a final reduction in ventilation between 12.1 ka
19 and 11.7 ka. This later reduction in ventilation coincided with the presence of brines
20 south of Iceland (Thornalley et al., 2010b). It is uncertain whether these brines
21 formed locally (as possibly suggested by low *Nps* $\delta^{18}\text{O}_{\text{sw-ivc}}$) or were exported from
22 the Nordic Seas, as suggested by Meland et al. (2008). The late YD reduction in
23 ventilation coincided with an abrupt increase in IRD south of Iceland, and Heinrich
24 Event 0 (H-0), recorded by cores in the Labrador Sea (e.g., Andrews and Tedesco
25 1992; Hillaire & deVernal, 2008). The increased presence of surface freshwater south

1 of Iceland would also account for increasing *Nps* temperatures during the late YD (H-
2 0), analogous to during HI when it is proposed that stratification of the upper water
3 column prevented atmospheric cooling of the subsurface waters inhabited by *Nps*.
4 (Additionally, an overall warming trend in the temperature of Atlantic Inflow waters
5 between ~13.4 ka (the onset of the IACP) and the early Holocene (~11.5 ka) may
6 indicate that during weakened AMOC, the subtropical waters that fed the Atlantic
7 Inflow warmed (Knorr & Lohman, 2007; Carlson et al., 2008a).

8

9 **5.1.5 The Early Holocene**

10 The onset of the Holocene was associated with an abrupt decrease in % *Nps* and IRD,
11 and an increase in benthic $\delta^{13}\text{C}$ suggests better ventilation of the deep ocean. These
12 changes were caused by the retreat of sea-ice, the resumption of open ocean
13 convection in the Nordic Seas and strong AMOC, that brought warm NAC water to
14 the northern North Atlantic (e.g. Dokken & Jansen, 1999; Sarnthein et al., 2000;
15 Robinson et al., 2005). Because of weak SPG circulation, the background salinity of
16 the Atlantic Inflow was still relatively high during the YD and early Holocene and
17 this may have assisted the resumption of strong AMOC at the onset of the Holocene
18 (figure 6 and Thornalley et al., 2010a).

19 Strong warming during the onset of the Holocene likely caused enhanced
20 melting of the circum-NE Atlantic ice-sheets, as recorded by low *Nps* $\delta^{18}\text{O}_{\text{sw-ivc}}$ south
21 of Iceland. Warming south of Iceland was interrupted by the Preboreal Oscillation
22 (PBO), a ~200 year cold interval at ~11.4 ka possibly triggered by a flood from
23 glacial Lake Agassiz into the Arctic Ocean (Fisher et al., 2002; Meissner and Clark,
24 2006). The PBO is recorded south of Iceland by a decrease in % *Nps* and an inferred
25 increase in salinity of *Nps*, presumably caused by decreased melting of the circum-NE

1 Atlantic ice sheets. Following the PBO, the continued melting and retreat of the LIS
2 triggered melt-water outbursts from Lake Agassiz (Teller et al., 2002), culminating in
3 the final drainage of Lake Agassiz at ~8.4 ka (Barber et al., 1999; Teller et al., 2002
4 and references therein), that is detected as a ~0.5‰ decrease (freshening) in the
5 $\delta^{18}\text{O}_{\text{sw-ivc}}$ of *G. inflata* (Atlantic Inflow water) south of Iceland. (Earlier events at 9.3
6 ka and 10.1 ka are possibly also detected south of Iceland.) Via a reduction in
7 AMOC, the release of Lake Agassiz freshwater at 8.4 ka triggered the ‘8.2 kyr cold
8 event’ (Barber et al., 1999; Ellison et al., 2006; Kleiven et al., 2007), and the
9 particularly low benthic $\delta^{13}\text{C}$ recorded in RAPiD-15-4P at 8.0 ka to 8.2 ka likely
10 indicate an increased incursion of southern source waters.

11 Although there was an abrupt increase in the strength of the AMOC at the
12 onset of the Holocene, the modern circulation regime was not reached until ~7-8 ka
13 (Marchitto et al., 1998; Oppo et al., 2003; Piotrowski et al., 2004; Renssen et al.,
14 2009; Thornalley et al., 2010b). This delay was likely a result of the remnant LIS
15 cooling the North Atlantic region via the ice-sheet’s high surface albedo, and the
16 continued input of LIS melt water preventing deep convection in the Labrador Sea
17 (Hillaire-Marcel et al., 2001; Carlson et al., 2007b; Carlson et al., 2008b; Renssen et
18 al., 2009). By ~7-8 ka benthic $\delta^{13}\text{C}$ south of Iceland had stabilised and reached
19 modern values, ISOW attained its present strength and/or depth on the South Iceland
20 Rise (Thornalley et al., 2010b), and upper ocean circulation resembled the modern
21 regime with strong SPG circulation influencing the Atlantic Inflow (figure 6) and
22 deep winter mixing events homogenising the upper ocean water column (Thornalley
23 et al., 2009).

24

1 **5.2 The role of circum-North Atlantic freshwater input on the centennial-**
2 **millennial structure of Termination I.**

3 Having determined the timing of deglacial freshwater inputs and their relationships
4 with changes in deep ocean ventilation south of Iceland (summarised in **table 1**), it is
5 now possible to examine the broader role of North Atlantic freshwater input in
6 determining the structure of T-I.

7

8 **5.2.1 The impact of early deglacial freshwater events**

9 Despite the early melt-water events (e.g. the 19 ka mwp and early NE Atlantic IRD
10 events) that were associated with a reduction in AMOC strength and warming of the
11 South Atlantic via the bipolar seesaw, it is not until the onset of H1B at ~17 ka that
12 there was a reduction in the ventilation of the glacial deep ocean (~2 km depth) south
13 of Iceland and a substantial increase in upwelling and CO₂ release from the Southern
14 Ocean (Anderson et al., 2009). It can therefore be inferred that early freshwater
15 inputs were sufficient to weaken AMOC, but further freshwater fluxes to the northern
16 North Atlantic (via the Atlantic Inflow), and extensive sea-ice formation associated
17 with widespread dispersal of freshwater across North Atlantic, were required to
18 strongly perturb North Atlantic climate and trigger significant changes in the Southern
19 Ocean, resulting in an abrupt increase in the release of CO₂ to the atmosphere.

20 The North Atlantic remained cool and the AMOC weak for at least a thousand
21 years after the end of the deposition of detrital carbonate during H1B (~16-17 ka).
22 Moreover, ventilation of the deep Atlantic was further reduced at ~16 ka (McManus
23 et al., 2004; Gherardi et al., 2005; Robinson et al., 2005; Thornalley et al., 2010b).
24 This later reduction has been attributed to freshwater associated with the second IRD
25 peak identified in the NE Atlantic (H1A, Bard et al., 2000), sourced from proximal

1 ice-sheets that weakened convection in the NE Atlantic (Gherardi et al., 2005),
2 although positive feedback processes associated with expanding sea-ice cover may
3 also have helped intensify climate deterioration and weakening of the AMOC. LIS
4 rerouting events after HS-1, but prior to the BA, may also have contributed towards a
5 weaker AMOC (Clark et al., 2001; Obbink et al., 2010; Thornalley et al., 2010a).
6 Whether the prolonged and intensified reduction of the AMOC during the late HS-1
7 necessitated a second freshwater injection from circum-NE Atlantic ice sheets, or if it
8 could have been maintained by sea-ice feedbacks alone, bears relevance upon what
9 controlled the timing of the climate amelioration at the end of HS-1, i.e. was the
10 primary control the abrupt cessation of freshwater input, as suggested by Liu et al.
11 (2009)?

12

13 **5.2.2 The onset of the Bølling-Allerød**

14 Numerous mechanisms have been called upon to explain the sudden strengthening of
15 the AMOC and North Atlantic warming at ~14.7 ka. For example: increased Aghulas
16 leakage of warm saline Indian Ocean water caused by southward migration of
17 Southern Ocean frontal systems in response to Antarctic sea-ice retreat (Knorr &
18 Lohman, 2003), freshening of AAIW caused by mwp-1a or retreat of Antarctic sea-
19 ice (Keeling & Stephens, 2001; Weaver et al., 2003), northward advection of salinity
20 that had accumulated in the tropics during the preceding interval of weakened AMOC
21 (e.g., Schmidt et al., 2004; Carlson et al., 2008a), or, an abrupt cessation or rerouting
22 of freshwater input to the North Atlantic (Clark et al., 2001; Liu et al., 2009; Obbink
23 et al., 2010). Several observations from the south of Iceland shed light upon possible
24 mechanisms responsible for the onset of the BA:

- 1 1. There was a gradual increase in the background salinity of the Atlantic Inflow
2 towards the onset of the BA, suggesting an increased salinity flux to the high
3 latitude North Atlantic. This flux would have then increased dramatically
4 during the abrupt resumption of strong AMOC because of the greater volume
5 of Atlantic Water that was drawn northwards (decreasing % *Nps* south of
6 Iceland).
- 7 2. Subsurface warming of local subpolar/polar waters, caused by surface ocean
8 stratification limiting atmospheric cooling of subsurface layers, and mixing
9 with warmer waters from the south, may have destabilised the water column
10 and triggered overturning, as suggested by modelling studies (e.g. Knorr &
11 Lohman, 2007).
- 12 3. Active sea-ice formation and the sinking of locally formed brines south of
13 Iceland between 16 ka and 14.8 ka (in addition to brine formation in the
14 Nordic Seas) will have drawn warm, saline Atlantic water northwards,
15 increasing local surface ocean salinity and triggering open ocean overturning
16 south of Iceland and in the Nordic Seas (Dokken & Jansen, 1999).
- 17 4. The abrupt strengthening of the AMOC at the onset of the BA coincides with
18 the end of active sea-ice formation south of Iceland, and the two processes are
19 probably coupled via a positive feedback (increasing AMOC causes sea-ice
20 retreat that allows further penetration of the Atlantic Inflow and invigoration
21 of the AMOC).

22
23 In order to simulate the abrupt warming of the BA, Liu et al. (2009) require
24 the continued input of freshwater to the North Atlantic (with a flux sufficient to
25 maintain the weakened AMOC state) until only a few centuries prior to the onset of

1 warming. We speculate that the necessary freshwater input was via H1A (~15-16 ka)
2 and the subsequent sea-ice formation in the NE Atlantic kept AMOC suppressed for
3 several further centuries, in a manner analogous to the prolonged sea-ice formation in
4 the Labrador Sea following the end of IRD deposition associated with H1B (Hillaire-
5 Marcel & deVernal, 2008). However, once the freshwater input had ended, it was
6 only a matter of time before the combined effects of subsurface warming and an
7 increasing salinity flux to the high latitudes overcame the effects of enhanced sea-ice
8 cover, leading to the reinvigoration of the AMOC, which coupled via a positive
9 feedback, caused rapid sea-ice retreat. Increasing atmospheric CO₂ levels may also
10 have helped melt northern North Atlantic sea-ice (Liu et al., 2009).

11

12 **5.2.3 Climate deterioration throughout the Bølling-Allerød and the onset of the** 13 **Younger Dryas**

14 To understand the relationship between North Atlantic freshwater input and climate
15 during the BA, it is instructive to consider the BA as a period of gradual climate
16 deterioration, accompanied by a reduction in deep ocean ventilation, upon which
17 centennial timescale cold intervals and sharp decreases in deep ocean ventilation are
18 superimposed. The centennial timescale cold intervals of the BA (i.e., the OD at
19 ~14.0 ka, a small event at ~13.6 ka, and the IACP at 13.4-13.0 ka) appear to have
20 been triggered by eastern rerouting of freshwater caused by the retreat of the LIS
21 during the preceding warm interval (Thornalley et al., 2010a). Based on the excellent
22 correlation between Nordic Seas IRD concentration and Greenland temperature,
23 circum-Nordic Seas ice-sheets presumably helped to amplify and/or sustain the
24 cooling in the Nordic Seas following the initial freshwater trigger delivered by the
25 Atlantic Inflow freshening.

1 What then determines the severity of the cold intervals and the ventilation of
2 the deep ocean south of Iceland during the BA? The magnitude of freshening of the
3 Atlantic Inflow is not the only consideration because the most prominent freshening
4 of the Atlantic Inflow during the BA, at ~14.2 ka, does not result in the coldest
5 conditions and poorest deep ocean ventilation of the BA – these events instead occur
6 during the IACP (13.0-13.4 ka). Knorr & Lohmann (2007) have hypothesised that the
7 extreme warmth of the Bølling was terminated by mwp-1a (or as previously reported
8 in Thornalley et al. (2010a), the eastward rerouting of the LIS contribution to mwp-1a
9 at ~14.2 ka), and the subsequent climate system shift towards cooler conditions may
10 have preconditioned the AMOC to be more sensitive to subsequent freshwater inputs.
11 Evidence from close to the LIS eastern outlets suggests background freshwater
12 discharge increased through these outlets during the BA (with rerouting events
13 superimposed), contributing to a gradually weakening AMOC (Obbink et al. 2010).
14 (Note - this background increase in freshwater discharge was apparently not advected,
15 via the NAC, as far north as the South Iceland Rise because no similar trend is
16 observed through the BA in our *G. inflata* $\delta^{18}\text{O}_{\text{sw-ivc}}$.)

17 Alternatively, Liu et al. (2009) have argued that the extreme warmth of the
18 Bølling was a transient event associated with an AMOC overshoot caused by basin-
19 wide salinity adjustments and the preceding North Atlantic subsurface warming, and
20 subsequent cooling may be an intrinsic relaxation of the climate system. Climate
21 cooling throughout the BA may also have facilitated increased ice-berg activity in the
22 Nordic Seas during centennial-scale cold intervals triggered by freshwater events in
23 the Atlantic Inflow. Considering Southern Ocean processes, the increasing sensitivity
24 of the AMOC to North Atlantic freshwater input through the BA may also have been
25 caused by a gradual increase in the salinity of AAIW, so that the density of AAIW

1 approached that of NADW, and any small freshwater input to the North Atlantic
2 caused NADW to be less dense than AAIW, resulting in a strong reduction in NADW
3 formation (Keeling & Stephens, 2001).

4 If the YD was indeed triggered by a freshwater input to the North Atlantic, the
5 preconditioning and gradual deterioration of the background climate throughout the
6 BA may explain why only a small freshwater input is detected south of Iceland at the
7 onset of the YD, sea-level shows little change (Bard et al., 2010), yet there is an
8 ensuing ~1200 year cold interval (Thornalley et al., 2010a). Of course, alternative
9 mechanisms for triggering the YD have been hypothesised (e.g., Tarasov & Peltier,
10 2005; Carlson et al., 2007a; Seager & Battisti, 2007; Clement & Peterson, 2008;
11 Eisenman et al., 2009).

12

13 **6. CONCLUSIONS**

14 Using multi-species planktic paired Mg/Ca- $\delta^{18}\text{O}$ measurements, the hydrography of
15 the Atlantic Inflow and local subpolar/polar water south of Iceland throughout the last
16 deglaciation has been reconstructed and compared to changes in deep ocean
17 circulation to elucidate the role freshwater input played in the climate evolution of
18 Termination I.

- 19 1. The strong reduction of the AMOC during the onset of HS-1 was presumably
20 related to the most extreme deglacial freshening of the Atlantic Inflow, caused
21 by a collapse of the LIS at ~16-17 ka. There was a further decrease in deep
22 ocean ventilation south of Iceland, coinciding with surface freshening and
23 possible sea-ice formation south of Iceland, that was likely caused by a later
24 collapse of a circum-NE Atlantic ice sheet (~15-16 ka).

- 1 2. Centennial-timescale cold intervals during the BA were caused by
2 fluctuations in the ice-margin of the LIS, which rerouted freshwater from
3 southern to eastern North American outlets. Iceberg discharge from circum-
4 Nordic Seas ice-sheets may have amplified or sustained cooling.
- 5 3. The gradual climate deterioration throughout the BA and into the YD,
6 combined with the absence of a strong freshwater signal south of Iceland at
7 the onset of the YD, suggests the YD may have been an intrinsic feature of
8 the deglaciation and only a relatively weak trigger was required once
9 background climate was preconditioned.
- 10 4. During the late HS-1 and YD, subsurface warming of local subpolar/polar
11 waters, brine formation in the northern North Atlantic, and a salty Atlantic
12 Inflow may all have helped promote the resumption of strong AMOC at the
13 onset of the BA and Holocene. The strengthening of AMOC during these
14 transitions is coupled with the retreat of sea-ice and a sharp increase in the
15 predominance of warm, saline Atlantic Inflow waters south of Iceland.
- 16 5. Freshwater input from the remnant LIS into the Labrador Sea likely caused
17 weak subpolar gyre circulation, resulting in a relatively warm and saline
18 Atlantic Inflow south of Iceland until ~8 ka. ISOW only attained its modern
19 strength and/or depth over the South Iceland Rise, and its characteristic,
20 stable, well-ventilated $\delta^{13}\text{C}$ signature by ~7-8 ka, possibly related to the upper
21 ocean reorganisation.

22

23 **7. ACKNOWLEDGEMENTS**

24 We thank the crew of CD-159; Linda Booth, Mervyn Greaves and Angela Huckle for
25 laboratory assistance; James Rolfe and Mike Hall for stable isotope analyses. We

1 would like to thank Anders Carlson and one anonymous reviewer for their positive
2 and helpful comments and guest editor Eric Wolff. Funding provided by NERC
3 RAPID grant NER/T/S/2002/00436.

4

5 **8. REFERENCES**

- 6 Alvarez-Solas, J., Charbit, S., Ritz, C., Paillard, D., Ramstein, G., Dumas, C., 2010.
7 Links between ocean temperature and iceberg discharge during Heinrich events.
8 *Nature Geoscience*, 3(2): 122-126.
- 9 Anand, P., Elderfield, H., Conte, M. H., 2003. Calibration of Mg/Ca thermometry in
10 planktonic foraminifera from a sediment trap time series. *Paleoceanography* 18:
11 10.1029/2002PA000846.
- 12 Andersen, R.F., Ali, S., Bradtmiller, L.I., Nielsen, S.H.H., Fleisher, M.Q., Anderson,
13 B.E., Burkle, L.H., 2009. Wind-driven upwelling in the Southern Ocean and the
14 deglacial rise in atmospheric CO₂. *Science*, 323: 1443-1448
- 15 Andrews, J.T. and Tedesco, K., 1992. Detrital carbonate rich sediments, Northwestern
16 Labrador Sea - Implications for ice-sheet dynamics and iceberg rafting (Heinrich)
17 events in the North Atlantic. *Geology*, 20(12): 1087-1090.
- 18 Bakke, J., Lie, Ø., Heegaard, E., Dokken, T., Haug, G., Birks, H., Dulski, P., Nilsen,
19 T., 2009. Rapid oceanic and atmospheric changes during the Younger Dryas cold
20 period. *Nature Geoscience*. 2: 202-205.
- 21 Barber, D.C., Dyke, A., Hillaire-Marcel, C., Jennings, A.E., Andrews, J.T., Kerwin,
22 M.W., Bilodeau, G., McNeely, R., Southon, J., Morehead, M.D., Gagnon, J.-M.,
23 1999. Forcing of the cold event of 8,200 years ago by catastrophic drainage of
24 Laurentide lakes. *Nature*, 400: 344-348.
- 25 Bard, E., Rostek, F., Turon, J. L., Gendreau, S., 2000. Hydrological impact of Heinrich
26 events in the subtropical northeast Atlantic. *Science* 289: 1321-1324.
- 27 Bard, E., Hamelin, B., Delanghe-Sabatier, D., 2010. Deglacial Meltwater Pulse 1B
28 and Younger Dryas Sea Levels revisited with boreholes at Tahiti. *Science*: doi:
29 10.1126/science.1180557.
- 30 Barker, S., Greaves, M. & H. Elderfield, H., 2003. A study of cleaning procedures
31 used for foraminiferal Mg/Ca paleothermometry. *Geochem. Geophys. Geosys.*, 4:
32 8407-8427.
- 33 Barker, S., Diz, P., Vautravers, M., Pike, J., Knorr, G., Hall, I.R., Broecker, W.S.,
34 2009. Interhemispheric Atlantic seesaw response during the last deglaciation.
35 *Nature*, 457: 1097-1102.
- 36 Bauch, D. and Bauch, H.A., 2001. Last glacial benthic foraminiferal delta O-18
37 anomalies in the polar North Atlantic: A modern analogue evaluation. *Journal of*
38 *Geophysical Research-Oceans*, 106(C5): 9135-9143.
- 39 Be, A.W.H., Tolderlund, D.S., 1971. Distribution and ecology of living planktonic
40 foraminifera in surface waters of the Atlantic and Indian Oceans, In: B.M.
41 Funnell, Riedel, W.R. (Eds.). *The micropaleontology of the oceans*, Cambridge,
42 United Kingdom, Cambridge University Press: 105-149.
- 43 Bemis, E. B., Spero, H.J., Bijma, J., Lea, D.W., 1998. Reevaluation of the oxygen
44 isotopic composition of planktonic foraminifera: Experimental results and revised
45 paleotemperature equation. *Paleoceanography*, 13: 150-160.

- 1 Bjorck, S. et al., 1996. Synchronized Terrestrial Atmospheric Deglacial Records
2 Around the North Atlantic. *Science*, 274(15 Nov.): 1155-1150.
- 3 Blindheim, J., Osterhus, S. (2005). The Nordic Seas, Main Oceanographic Features.
4 In: Drange, H. D., Dokken, T., Furevik, T., Gerdes, R., Berger, W. (Ed.), *The*
5 *Nordic Seas: An Integrated Perspective*. American Geophysical Union.
- 6 Bond, G. C., Lotti, R., 1995. Iceberg discharges into the North Atlantic on millennial
7 time scales during the last glaciation. *Science*, 267: 1005-1010.
- 8 Boyle, E.A. and Keigwin, L., 1987. North Atlantic thermohaline circulation during
9 the past 20,000 years linked to high-latitude surface temperature. *Nature*, 330: 35-
10 40.
- 11 Broecker, W. S., Kennett, J.P., Flower, B.P., Teller, J.T., Trumbore, S., Bonani, G.,
12 Wolfli, W., 1989. Routing of meltwater from the Laurentide ice sheet during the
13 Younger Dryas cold episode. *Nature*, 34: 318-321.
- 14 Carlson, A.E., Clark, P.U., Haley, B.A., Klinkhammer, G.P., Simmons, K., Brook,
15 E.J., Meissner, K.J., 2007a. Geochemical proxies of North American freshwater
16 routing during the Younger Dryas cold event. *Proc. Natl. Acad. Sci.*, 104: 6556-
17 6561.
- 18 Carlson, A.E., Clark, P.U., Raisbeck, G.M. and Brook, E.J., 2007b. Rapid Holocene
19 deglaciation of the Labrador sector of the Laurentide Ice Sheet. *Journal of*
20 *Climate*, 20: 5126-5133.
- 21 Carlson A.E., Oppo, D.W., Came, R.W., LeGrande, A.N., Keigwin, L.D., Curry,
22 W.B., 2008a. Subtropical Atlantic salinity variability and Atlantic meridional
23 circulation during the last deglaciation. *Geology*, 36: 991-994. Carlson, A.E.,
24 2008. Why there was not a Younger Dryas-like event during the Penultimate
25 Deglaciation. *Quaternary Science Reviews*, 27(9-10): 882-887.
- 26 Carlson, A.E. et al., 2008b. Rapid early Holocene deglaciation of the Laurentide ice
27 sheet. *Nature Geoscience*, 1(9): 620-624.
- 28 Carlson, A.E., and Clark, P.U., 2008. Rapid climate change and Arctic Ocean
29 freshening. *Comment: Geology*, doi: 10.1130/G24786C.1.
- 30 Carlson, A.E., Clark, P.U., Haley, B.A. and Klinkhammer, G.P., 2009. Routing of
31 western Canadian Plains runoff during the 8.2 ka cold event. *Geophysical*
32 *Research Letters*, 36.
- 33 Cheng, H., Edwards, R.L., Broecker, W.S., Denton, G.H., Kong, X., Wang, Y.,
34 Zhang, R., Wang, X., 2009. Ice Age Terminations. *Science* 326, 248-252.
- 35 Clement, A.C., Peterson, L.C., 2008. Mechanisms of abrupt climate change of the last
36 glacial period. *Rev. Geophys.*, 46: RG4002, doi:10.1029/2006RG000204.
- 37 Clark, P.U., Alley, R.B., Pollard, D., 1999. Northern Hemisphere ice-sheet influences
38 on global climate change. *Science*, 286: 1104-1111.
- 39 Clark, P.U., Marshall, S.J., Clarker, G.K.C., Hostetler, S.W., Licciardi, J.M., Teller,
40 J.T., 2001. Freshwater forcing of abrupt climate change during the Last
41 Glaciation. *Science*, 293: 283-287.
- 42 Clark, P.U., McCabe, A.M., Mix, A.C., Weaver, A.J., 2004. Rapid rise of sea level
43 19,000 years ago and its global implications. *Science*, 304: 1141-1144.
- 44 Clark, P.U., Hostetler, S.W., Pisias, N.G., Schmittner, A., and Meissner, K.J., 2007.
45 Mechanisms for a ~7-kyr climate and sea-level oscillation during marine isotope
46 stage 3, in Schmittner, A., Chiang, J., and Hemming, S., Eds., *Ocean Circulation:*
47 *Mechanisms and Impacts*, American Geophysical Union, *Geophysical Monograph*
48 173, Washington, D.C., pp. 209-246.
- 49 Clark, P.U., Dyke, A.S., Shakun, J.D., Carlson, A.E., Clark, J., Wohlfarth, B.,
50 Mitrovica, J.X., Hostetler, S.W., McCabe, A.M., 2009. The Last Glacial

1 Maximum. *Science*, 325: 710-714.

2 Cléroux, C., Cortijo, E., Duplessy, J.-C., Zahn, R., 2007. Deep-dwelling foraminifera
3 as thermocline temperature recorders. *Geochem. Geophys. Geosys.*, 8:
4 doi:10.1029/2006GC001474.

5 deMenocal, P., Ortiz, J., Guilderson, T., Sarnthein, M., 2000. Coherent high- and low-
6 latitude climate variability during the Holocene Warm Period. *Science* 288: 2198-
7 2202.

8 deVernal, A., Hillaire-Marcel, C. & Bilodeau, G., 1996. Reduced meltwater outflow
9 from the Laurentide ice margin during the Younger Dryas. *Nature*, 381: 774-777.

10 Dokken, T., Jansen, E., 1999. Rapid changes in the mechanism of ocean convection
11 during the last glacial period. *Nature*, 401: 458-461.

12 Eisenman, I., Bitz, C.M., Tziperman, E., 2009. Rain driven by receding ice sheets as a
13 cause of past climate change. *Paleoceanography*, 24: PA4209,
14 doi:10.1029/2009PA001778.

15 Elderfield, H., Ganssen, G., 2000. Past temperature and $\delta^{18}\text{O}$ of surface ocean waters
16 inferred from foraminiferal Mg/Ca ratios. *Nature*, 405: 442-445.

17 Elliot, M., Labeyrie, L., Bond, G., Cortijo, E., Turon, J.-L., Tisnerat, N., Duplessy,
18 J.-C., 1998. Millennial-scale iceberg discharges in the Irminger Basin during the
19 last glacial period: Relationship with the Heinrich events and environmental
20 settings. *Paleoceanography*, 13: 433-446.

21 Ellison, C.R.W., Chapman, M.R., Hall, I.R., 2006. Surface and deep ocean
22 interactions during the cold climate event 8,200 years ago. *Science*, 312: 1929-
23 1932.

24 Fairbanks, R.G., 1989. A 17,000 year glacio-eustatic sea level record: influence of
25 glacial melting rates on the Younger Dryas event and deep ocean circulation.
26 *Nature*, 342: 637-642.

27 Fisher, T.G., Smith, D.G., Andrews, J.T., 2002. Preboreal oscillation caused by a
28 glacial Lake Agassiz flood. *Quat. Sci. Rev.* 21: 873-878.

29 Ganssen, G.M., Kroon, D., 2000. The isotopic signature of planktonic foraminifera
30 from NE Atlantic surface sediments: implications for the reconstruction of past
31 oceanic conditions. *Journal of the Geological Society London*, 157: 693-699.

32 Gherardi, J.-M., Labeyrie, L., McManus, J.F., Francois, R., Skinner, L.C., Cortijo, E.,
33 2005. Evidence from the Northeastern Atlantic basin for the variability in the rate
34 of meridional overturning circulation through the last deglaciation. *Earth and
35 Planetary Science Letters*, 240: 710-723.

36 Hall, I.R., Moran, S.B., Zahn, R., Knutz, P.C., Shen, C.-C., Edwards, R.L., 2006.
37 Accelerated drawdown of meridional overturning in the late-glacial Atlantic
38 triggered by pre-H event freshwater perturbation. *Geophysical Research Letters*,
39 33: L16616, doi:10.1029/2006GL026239.

40 Hathorne, E.C., James, R.H., Lampitt, R.S., 2009. Environmental versus
41 biomineralization controls on the intratest variation in the trace element
42 composition of the planktonic foraminifera *G. inflata* and *G. scitula*.
43 *Paleoceanography*, 24: PA4204, doi:10.1029/2009PA001742.

44 Hatun, H., Sando, A. B., Drange, H., Hansen, B., Valdimarsson, H., 2005. Influence
45 of the Atlantic subpolar gyre on the thermohaline circulation. *Science*, 309: 1841-
46 1844.

47 Hemming, S.R., 2004. Heinrich Events: Massive Late Pleistocene detritus layers of
48 the North Atlantic and their global climate imprint. *Rev. Geophys.*, 42:
49 doi:10.1029/2003RG000128.

- 1 Hilbrecht, H., 1996. Extant planktic foraminifera and the physical environment in the
2 Atlantic and Indian Oceans.- Mitteilungen aus dem Geologischen Institut der
3 Eidgen. Technischen Hochschule und der Universität Zürich, Neue Folge. No.
4 300, 93 pp.; Zürich
- 5 Hillaire-Marcel, C., deVernal, A., Bilodeau, G., Weaver, A., 2001. Absence of deep-
6 water formation in the Labrador Sea during the last interglacial period. *Nature*,
7 410: 1073-1077.
- 8 Hillaire-Marcel, C., deVernal, A., 2008. Stable isotope clue to episodic sea ice
9 formation in the glacial North Atlantic. *Earth and Planetary Science Letters*, 268:
10 143-150.
- 11 Jouzel, J. et al., 2007. Orbital and Millennial Antarctic Climate Variability over the
12 past 800,000 years. *Science*, 317: 793-796.
- 13 Keeling, R.F., Stephens, B.B., 2001. Antarctic sea ice and the control of Pleistocene
14 climate instability. *Paleoceanography*, 16: 112-131.
- 15 Keigwin, L.D., Jones, G.A., Lehman, S.J., Boyle, E.A., 1991. Deglacial meltwater
16 discharge, North Atlantic deep circulation, and abrupt climate change. *Journal of*
17 *Geophysical Research*, 96 (C9): 16811-16826.
- 18 Kim, S.-T., O'Neil, J.R., 1997. Equilibrium and non-equilibrium oxygen isotope
19 effects in synthetic carbonates. *Geochim. Cosmochim. Acta*, 61: 3461– 3475.
- 20 Kleiven, H. F., Kissel, C., Laj, C., Ninnemann, U. S., Richter, T. O., et al., 2008.
21 Reduced North Atlantic Deep Water coeval with the glacial Lake Agassiz
22 freshwater outburst. *Science*, 319: 60-64.
- 23 Knorr, G., Lohmann, G., 2003. Southern Ocean origin for the resumption of Atlantic
24 thermohaline circulation during deglaciation. *Nature*, 424: 532-536.
- 25 Knorr, G., Lohmann, G., 2007. Rapid transitions in the Atlantic thermohaline
26 circulation triggered by global warming and meltwater during the last
27 deglaciation. *Geochemistry Geophysics Geosystems*, 8: doi:
28 10.1029/2007GC001604.
- 29 Kozdon, R., Eisenhauer, A., Weinelt, M., Meland, M.Y., Nürnberg, D., 2009a.
30 Reassessing Mg/Ca temperature calibrations of *Neogloboquadrina pachyderma*
31 (sinistral) using paired $\delta^{44/40}\text{Ca}$ and Mg/Ca measurements. *Geochem. Geophys.*
32 *Geosyst.*, 10: Q03005, doi:10.1029/2008GC002169.
- 33 Kozdon, R. et al. (2009b), Intratest oxygen isotope variability in the planktonic
34 foraminifer *N. pachyderma*: Real vs. apparent vital effects by ion microprobe,
35 *Chemical Geology*, 258, 327-337.
- 36 Jonkers, L., Brummer, G.J.A., Peeters, F.J.C., van Aken, H.M. and De Jong, M.F.,
37 2010. Seasonal stratification, shell flux, and oxygen isotope dynamics of left-
38 coiling *N. pachyderma* and *T. quinqueloba* in the western subpolar North Atlantic.
39 *Paleoceanography*, 25.
- 40 Labeyrie, L., Duplessey, J.-C., Duprat, J., Juillet-Leclerc, A., Moyes, J., Michel, E.,
41 Kallel, N., Shackleton, N.J., 1992. Changes in the vertical structure of the North
42 Atlantic Ocean between glacial and modern times. *Quaternary Science Reviews*,
43 11: 401-414.
- 44 Lea, D.W., 2003. Elemental and isotopic proxies of past ocean temperatures. In:
45 Elderfield, H. (Ed.), *Treatise on Geochemistry*, v6: 366-390.
- 46 Lehman, S.J., Keigwin, L.D., 1992. Sudden changes in North Atlantic circulation
47 during the last deglaciation. *Nature*, 356: 757-762.
- 48 Liu, Z. et al., 2009. Transient simulation of last deglaciation with a new mechanism
49 for Bølling- Allerød warming. *Science*, 325: 310-314.
- 50 Lüthi, D. et al., 2008. High-resolution carbon dioxide concentration record 650,000-

1 800,000 years before present. *Nature*, 453: 379-382.

2 Lynch-Stieglitz, J. et al., 2007. Atlantic meridional overturning circulation during the
3 Last Glacial Maximum. *Science*, 316: 66-69.

4 Marchitto, T. M., Curry, W.B., Oppo, D.W., 1998. Millennial-scale changes in North
5 Atlantic circulation since the last glaciation. *Nature*, 393: 557-561.

6 McCave, I.N., 2005. Cruise Report RRS Charles Darwin 159 – RAPID Climate
7 Change, University of Cambridge, Cambridge. 49 pp.

8 McManus, J. F., Francois, R., Gherardi, J.-M., Keigwin, L.D., Brown-Leger, S., 2004.
9 Collapse and rapid resumption of Atlantic meridional circulation linked to
10 deglacial climate changes. *Nature*, 428: 834-837.

11 Meissner, K.J. and Clark, P.U., 2006. Impact of floods versus routing events on the
12 thermohaline circulation. *Geophysical Research Letters*, 33(15).

13 Meland, M. Y., Dokken, T.M., Jansen, E., Hevroy, K., 2008. Water mass properties
14 and exchange between the Nordic seas and the northern North Atlantic during the
15 period 23-6 ka: Benthic oxygen isotopic evidence. *Paleoceanography*, 23: doi:
16 10.1029/2007PA001416.

17 Menot, G. et al., 2006. Early reactivation of European rivers during the last
18 deglaciation. *Science*, 313(5793): 1623-1625.

19 Murton, J.B., Bateman, M.D., Dallimore, S.R., Teller, J.T. and Yang, Z.R., 2010.
20 Identification of Younger Dryas outburst flood path from Lake Agassiz to the
21 Arctic Ocean. *Nature*, 464(7289): 740-743.

22 NGRIP members, 2004. High-resolution record of Northern Hemisphere climate
23 extending into the last interglacial period. *Nature*, 431: 147-151.

24 Nyland, B. F., Jansen, E., Elderfield, H., Andersson, C., 2006. *Neogloboquadrina*
25 *pachyderma* (dex. and sin.) Mg/Ca and delta O-18 records from the Norwegian
26 Sea. *Geochemistry Geophysics Geosystems* 7: doi: 10.1029/2005GC001055.

27 Obbink, E.A., Carlson, A.E. and Klinkhammer, G.P., 2010. Eastern North American
28 freshwater discharge during the Bolling-Allerod warm periods. *Geology*, 38(2):
29 171-174.

30 Oppo, D.W., Lehman, S.J., 1993. Mid-depth circulation of the subpolar North
31 Atlantic during the Last Glacial Maximum. *Science*, 259: 1148-1152.

32 Oppo, D.W., Horowitz, M. and Lehman, S.J., 1997. Marine core evidence for reduced
33 deep water production during Termination II followed by a relatively stable
34 substage 5e (Eemian). *Paleoceanography*, 12(1): 51-63.

35 Oppo, D.W., McManus, J.E., Cullen, J.L., 2003. Deepwater variability in the
36 Holocene epoch. *Nature*, 422: 277-278.

37 Peck, V.L., Hall, I.R., Zahn, R., Grousset, F., Hemming, S.R., Scourse, J.D., 2007.
38 The relationship of Heinrich events and their European precursors over the past 60 ka
39 BP: a multi-proxy ice-rafted debris provenance study in the North East Atlantic.
40 *Quat. Sci. Rev.*, 26:862-875.

41 Peltier, W.R., deVernal, A., Hillaire-Marcel, C., 2008. Reply to comments by Carlson
42 and Clark on “Rapid climate change and Arctic Ocean freshening”. *Geology*, doi:
43 10.1130/G24971Y.1

44 Pflaumann, U., Sarnthein, M., Chapman, M., d'Abreu, L., Funnell, B., Huels, M.,
45 Kiefer, T., Maslin, M., Schulz, H., Swallow, J., van Kreveld, S., Vautravers, M.,
46 Vodelsang, E. & Weinelt, M., 2003. Glacial North Atlantic: Sea-surface
47 conditions reconstructed by GLAMAP 2000. *Paleoceanography*, 18: doi:
48 10.1029/2002PA000774.

49 Piotrowski, A. M., Goldstein, S.L., Hemming, S.R., Fairbanks, R.G., 2004.
50 Intensification and variability of ocean thermohaline circulation through the last

- 1 deglaciation. *Earth and Planetary Science Letters*, 225: 205-220.
- 2 Rasmussen, T.L., Thomsen, E., 2004. The role of the North Atlantic Drift in the
3 millennial timescale glacial climate fluctuations. *Palaeogeography*
4 *Palaeoclimatology Palaeoecology*, 210: 101-116.
- 5 Rasmussen, T.L. and Thomsen, E., 2009a. Stable isotope signals from brines in the
6 Barents Sea: Implications for brine formation during the last glaciation. *Geology*,
7 37(10): 903-906.
- 8 Rasmussen, T.L. and Thomsen, E., 2009b. Ventilation changes in intermediate water
9 on millennial time scales in the SE Nordic seas, 65-14 kyr BP. *Geophysical*
10 *Research Letters*, 36(1).
- 11 Renssen, H. Seppa, H., Heiri, O., Roche, D.M., Goose, H., Fichet, T., 2009. The
12 spatial and temporal complexity of the Holocene thermal maximum. *Nature*
13 *Geoscience*, 2: 411-414.
- 14 Rickaby, R.E.M., Elderfield, H., 2005. Evidence from the high-latitude North Atlantic
15 for variations in Antarctic Intermediate water flow during the last deglaciation.
16 *Geochemistry Geophysics Geosystems*, 6: doi: 10.1029/2004GC000858.
- 17 Rinterknecht, V.R. et al., 2006. The last deglaciation of the southeastern sector of the
18 Scandinavian Ice Sheet. *Science*, 311(5766): 1449-1452.
- 19 Robinson, L.F., Adkins, J.F., Keigwin, L.D., Southon, J., Fernandez, D.P., Wang, S.-
20 L., Schrier, D.S., 2005. Radiocarbon variability in the western North Atlantic
21 during the last deglaciation. *Science*, 310: 1469-1473.
- 22 Ruddiman, W.F., Molino, B., Esmay, A. and Pokras, E., 1980. Evidence bearing on
23 the mechanism of rapid deglaciation. *Climatic Change*, 3(1): 65-87.
- 24 Sarnthein, M. et al. (2000), *Fundamental Modes and Abrupt Changes in North*
25 *Atlantic Circulation and Climate over the last 60 ky - Concepts, Reconstruction*
26 *and Numerical Modeling*. In: Schafer, P. W. Ritzrau, W. Schluter, J. Thiede (Ed.),
27 *The Northern North Atlantic: A Changing Environment*. Springer, Berlin.
- 28 Seager, R., Battisti, D.S., 2007. Challenges to our understanding of the general
29 circulation: abrupt climate change. In: Schneider and Sobel (Eds.), *Global*
30 *Circulation of the Atmosphere*. Princeton Univ. Press, pp. 331-371.
- 31 Schmidt, M.W., Spero, H.J., Lea, D.W., 2004. Links between salinity variation in the
32 Caribbean and North Atlantic thermohaline circulation. *Nature*, 428: 160-163.
- 33 Shaffer, G., Olsen, S.M. and Bjerrum, C.J., 2004. Ocean subsurface warming as a
34 mechanism for coupling Dansgaard-Oeschger climate cycles and ice-rafting
35 events. *Geophysical Research Letters*, 31(24).
- 36 Siddall, M., Rohling, E.J., Almogi-Labin, A., Hemleben, Ch., Meischner, D.,
37 Schmeizer, I., Smeed, D.A., 2003. Sea-level fluctuations during the last glacial
38 cycle. *Nature*, 423: 853-858.
- 39 Simstich, J., Sarnthein, M., Erlenkeuser, H., 2003. Paired $\delta^{18}\text{O}$ signals of
40 *Neogloboquadrina pachyderma* (s) and *Turborotalita quinqueloba* show thermal
41 stratification structure in Nordic Seas. *Marine Micropalaeontology*, 48: 107-125.
- 42 Smith, L. M., Andrews, J. T., Castaneda, I. S., Kristjansdottir, G. B., Jennings, A. E.,
43 et al., 2005. Temperature reconstructions for SW and N Iceland waters over the
44 last 10 cal ka based on $\delta^{18}\text{O}$ records from Planktic and Benthic Foraminifera.
45 *Quaternary Science Reviews*, 24: 1723-1740.
- 46 Solignac, S., de Vernal, A., Hillaire-Marcel, C., 2004. Holocene sea-surface
47 conditions in the North Atlantic - contrasted trends and regimes in the western and
48 eastern sectors (Labrador Sea vs. Iceland Basin). *Quaternary Science Reviews*, 23:
49 319-334.
- 50 Stangeew, E., 2001. *Distribution and Isotopic Composition of Living Planktonic*

- 1 Forminifera *N. pachyderma* (sinistral) and *T. quiqueloba* in High Latitude North
2 Atlantic. Dr. Scient., der Christian-Albrechts-Universität zu Kiel.
- 3 Stocker, T. F., Johnsen, S.J., 2003. A minimum thermodynamic model for the bipolar
4 seesaw. *Paleoceanography*, 18(4): 1087, doi:10.1029/2003PA000920.
- 5 Tarasov, L., Peltier, W.R., 2005. Arctic freshwater forcing of the Younger Dryas cold
6 reversal. *Nature*, 435: 662-665.
- 7 Teller, J.T., Leverington, D.W., Mann, J.D., 2002. Freshwater outbursts to the oceans
8 from glacial Lake Agassiz and their role in climate change during the last
9 deglaciation. *Quat. Sci. Rev.*, 21: 879-887.
- 10 Thornalley, D.J.R., Elderfield, H., McCave, I.N., 2009. Holocene oscillations in
11 temperature and salinity of the subpolar North Atlantic. *Nature*, 457: 711-714.
- 12 Thornalley, D. J. R., McCave, I.N., Elderfield, H., 2010a. Freshwater input and abrupt
13 deglacial climate change in the North Atlantic. *Paleoceanography*, 25: PA1201,
14 doi:10.1029/2009PA001772.
- 15 Thornalley, D.J.R., Elderfield, H., McCave, I.N., 2010b. Intermediate and deep water
16 paleoceanography of the northern North Atlantic over the past 21,000 years.
17 *Paleoceanography*, 25: PA1211, doi:10.1029/2009PA001833.
- 18 Toggweiler, J.R., Russell, J.L., Carson, S.R., 2006. Midlatitude westerlies,
19 atmospheric CO₂, and climate change during the ice ages. *Paleoceanography*, 21:
20 PA2005, doi:10.1029/2005PA001154.
- 21 Waelbroeck, C., Labeyrie, L., Duplessy, J. C., Guiot, J., Labracherie, M., Leclaire, H.,
22 Duprat, J., 1998. Improving past sea surface temperature estimates based on
23 planktonic fossil faunas. *Paleoceanography*, 13: 272-283.
- 24 Waelbroeck, C., Labeyrie, L., Michel, M., Duplessy, J.C., McManus, J.F., Lambeck
25 K., Balbon, E. & Labracherie, M., 2002. Sea-level and deep water temperature
26 changes derived from benthic foraminifera isotopic records. *Quat. Sci. Rev.*, 21:
27 295-305.
- 28 Waelbroeck, C., Duplessy, J.-C., Michel, E., Labeyrie, L., Paillard, D., Duprat, J.,
29 2001. The timing of the last deglaciation in North Atlantic climate records.
30 *Nature*, 412: 724-727.
- 31 Weaver, A.J., Saenko, O.A., Clark, P.U., Mitrovica, J.X., 2003. Meltwater pulse 1a
32 from Antarctica as a trigger of the Bølling-Allerød warm interval. *Science*, 299:
33 1709-1713.

1 **9. FIGURE LEGENDS, TABLES**

2 **Fig. 1** Modern northern North Atlantic ocean circulation and the location of the
3 study area (black star). Grey arrows indicate deep ocean currents; solid black arrows,
4 warm surface currents; dashed black arrows, cold surface currents. SPG, subpolar
5 gyre; NAC, North Atlantic Current; EGC, East Greenland Current; EIC, East
6 Icelandic Current.

7 **Fig. 2.** Oxygen isotope data from cores RAPID-15-4P (grey) and RAPID-17-5P
8 (black). **(a)** *G. bulloides* **(b)** *C. wuellerstorfi* (Thornalley et al., 2010b).

9 **Fig. 3.** Paired $\delta^{18}\text{O}$ -Mg/Ca data for *N. pachyderma* (s) from RAPID-10-1P (grey)
10 and RAPID-15-4P (black). **(a)** % abundance of *Nps*. **(b)** *Nps* $\delta^{18}\text{O}$, Thornalley et al.,
11 2010b **(c)** *Nps* Mg/Ca ratios **(d)** 'Ice-volume corrected' seawater $\delta^{18}\text{O}$.

12 **Fig. 4.** Bar chart illustrating the range of reconstructed Mg/Ca-temperatures for the
13 different planktic species analysed.

14 **Fig. 5.** Comparison of surface and deep ocean records. Light grey bars, Greenland
15 cold intervals; dark grey bars, Heinrich events H-0 and H1. **(a)** Records of ice-rafted
16 debris (IRD) from south of Iceland (black line, 150-250 μm fraction, excluding
17 volcanic grains, from RAPID-17-5P, Thornalley et al., 2010b) and the Nordic Seas
18 (grey line, ratio of lithics:(lithics+foraminifera), from core MD99-2284, Bakke et al.,
19 2009). **(b)** Derived temperatures and 'ice-volume corrected' seawater $\delta^{18}\text{O}$ from
20 planktic paired Mg/Ca- $\delta^{18}\text{O}$ data. Data from 7-11.4 ka is from RAPID-12-1K
21 (Thornalley et al., 2009), data from 11.4-18 ka is from RAPID-15-4P (Thornalley et
22 al., 2010a). *G. bulloides*, open symbols, *G. inflata*, filled diamonds. **(c)** Derived
23 temperatures and 'ice-volume corrected' seawater $\delta^{18}\text{O}$ from *Nps* paired Mg/Ca- $\delta^{18}\text{O}$
24 data. All data is from RAPID-15-4P. **(d)** Benthic $\delta^{13}\text{C}$ and 'ice-volume corrected'


1 $\delta^{18}\text{O}$ for *C. wuellerstorfi* (>212 μm) from RAPiD-15-4P (2133 m), Thornalley et al.,
2 2010b. (e) NGRIP $\delta^{18}\text{O}$ (NGRIP members, 2004).

3 **Fig. 6.** Proxy records indicating changes in the strength of the subpolar gyre (SPG) –
4 increasing SPG strength downwards. (a) ‘Ice-volume corrected’ seawater $\delta^{18}\text{O}$ from
5 RAPID-12-1K (0-11.4 ka) and RAPiD-15-4P (11.4-18 ka), inferred salinity increases
6 upwards. (b) Labrador Sea salinity, based on dinocyst assemblage data from core P-
7 013 (Solignac et al., 2004) – salinity increases downwards. (c) Linear detrended
8 temperature anomalies based on planktic foraminiferal assemblages, from the eastern
9 subtropical Atlantic, ODP site 658C (deMenocal et al., 2000).

1 **Table 1.** Synthesis of deglacial palaeoceanography south of Iceland. (f.w., freshwater; SSW, Southern Source Water).


	Events south of Iceland				Controls elsewhere	
	% <i>Nps</i> - (sub)polar water	Atlantic Inflow	Local subpolar/polar water	Deep ocean (~2 km)		
Mid-Late Holo. (0-9 ka)	<1%	Oscillating <i>TS</i> changes – SPG control Millennial surface f.w. events	-	Modern, stable ISOW $\delta^{13}\text{C}$ and flow strength/depth by ~7-8 ka	Oscillating SPG strength. Atmospheric circulation changes.	
8.2 ka event	<1%	High background salinity caused by weak SPG	0.5 psu f.w. event at 8.4-8.2 ka	-	Poor ventilation ($\delta^{13}\text{C}$ ~0.5‰)	Drainage of Lake Agassiz
Preboreal (~9-11.7 ka)	60% to <1% (50% in PBO)		Warm and salty, gradually decreasing.	Warm. f.w. input from local melting Salty during cold interval	Variable $\delta^{13}\text{C}$ – preformed or entrained signal, or increased SSW. Brines ~10.5-11.7 ka	f.w. event at 11.3 ka – LIS source via Arctic (?)
YD (11.7-12.9 ka)	85-90%		Warm and salty, 100-yr f.w. event at onset	Warming through YD f.w. or brine formation with high IRD at 12.1-11.7 ka	Poor ventilation at 12.8-12.6 ka and 12.1-11.7 ka. Brines at 12.1-11.7 ka.	LIS f.w. event at onset (?) H-0 ~12.1-11.7 ka. Abundant IRD in Nordic Seas.
BA (12.9-14.7 ka)	Fluctuating 20-80%		f.w. events caused by LIS rerouting, trigger cold intervals	Warmer and fresher (enhanced melting) during climate warm intervals	Well ventilated, decreasing through BA. Poorer ventilation in cold intervals.	Open ocean convection Nordic Seas IRD in cold intervals Fluctuating LIS margin
HS-1 (14.7-17 ka)	85-95%		Warm. f.w. from LIS 17.5-16 ka	Sub-surface warming to 5°C. Active sea-ice formation from 16-14.8 ka.	Decreasing ventilation. Increasing brine influence	H1B (~17 ka) – LIS IRD H1A (~16 ka) – NE Atlantic IRD Heat/salinity retention in (sub)tropics
LGM and early deglacial (17-21 ka)	85-95%		-	Cold (3-4.5 °C). No signal of f.w. events or sea-ice formation	GNAIW and GAABW, boundary at 2-2.5 km.	Nordic Seas sea-ice and ice-bergs exported to south of Iceland. 19 ka melt-water pulse

1 **Fig. 1**




2

1 Fig.2




2

1 **Fig. 3**




2

1 **Fig. 4**




2

1 **Fig. 5**



2

1 **Fig. 6**



2