

Diagnostic First and Last Occurrences of Mesozoic and Cenozoic Planktonic Foraminifera

MARCELLE K. BOUDAGHER-FADEL

OVPR, UCL, Gower Street, London WC1E 6BT, UK

Email: m.fadel@ucl.ac.uk

Abstract: The diagnostic first and last occurrences of Cretaceous, Paleogene and Neogene planktonic foraminifera species are given, and have been calibrated against the most recent biostratigraphical time scale and radio-isotope data.

The major planktonic foraminiferal bioevents for the Late Mesozoic and the Cenozoic (as defined by BouDagher-Fadel, 2012) are calibrated against the biostratigraphical time scale and the radio-isotopes (as defined by Gradstein et al., 2012). Those of the Cenozoic are compared with the earlier zonations of Berggren and Pearson (2005) and Wade et al. (2011). The biostratigraphic resolution of the alpha-numeric biozonations is determined by distinct evolutionary lineages of morphospecies and has been deduced through extensive industrial work, calibrated against Sr-isotopes and other biostratigraphic markers, involving global offshore wells by the present author and compared to already published works (for further details see BouDagher-Fadel, 2012).

Three figures are presented. Fig. 1 shows the diagnostic calibration based on species of planktonic foraminifera for the Cretaceous biozonations as defined by BouDagher-Fadel (2012). First occurrences and last appearances of index foraminiferal taxa are plotted against the biostratigraphical time scale of Gradstein *et al.* (2012).

Figs 2 and 3 shows respectively the diagnostic calibration of planktonic foraminifera of the Paleogene and Neogene biozonations as defined by BouDagher-Fadel (2012). First

occurrences and last appearances of index foraminiferal taxa are plotted against the biostratigraphical time scale of Gradstein *et al.* (2012) and compared against the earlier zonations of Berggren and Pearson (2005) and Wade *et al.* (2011).

References

Berggren, W.A., Pearson, P.N., 2005. A revised tropical and subtropical Paleogene planktonic foraminiferal zonation. *J. Foraminiferal Res.*, **35**, 279–298.

BouDagher-Fadel, M.K., 2012. Biostratigraphic and Geological Significance of Planktonic Foraminifera. Developments in Palaeontology and Stratigraphy: Vol.22. Elsevier: Amsterdam, 312p.

Gradstein, F. M., Ogg, J. G., Schmitz, M. D., Ogg, G. M., 2012. *The Geologic Time Scale* 2012. Elsevier, 1,144 p.

Wade, B.S., Pearson, P.N., Berggren, W.A., Paelike, H., 2011. Review and revision of Cenozoic tropical planktonic foraminiferal biostratigraphy and calibration to the geomagnetic polarity and astronomical time scale. *Earth Sci. Rev.*, **104**, 111-142.

Age (Ma)	Age (Ma) Period, Epoch,			Planktonic zonation		Diagnostic First Occurrence of Planktonic Foraminifera		Diagnostic Last Occurrence of Planktonic Foraminifera
99			\dashv	2 b	66.2	Ventilabrella multicamerata	4 66.0 1	Globotruncana spp./Rugoglobigerina spp.
99			نډ	3 a	67.0	Plummerita reicheli	4	Contusotruncana plicata
			Maast.	2	70.0	Racemiguembelina powelli		Rugotruncana subpennyi
			Σ	1 b	71.0	Contusotruncana plicata Plummerita hantkeninoides	70.0	Globotruncana lapparenti
			\dashv	b	72.1	!		Radotruncana subspinosa
				3 <u>a</u>	74.5	Rugotruncana subpennyi Globotruncanita stuarti		Globotruncanita atlantica
		Φ	Camp.	2	₹ 77.0		77.0	Radotruncana calcarata
		Late	ပိ	1	₹ 80.3	Radotruncana calcarata, Globotruncanita atlantica Globotruncana mariei	₹ 80.3	Rugoglobigerina subrugosa
				2	₹ 83.6		83.6	Concavatotruncana asymetrica
			Sant.		84.9 85.6	Rugoglobigerina subrugosa Globotruncana linneiana		
			S	1 a	86.3	Concavatotruncana asymetrica	86.3	Marginotruncana paraconcavata
			Con.	2	₹ 88.0	Globotruncana lapparenti	00.5	
			0	1	89.8	Marginotruncana paraconcavata	<u> </u>	Circultura and Alice and Circultural
			ن	3	91.3	Dicarinella primitiva	89.8 91.3	Sigalitruncana biconvexiformis Helvetoglobotruncana helveticae
	Cretaceous Cen. Tur.		Tu	1	92.8	l Helvetoglobotruncana helveticae Sigalitruncana biconvexiformis	4	Concavatotruncana elata
				3 5	93.9 95.1	Rotalipora cushmani	▼ 93.9 •	Rotalipora cushmani
			Sen	2 b	97.2 98.8	Thalmanninella reicheli Thalmanninella greenhornensis	33.1	Favusella washitensis
LO	rets		$\overset{\smile}{-}$	1 b	100.5 101.5	Thalmanninella globotruncanoides Thalmanninella gandolfi	99.6 100.5	Alanlordella bentonensis Hedbergella rischi
100.5	Ō		H	4 a	102.2 104.4	Thalmanninella gandolfi Ticinella madecassiana Pseudothalmanninella subticinensis	102.2	Ticinella praeticinensis
_			Alb.	3 a	106.7	Ticinella praeticinensis	4	
			1	2	109.8	Ticinella roberti Ticinella primula	106.7	Blefuscuiana aptiana
			-	1 a	113.0	Hedbergella rischi	4 113.0	B. maslakovae/ L. globulifera
				4 a	115.0 116.5	l Blefuscuiana kuznetsovae Lilliputianella kuhryi		
			Apt.	3	119.5	Praehedbergella tatianae	L	
			Ā	2	122.9		119.5	Praehedbergella ruka
		arly		1		Blefuscuiana maslakovae	122.9	Gorbachikella anteroapertura
		Ear		3	127.0	ј Бјејиѕсијана таѕјакоуае 	126.3	Gorbachikella grandiapertura
			Barr.	2	129.0	Blefuscuiana daminiae		
			-	1	130.8	B. aptiana/Pr. tuschepsensis	129.0	Praehedbergella tuschepsensis
			Haut.	2		Gorbachikella anteroapertura	4 122 0 □	0 - 1 - 11 11 - 1 1
			_	1	_	Praehedbergella handousi	133.9	Praehedbergella handousi
			Val.	2	◀ 136.6	Gorbachikella grandiapertura	136.65	Conoglobigerina gulekhensis
				1	-		139.4	Conoglobigerina sp. A
5			Berr.				1 233.4	Conograpiyerina sp. A
14	_		\blacksquare		145.0	Conoglobigerina gulekhensis	145.0	Conoglobigerina conica
Ju	Jurassic				152.1	Conoglobigerina conica	143.0	conography control

Fig. 1. Diagnostic First and Last Occurrences of Cretaceous Planktonic Foraminifera.

Diagnostic Last Occurrence of Planktonic Foraminifera	7≥6 □ Globigerina sellii 7≥8 □ Tenuirella gemma/Paragloborotalia opima		28.1 I Globoturborotalita angulioffinalis				339 Hantkenina alabamensis	Marazovella snimulosa/Truncoratalnidas trodiensis	38.0	* 38.2 Orbulinoides beckmanni/Monozovalla lebneri	_	412 Globigerapsis kugleri/Guembelitrioides higginsi	◆ 42.3		44.9 Acarinina decepta		47.0 Morozovella caucasica			Anadarina milanonaria	510 - Acanima wincaxansis		54.0 Morozovella occlusa		se.o I Subbotina velascoensis		Esa Zanuiradina rauria		e.o.1 Parasubbotina variospira/Igorina pusilla eo.s.1 Eoglobigerina spiralis	• 61.6 • Eoglobigerina eobulloides /Globoconusa daubjergensis			Fe4.7 Hedbergella holmedelensis/H. monmouthensis	65.2 1 Parvularugoglobigerina eugubina 65.451 Guembelitria cretacea	
Diagnostic First Occurrence of Planktonic Foraminifera	عدم العدم ا	28. 1 Globigerinita praestainforthii. Turborotalita praequinqueloba		28.7 Paragloborotalla opima 30.3 Tenutella neoclemenciae	$\neg \neg$		sa. Calobigerina ampliapertura sa. Camolundordalia curalensis sa. L'indoordalia pseuchoampliapertura	_	38.0 Praetenutetta impanapertura	38.5 Subbound angliporordes 38.2 Carinoturborotalia cocoaensis	40.2 Orbulinoides beckmanni		42.3 Hantkenina alabamensis/Truncorotaloides topilensis	<u> 432</u> Hantkenina mexicana	44.9 Acarinina spinuloinflata, Turborotalia centralis			478 Igorina anapetes, Aragonella nuttalli 480 Globigerina hagni, Morozovella caucasica	50.1 Dentoglobigerina galavisi, Acarinina collactea	$\overline{}$	4.7.1 Acarinina pentacamerata 52.3 Monzovella aradorensis	_		ss.o Morozovella subbotinae, Igorina broedermanni	88.4 Globanomalina australiformis, Acarinina wilcoxensis	szo Acarinina esnaensis, A. primitiva	882 Acarinina mckannai, Pseudomenardella pseudomargaritae	60.0 Morozovella acuta, Morozovella velascoensis			62.0 Subbotina triangularis, Praemurica uncinata	83.5 Subbotina fringa. S. triloculinoides, Parasubbotina varianta	7-	es.मs Parvularugoglobigerina eugubina,Globanomalina planocompressa	
Planktonic Zonation (this study)	а	q	a	ற ம	. σ Ω	аΩ	a 2 a	ه م	υ <u>τ</u>	4	- m-c	a c	ŀ	<u>о</u>	,	٩	a	q	w t	эœ	a p	q	æ	q	а	U	٩	а	ه و	σ		O A	σ	0	
	P22	P21		P20	P19	P18	P17	P15	\downarrow	P14	P13	P12	+	<u> </u>		1	Ĺ	å	<u>:</u>	88	P7	g	2	4	2		P4		E	6	72	<u>7</u>	Τ	۵	
Wade <i>et al</i> 2011	90	8	04	03	05	10	E15	E14		E13	E12	E11	E10	63 1	8		E7b		E7a	E6	E5	E4	E3	E2 E1	P5		P4		B 3	ь	27 2	P1c	P1a	Pa and P0	
Berggren and Pearson 2005	90	05	04	03	02	01	E16 E15	E14		E13	E12	E11	E10	E9			E8		E7	E6	E5	E4	E3	E2 E1	P5		P4		P3	8	2 2	P1c	P1a	Pa and P0	
iod, och, ige	neitte			- "	edn _B		nsinod		ue	inoh	Ba	Ľ			eitetu	п		Γ			presian	λ			вп	sitensr	11		elano	3	_		Dan		
Period, Epoch, Stage	əte		oogil	Oligo Early			əte	1	Eocene Middle										Евну							Late			aleo Midd	Д_	Early				
Age (Ma)	0.62	1.82	11	_		9.66		8.78			Z.11	$\overline{}$					8.7	<u>'</u> †					(0.98		ē	Z [*] 69			19				0.88	

Fig. 2. Diagnostic First and Last Occurrences of Paleogene Planktonic Foraminifera.

Period	Diagnostic Last Occurrence of Planktonic Foraminifera	Globorotalia flexuosa 1.2 Truncorotalia hess/Truncorotalia tosaensis	Neogloboquadrina acostaensis Se Globorotalia cibaoensis Se Globorotalia juanai	42 Globorotalia conomiozea			8.6 Paragloborotalia paralenguaensis			Г	120 readyoborotalia peripreroacuta 128 Globorotalia praefohsi	◆ 13.6 Paragloborotalia peripheroronda	Globorotalia archeomenardii 14.5 Globigerinoides diminutus	15.0 Paragloborotalia semivera, Globoturborotalita brazieri 15.9 Dentoglobigerina praedehiscens	г		18.0 Globigerinita praestainforthi	,	21.0 Globigerina ciperoensis	22.0 1 Globigerina angulisuturalis
Period, Berggren Stage et al., Stage beist.	Diagnostic First Occurrence of Planktonic Foraminifera	- ככל	י חדרו	7 7-	ר ו	_	וח-	ו ו	\neg			ירו	Paragloborotalia peripheroacuta, Globorotalia rifensis S Orbulina universa Orbulina suriversa		7	ררו	i	¬-	i –	Paragloborotalia kugleri, Globoturborotalita brazieri,
Early Miocene Middle Miocene Discene Pleist. May Miocene Middle Miocene	ic (y	 	* 	† †		,	 	╅┰╅	$\overline{}$	/ 	, †	╫	4 4 4	╅╅┷	* * `	* * *		+		
Early Miocene Middle Miocene Discene Pleist. May Miocene Middle Miocene Miocen	Planktor Zonation his stud	ЩШ	$\sqcup \sqcup \sqcup$	\perp	٩		Ш	2	4	3		Ш	0 g		#			a	q	
Early Miocene Middle Miocene Late Miocene Pleist. M M M M M M M M M M M M M M M M M M M		Π	 ' T							M10			•			Т		7		
Early Miocene Middle Miocene Late Miocene Pleist.		PT1	PL3-PL6	PL1		M13-M14		M12	M11	M10	M8-M9	M7	We We	M5	M4	M3	. M2			Ž
H I I I I I I I I I I I I I I I I I I I	eriod, 3och, tage		Gela.	elonsZ nsi	nissəM		nsino	Tort		lian	Serraval	6	nsidbi	пе		nsilsgi	Burd	ı	isinsti	nb∀
(\$M) 9QA [18.1] 88.3 89.11 79.31 80	(sM) egA 页 页 众	Η				liocene	V əte		CO.11		eueo	oiM əl	IbbiM				Miocene	Early		50.62

Fig. 3. Diagnostic First and Last Occurrences of Neogene Planktonic Foraminifera.

This article should be referred to as:

BouDagher-Fadel, M.K. Diagnostic First and Last Occurrences of Mesozoic and Cenozoic Planktonic Foraminifera. *UCL Office of the Vice-Provost Research, Professional Papers Series*, **No.1**, 2013, pp.1-4. And can be found at:

http://discovery.ucl.ac.uk/1396461/