



RESEARCH ARTICLE

**REVISED** From zebrafish heart jogging genes to mouse and human orthologs: using Gene Ontology to investigate mammalian heart development. [version 2; referees: 2 approved]

Varsha K Khodiyar<sup>1</sup>, Doug Howe<sup>2</sup>, Philippa J Talmud<sup>1</sup>, Ross Breckenridge<sup>3</sup>, Ruth C Lovering<sup>1</sup>

<sup>1</sup>Cardiovascular GO Annotation Initiative, Centre for Cardiovascular Genetics, Institute of Cardiovascular Science, University College London, London, WC1E 6JF, UK

<sup>2</sup>The Zebrafish Model Organism Database, University of Oregon, Eugene, OR, 97403-5291, USA

<sup>3</sup>Centre for Metabolism and Experimental Therapeutics, University College London, London, WC1E 6JF, UK

**v2** First published: 13 Nov 2013, 2:242 (doi: [10.12688/f1000research.2-242.v1](https://doi.org/10.12688/f1000research.2-242.v1))  
 Latest published: 10 Feb 2014, 2:242 (doi: [10.12688/f1000research.2-242.v2](https://doi.org/10.12688/f1000research.2-242.v2))

**Abstract**

For the majority of organs in developing vertebrate embryos, left-right asymmetry is controlled by a ciliated region; the left-right organizer node in the mouse and human, and the Kupffer’s vesicle in the zebrafish. In the zebrafish, laterality cues from the Kupffer’s vesicle determine asymmetry in the developing heart, the direction of ‘heart jogging’ and the direction of ‘heart looping’. ‘Heart jogging’ is the term given to the process by which the symmetrical zebrafish heart tube is displaced relative to the dorsal midline, with a leftward ‘jog’. Heart jogging is not considered to occur in mammals, although a leftward shift of the developing mouse caudal heart does occur prior to looping, which may be analogous to zebrafish heart jogging. Previous studies have characterized 30 genes involved in zebrafish heart jogging, the majority of which have well defined orthologs in mouse and human and many of these orthologs have been associated with early mammalian heart development.

We undertook manual curation of a specific set of genes associated with heart development and we describe the use of Gene Ontology term enrichment analyses to examine the cellular processes associated with heart jogging. We found that the human, mouse and zebrafish ‘heart jogging orthologs’ are involved in similar organ developmental processes across the three species, such as heart, kidney and nervous system development, as well as more specific cellular processes such as cilium development and function. The results of these analyses are consistent with a role for cilia in the determination of left-right asymmetry of many internal organs, in addition to their known role in zebrafish heart jogging.

This study highlights the importance of model organisms in the study of human heart development, and emphasises both the conservation and divergence of developmental processes across vertebrates, as well as the limitations of this approach.

**Open Peer Review**

Referee Status:

|   | Invited Referees |            |
|---|------------------|------------|
|   | 1                | 2          |
| <b>REVISED</b><br>version 2<br>published<br>10 Feb 2014 |                  |            |
| version 1<br>published<br>13 Nov 2013                   | <br>report       | <br>report |

- 1 **Vincent VanBuren**, Texas A&M University  
USA
- 2 **Jeroen Bakkers**, Hubrecht Institute  
Netherlands

**Discuss this article**

Comments (0)

**Corresponding author:** Ruth C Lovering ([r.lovering@ucl.ac.uk](mailto:r.lovering@ucl.ac.uk))

**How to cite this article:** Khodiyar VK, Howe D, Talmud PJ *et al*. **From zebrafish heart jogging genes to mouse and human orthologs: using Gene Ontology to investigate mammalian heart development.** [version 2; referees: 2 approved] *F1000Research* 2014, 2:242 (doi: [10.12688/f1000research.2-242.v2](https://doi.org/10.12688/f1000research.2-242.v2))

**Copyright:** © 2014 Khodiyar VK *et al*. This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

**Grant information:** The Cardiovascular GO Annotation Initiative is funded by the British Heart Foundation (SP/07/007/23671 and RG/13/5/30112) and PJT is supported by the British Heart Foundation (RG08/014). The Zebrafish Model Organism Database is funded by the National Human Genome Research Institute (P41 HG002659) of the National Institutes of Health.

*The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

**Competing interests:** The authors declare that they have no conflict of interests. VKK is currently employed by *F1000Research*. Her role at the journal does not include any involvement in the pre-publication editorial checks, or with the refereeing process.

**First published:** 13 Nov 2013, 2:242 (doi: [10.12688/f1000research.2-242.v1](https://doi.org/10.12688/f1000research.2-242.v1))

**REVISED Amendments from Version 1**

The main changes from the previous version were included to address comments from the reviewers:

The methods subsection - Generation of the list of zebrafish jogging genes section: edited to make it clearer how the list of zebrafish jogging genes was generated and the limitation of the approach taken to create a list of 'heart jogging genes'.

In Discussion - paragraph 4: edited to create more positive statements about the role of the 'jogging ortholog' genes, in human heart development.

**See referee reports**

## Introduction

An understanding of heart development is important for the treatment of both congenital and acquired heart disease. The majority of heart development studies use model organisms for ethical and practical reasons. Transparent fish embryos, as well chick embryos, enable the developing heart to be studied in real time<sup>1</sup>, and the mouse continues to be a key model organism used to investigate mammalian heart development<sup>2</sup>. Although there is substantial evolutionary conservation in the development of left-right axis asymmetry, there is divergence between species<sup>3</sup>. The earliest events in mammalian heart development are of great interest, but are poorly understood relative to externally developing organs, due to practical constraints.

For the majority of developing vertebrate embryos left-right asymmetry is controlled by a ciliated region; the left-right organizer node in the mouse and human, and the Kupffer's vesicle in the zebrafish<sup>4,5</sup>. In the zebrafish, laterality cues from the Kupffer's vesicle determine asymmetry in the developing heart, and consequently the direction of heart jogging and heart looping. At 24 hours post-fertilization (hpf) the symmetrical zebrafish heart tube is displaced relative to the dorsal midline, with a leftward 'jog'. At 36hpf the heart tube then loops to the right to create the asymmetric heart<sup>5,6</sup>. Cilia within the Kupffer's vesicle are known to be instrumental in establishing left-right asymmetry and consequently play a significant role in determining the direction of heart jogging<sup>7</sup> and heart looping<sup>8</sup>. However, a failure of heart jogging does not necessarily imply that there will be a failure in heart looping, and vice versa. In addition, asymmetric cell migration has been implicated as a key factor in the process of heart jogging<sup>9-13</sup>. Several of the genes involved in zebrafish heart jogging have been identified from mutation, morpholino and functional complementation studies<sup>6,10,13-26</sup>.

We sought to determine whether the use of Gene Ontology (GO) annotation could offer mechanistic clues to early mammalian heart development. GO is a controlled vocabulary that is used to describe gene product function<sup>27</sup>. GO describes three aspects of a gene product's biology: the *biological process* that the gene product is involved in, the specific *molecular function* of the gene product and the *cellular component* that the gene product is located in. GO terms are associated in a directed acyclic graph (DAG), and thus have defined relationships to each other.

The process of heart looping has been described in a variety of higher eukaryotes<sup>2,28,29</sup>, and the occurrence of dextral-looping, the

early phase of heart looping, appears to be conserved from zebrafish to chicken to humans. In addition, many congenital heart abnormalities, such as dextrocardia and isomerisms are thought to be due to abnormal heart looping<sup>2,30</sup> and ciliary dysfunction has been associated with 50% of patients with congenital heart disease and heterotaxy<sup>31</sup>. However, the process of heart jogging has only been described in zebrafish<sup>6</sup>. Biben and Harvey describe a leftward shift of the developing mouse caudal heart prior to looping, which may be analogous to heart jogging in zebrafish<sup>28</sup>, but to our knowledge this has not been investigated further, and heart jogging is not considered to occur in mammals. Consequently, when the ontology describing heart development was expanded<sup>32</sup>, limitations were included to prevent the association of the GO term 'heart jogging' to mammalian gene products<sup>33</sup>. However, an absence of evidence is not evidence of absence, hence it remains a possibility that heart jogging also occurs in mammalian systems.

Although there has been substantial progress in heart development research<sup>1,3,4,29</sup>, there are clearly gaps in our understanding of early heart development, particularly in the mammal. Functional enrichment analysis of genes known to be involved in zebrafish heart jogging, and also of the human and mouse orthologs of these zebrafish heart jogging genes, identifies many conserved biological processes, functions and cellular locations across these three species. The results of these analyses support the role of cilia in symmetry breaking and the importance of cell signalling in early heart development.

## Methods

### Generation of the list of zebrafish heart jogging genes

A list of 30 zebrafish genes that affect heart jogging was compiled using a variety of approaches. Twelve zebrafish proteins were identified as they were already annotated to the 'heart jogging' GO terms, the remaining 18 proteins were then identified using the ZFIN (<http://zfin.org/>) Site Search, with the search phrase 'heart jogging', and filtering using the 'Expression/Phenotypes' category. This search retrieves figures from papers that have 'heart jogging' in the figure legend, and thus are likely to be describing specific zebrafish genes (and proteins) involved in this process. Many of these genes had not yet been curated with GO terms. Each of the papers identified in this way were reviewed; of the 23 zebrafish genes identified in these papers five (Bmpr1aa, Tbx1, unnm\_hu119, unnm\_hu202, unnm\_hu304) were eliminated, as none of these papers provided experimental evidence for the involvement of these genes in heart jogging. This left 30 zebrafish proteins with strong evidence for a role in the heart jogging process (Table 1). The experimental evidence describing the association of each gene to the process of heart jogging was manually reviewed, to ensure consistent criteria were applied.

### Generation of the list of human and mouse 'jogging ortholog' genes

The HUGO Gene Nomenclature Committee Comparison of Orthology Predictions (HCOP) search tool (<http://www.genenames.org/cgi-bin/hcop.pl>) was used to identify the closest possible human and mouse ortholog for each of the 30 zebrafish genes. HCOP displays predictions from 11 homology prediction tools, including EnsemblCompara, Homologene and Inparanoid<sup>34</sup>. For all but one gene, *southpaw*, HCOP returned human or mouse homologs for the zebrafish genes. The lack of a close mammalian ortholog of *southpaw*

**Table 1. Proteins included in zebrafish 'jogging' gene list and the human and mouse 'jogging ortholog' gene lists.** The evidence for these 30 zebrafish proteins having a role in heart jogging comes from mutant, morpholino or functional complementation studies, as described in the associated publications.

| Zebrafish gene symbol (protein ID)       | Human gene symbol (protein ID) | Mouse gene symbol (protein ID) |
|--|--------------------------------|--------------------------------|
| <i>acvr1<sup>F</sup></i> (Q9DGI6)        | <i>ACVRL1</i> (P37023)         | <i>Acvr11</i> (Q61288)         |
| <i>apc<sup>57</sup></i> (F1QN37)         | <i>APC</i> (P25054)            | <i>Apc</i> (Q61315)            |
| <i>bmp4<sup>4,13</sup></i> (O57574)      | <i>BMP4</i> (P12644)           | <i>Bmp4</i> (P21275)           |
| <i>bmp7a<sup>6</sup></i> (Q9PTF9)        | <i>BMP7</i> (P18075)           | <i>Bmp7</i> (P23359)           |
| <i>bmpr2a<sup>20</sup></i> (Q288P3)      | <i>BMPR2</i> (Q13873)          | <i>Bmpr2</i> (O35607)          |
| <i>bmpr2b<sup>20</sup></i> (Q288P2)      |                                |                                |
| <i>camk2a<sup>14</sup></i> (Q32PV2)      | <i>CAMK2A</i> (Q9UQM7)         | <i>Camk2a</i> (P11798)         |
| <i>camk2b<sup>214</sup></i> (E7F012)     | <i>CAMK2B</i> (Q13554)         | <i>Camk2b</i> (P28652)         |
| <i>camk2g<sup>114</sup></i> (Q4V9P8)     | <i>CAMK2G</i> (Q13555)         | <i>Camk2g</i> (Q923T9)         |
| <i>ccdc103<sup>3</sup></i> (Q6DGB6)      | <i>CCDC103</i> (Q8IW40)        | <i>Ccdc103</i> (Q9D9P2)        |
| <i>ccdc40<sup>5</sup></i> (Q56A40)       | <i>CCDC40</i> (Q4G0X9)         | <i>Ccdc40</i> (Q8B179)         |
| <i>cobl<sup>23</sup></i> (I1X3U9)        | <i>COBL</i> (O75128)           | <i>Cobl</i> (Q5NBX1)           |
| <i>dand5<sup>15</sup></i> (Q76C29)       | <i>DAND5</i> (Q8N907)          | <i>Dand5</i> (Q76LW6)          |
| <i>dnaaf1<sup>6,10</sup></i> (Q7ZV84)    | <i>DNAAF1</i> (Q8NEP3)         | <i>Dnaaf1</i> (Q9D2H9)         |
| <i>dub<sup>22</sup></i> (Q0P484)         | <i>RCSL1</i> (Q6JBY9)          | <i>Rcsd1</i> (Q3UZA1)          |
| <i>fgfr2<sup>18</sup></i> (Q8JG38)       | <i>FGFR2</i> (P21802)          | <i>Fgfr2</i> (P21803)          |
| <i>foxh1<sup>6,58</sup></i> (Q9I9E1)     | <i>FOXH1</i> (O75593)          | <i>Foxh1</i> (O88621)          |
| <i>foxj1a<sup>25</sup></i> (Q08CI2)      | <i>FOXJ1</i> (Q92949)          | <i>Foxj1</i> (Q61660)          |
| <i>foxj1b<sup>25</sup></i> (F1R8Z9)      |                                |                                |
| <i>fzd2<sup>22</sup></i> (Q90YL7)        | <i>FZD2</i> (Q14332)           | <i>Fzd2</i> (Q9JIP6)           |
| <i>gsk3b<sup>17</sup></i> (Q9IBD2)       | <i>GSK3B</i> (P49841)          | <i>Gsk3b</i> (Q9WV60)          |
| <i>has2<sup>13</sup></i> (Q9DG41)        | <i>HAS2</i> (Q92819)           | <i>Has2</i> (P70312)           |
| <i>lrrc6<sup>5</sup></i> (B3DH20)        | <i>LRRC6</i> (Q86X45)          | <i>Lrrc6</i> (O88978)          |
| <i>nipbla<sup>21</sup></i> (F5HSE3)      | <i>NIPBL</i> (Q6KC79)          | <i>Nipbl</i> (Q6KCD5)          |
| <i>Nipblb<sup>21</sup></i> (F1QBY1)      |                                |                                |
| <i>nkd1<sup>24</sup></i> (Q2TJA6)        | <i>NKD1</i> (Q969G9)           | <i>Nkd1</i> (Q99MH6)           |
| <i>nphp3<sup>26</sup></i> (POCI65)       | <i>NPHP3</i> (Q7Z494)          | <i>Nphp3</i> (Q7TNH6)          |
| <i>pkd2<sup>8</sup></i> (Q6IVV8)         | <i>PKD2</i> (Q13563)           | <i>Pkd2</i> (O35245)           |
| <i>ptpn11a<sup>16</sup></i> (Q7ZW17)     | <i>PTPN11</i> (Q06124)         | <i>Ptpn11</i> (P35235)         |
| <i>southpaw<sup>10,19</sup></i> (Q7ZZT5) | no mammalian orthologs         |                                |

was confirmed with a UCSC BLAT analysis against the human and mouse genomes<sup>35</sup>. BLAST analysis<sup>36</sup> showed that the closest possible human and mouse homolog for the zebrafish *southpaw* gene was *Nodal* (33% identity). Indeed, both *southpaw* and *nodal* are specifically expressed in the left lateral plate mesoderm<sup>5,37</sup> and knockdown of murine *Nodal* in this region leads to a disruption of cardiac asymmetry, as does injection of *southpaw* morpholinos, suggesting a functional orthology between *southpaw* and *Nodal*<sup>5,37</sup>. However a reciprocal HCOP search showed that the zebrafish genes *nodal-related 1* and *2* are the closest orthologs of human *NODAL*. Hence we have not included a human or mouse ortholog for zebrafish *southpaw* (Table 1). Three pairs of zebrafish paralogs (*bmpr2a/bmpr2b*; *foxj1a/foxj1b*; *nipbla/nipblb*) have a single corresponding ortholog in human and mouse. Therefore, there are

26 human and 26 mouse orthologs to the 30 zebrafish genes identified as relevant to zebrafish heart jogging (Table 1).

#### Gene ontology annotation

The human 'jogging ortholog' genes were fully manually annotated, by an experienced GO curator<sup>38</sup>. Individual PubMed queries were run for each gene using the approved human gene symbol and filtering on 'human'. To achieve full annotation, all of the relevant publications (a total of 232) containing unique functional data for each gene were annotated, regardless of the specific biology described in each paper. This approach enabled consistent annotation of all experimental data relating to each gene, thus ensuring an unbiased overview of any common processes associated with these genes. In addition, the GO term 'heart looping' was associated with a 'jogging

ortholog' human gene if dextrocardia or *situs inversus totalis* phenotypes had been associated with a mutation in the gene, in order to follow the generally agreed view that leftward heart looping will have resulted in these phenotypes<sup>2</sup>.

### Functional enrichment analysis

The Mouse Genome Informatics functional enrichment tool VLAD (Visual Annotation Display; <http://proto.informatics.jax.org/prototypes/vlad-1.0.3/>) was used to look for overrepresentation of GO terms in each gene list relative to the whole genome of the organism. The annotation datasets used for the analysis were zfin (4th March 2013), goa\_human (5th March 2013) and mgi (7th March 2013) for the zebrafish, human and mouse analyses respectively, and the ontology dataset used was dated 10th March 2013. The query gene lists (as UniProt IDs) were pasted into the 'Query Set' field, the 'Universe Set' field was left blank (to specify all genes in species specific annotation file) and the 'Display Settings' options selected were 'pruning threshold':3 and 'collapsing threshold':6. No evidence codes were excluded from the analyses. For this analysis the total number of genes (universe set size) having annotations in the *biological process* ontology were 14,577, 30,441 and 24,813 for zebrafish, human and mouse respectively. In line with common practice, when using functional analysis tools, enriched GO terms with 1 or 2 associated query genes were excluded from the final results table.

### Creation of an 'early heart development' mouse gene list

A list of 103 mouse genes likely to play a role in early heart development was created by combining gene lists derived from three sources: The Mouse Genome Informatics Mammalian Phenotype Ontology browser [http://www.informatics.jax.org/searches/MP\\_form.shtml](http://www.informatics.jax.org/searches/MP_form.shtml)<sup>39</sup>, the QuickGO browser <http://www.ebi.ac.uk/QuickGO/><sup>40</sup> and the 'jogging ortholog' gene list described above (see Mousegenelist.csv in [Data File](#)). The Mammalian Phenotype Ontology browser was queried for genotypes annotated with the terms 'abnormal direction of heart looping', '*situs inversus totalis*', 'dextrocardia' and 'mesocardia', creating a list of 180 genotypes with an associated gene. Due to the multiple phenotypes associated with each of these genotypes only 58 genes were identified through this approach, and of these only 5 overlap with the 26 'jogging ortholog' genes. Thirty-five genes were identified by filtering on the GO term 'determination of heart left/right asymmetry' and its child terms, the evidence code IMP (Inferred by Mutant Phenotype), and the mouse taxon. Of these only two are also present in the 'jogging ortholog' gene lists and 11 are present in the phenotype gene list. Twenty-six mouse 'jogging ortholog' genes were added to this combined gene list, and any duplicated genes were removed.

## Results

### Annotation of the zebrafish heart jogging genes and the human 'jogging ortholog' genes

Thirty zebrafish genes were annotated to the GO term 'heart jogging' or one of its child terms based on experimental data from the literature ([Table 1](#)). Human and mouse orthologs of these genes were identified, as described in the Methods section, resulting in a list of 26 mammalian 'jogging orthologs'.

The human 'jogging ortholog' genes were then fully annotated with GO terms based on published experimental data. All manual

annotations to the human, mouse and zebrafish genes can be visualized with the QuickGO Gene Ontology browser <http://tinyurl.com/humanortholog>, <http://tinyurl.com/mouseortholog> and <http://tinyurl.com/zebrafishgenes>.

### Functional enrichment analysis

The zebrafish heart jogging gene list and the human and mouse 'jogging ortholog' gene lists were analysed using the VLAD enrichment tool. This identified 155 *biological process* GO terms that were significantly enriched in the zebrafish (see Human\_data.csv in [Data File](#)), 431 in the human (see Human\_data.csv in [Data File](#)) and 402 in the mouse (see Mouse\_data.csv in [Data File](#)) gene lists. The enriched GO terms from all three species were grouped into five biological areas: Development, Patterning, Cellular Process, Signalling and Movement. The relative enrichment of key GO terms from each area was compared across all three species (see [Biological\\_process\\_summary.csv](#) in [Data File](#); summarized in [Table 2](#)).

**Enrichment of heart development terms.** As expected there was a significant enrichment of developmental process terms in all three gene lists, including an enrichment of the GO term 'heart development'. However, there was also enrichment of terms such as 'renal system development' and 'nervous system development', indicating the role of these proteins in regulating the development of a range of organ systems and tissues. These data analyses also show an enrichment of terms describing specific, but universal, cellular processes, such as signalling and regulation of transcription ([Table 2](#)). These terms represent essential aspects of development, but are grouped discretely due to their roles in many other biological processes.

'Pattern specification', described in GO as a 'developmental process that results in the creation of defined areas or spaces within an organism to which cells respond and eventually are instructed to differentiate' and several of its more specific child terms (such as 'specification of symmetry'), were also enriched in all three gene lists. Within the symmetry ontology, the GO term 'determination of heart left/right asymmetry' is annotated to all 30 genes in the zebrafish jogging gene list, however, it is only associated with 8 and 4 jogging ortholog genes in the human and mouse respectively. Of the 97 zebrafish genes associated with 'determination of heart left/right asymmetry' 31% are also present in the zebrafish jogging gene list. In contrast, only 11% of the human and 8% of the mouse genes associated with this term are also 'jogging orthologs'. These results confirm an overlap in the functional role of the zebrafish jogging genes and the human and mouse orthologs in the determination of heart left/right symmetry. However, this relatively low level of overlap may reflect the limitations of model organism and human research in this area, rather than a lack of functional conservation of these genes.

In addition, there were some differences in the developmental terms that were enriched between species. For example, the GO terms 'vasculature development' and 'sensory organ development' are enriched in both the human and mouse 'jogging ortholog' gene lists ([Table 2](#)), but neither of these processes are enriched in the zebrafish jogging ortholog genes. This difference may reflect the type of experiments zebrafish are used for, rather than reflecting a difference between zebrafish and mammals in the genes required for these developmental processes.


**Table 2. Comparison of enriched Gene Ontology terms across orthologous gene lists from zebrafish, human and mouse.** The enriched GO terms were grouped into specific ontology areas, with a selection of more specific child term (preceded with a dash) also included. The full list of grouped GO terms can be found in Table S4, which also shows the genes annotated to each term from each of the three species. k: the number of genes in each gene list annotated to the GO term; M: the number of genes in the species proteome annotated to the GO term.

| Gene Ontology terms  | Zebrafish |      |          | Human |       |          | Mouse |      |          |
|--|-----------|------|----------|-------|-------|----------|-------|------|----------|
|  | k         | M    | k/M as % | k     | M     | k/M as % | k     | M    | k/M as % |
| <b>DEVELOPMENT</b>   |           |      |          |       |       |          |       |      |          |
| GO:0032502 developmental process                                       | 30        | 2357 | 1.3%     | 24    | 6803  | 0.4%     | 22    | 3945 | 0.6%     |
| - GO:0009888 tissue development  | 30        | 673  | 4.5%     | 17    | 1849  | 0.9%     | 14    | 1138 | 1.2%     |
| - GO:0072358 cardiovascular system development                         | 30        | 487  | 6.2%     | 15    | 1095  | 1.4%     | 10    | 679  | 1.5%     |
| - GO:0001944 vasculature development                                   | -         | -    | -        | 7     | 666   | 1.1%     | 7     | 436  | 1.6%     |
| - GO:0007507 heart development   | 30        | 268  | 11.2%    | 13    | 634   | 2.1%     | 8     | 390  | 2.1%     |
| - GO:0001947 heart looping   | 22        | 83   | 26.5%    | 7     | 68    | 10.3%    | -     | -    | -        |
| - GO:0007399 nervous system development                                | 8         | 700  | 1.1%     | 15    | 2802  | 0.5%     | 12    | 1486 | 0.8%     |
| - GO:0072001 renal system development                                  | 6         | 100  | 6.0%     | 9     | 386   | 2.3%     | 6     | 200  | 3.0%     |
| - GO:0007423 sensory organ development                                 | -         | -    | -        | 10    | 738   | 1.4%     | 8     | 480  | 1.7%     |
| - GO:0048736 appendage development                                     | 4         | 123  | 3.3%     | 4     | 237   | 1.7%     | 4     | 159  | 2.5%     |
| - GO:0050793 regulation of developmental process                       | 6         | 267  | 2.2%     | 14    | 2383  | 0.6%     | 14    | 1542 | 0.9%     |
| <b>PATTERNING</b>  |           |      |          |       |       |          |       |      |          |
| GO:0007389 pattern specification process                               | 30        | 435  | 6.9%     | 15    | 644   | 2.3%     | 12    | 414  | 2.9%     |
| - GO:0009799 specification of symmetry                                 | 30        | 177  | 16.9%    | 10    | 147   | 6.8%     | 7     | 90   | 7.8%     |
| - GO:0061371 determination of heart left/right asymmetry               | 30        | 97   | 30.9%    | 8     | 71    | 11.3%    | 4     | 48   | 8.3%     |
| <b>CELLULAR PROCESS</b>  |           |      |          |       |       |          |       |      |          |
| GO:0071840 cellular component organization or biogenesis               | 12        | 1124 | 1.1%     | 17    | 5991  | 0.3%     | 15    | 3306 | 0.5%     |
| - GO:0030030 cell projection organization                              | 10        | 274  | 3.6%     | 12    | 1311  | 0.9%     | 11    | 644  | 1.7%     |
| - GO:0051128 regulation of cellular component organization             | -         | -    | -        | 10    | 1894  | 0.5%     | 10    | 1293 | 0.8%     |
| - GO:0031344 regulation of cell projection organization                | -         | -    | -        | 5     | 452   | 1.1%     | 5     | 311  | 1.6%     |
| GO:0006468 protein phosphorylation                                     | 8         | 696  | 1.1%     | 8     | 842   | 1.0%     | 9     | 748  | 1.2%     |
| - GO:0001932 regulation of protein phosphorylation                     | -         | -    | -        | 8     | 1124  | 0.7%     | 8     | 722  | 1.1%     |
| GO:0006357 regulation of transcription from RNA polymerase II promoter | 8         | 166  | 4.8%     | 10    | 1939  | 0.5%     | 12    | 1268 | 0.9%     |
| GO:0042127 regulation of cell proliferation                            | 5         | 70   | 7.1%     | 9     | 1769  | 0.5%     | 9     | 1123 | 0.8%     |
| GO:0007049 cell cycle  | -         | -    | -        | 9     | 1644  | 0.5%     | 8     | 915  | 0.9%     |
| - GO:0051726 regulation of cell cycle                                  | -         | -    | -        | 6     | 934   | 0.6%     | 6     | 538  | 1.1%     |
| <b>SIGNALLING</b>  |           |      |          |       |       |          |       |      |          |
| GO:0023052 signaling   | 12        | 2131 | 0.6%     | 16    | 6682  | 0.2%     | 16    | 4568 | 0.4%     |
| - GO:0023051 regulation of signaling                                   | 7         | 521  | 1.3%     | 16    | 3227  | 0.5%     | 16    | 1931 | 0.8%     |
| GO:0050896 response to stimulus  | -         | -    | -        | 20    | 10767 | 0.2%     | 21    | 6679 | 0.3%     |
| - GO:0048583 regulation of response to stimulus                        | 7         | 557  | 1.3%     | 15    | 3799  | 0.4%     | 15    | 2140 | 0.7%     |
| <b>MOVEMENT</b>  |           |      |          |       |       |          |       |      |          |
| GO:0007017 microtubule-based process                                   | 6         | 188  | 3.2%     | 6     | 873   | 0.7%     | 4     | 358  | 1.1%     |
| GO:0040011 locomotion  | 10        | 289  | 3.5%     | 10    | 1448  | 0.7%     | 8     | 753  | 1.1%     |
| - GO:0040012 regulation of locomotion                                  | -         | -    | -        | 6     | 657   | 0.9%     | 5     | 473  | 1.1%     |
| - GO:2000145 regulation of cell motility                               | -         | -    | -        | 6     | 608   | 1.0%     | 5     | 439  | 1.1%     |
| - GO:0016477 cell migration  | 8         | 200  | 4.0%     | 6     | 931   | 3.6%     | 5     | 504  | 1.0%     |

**Enrichment of cilia terms.** Terms in the cellular component organization or biogenesis ontology were enriched across all three gene lists (Table 2 and Biological\_process\_summary.csv in Data File). Specifically there was an enrichment of terms describing ‘cilium morphogenesis’ and ‘protein complex assembly’ (Figure 1). Within each of these, some more specific terms were enriched, for example the human and mouse ‘jogging ortholog’ gene lists were enriched for the term ‘axonemal dynein complex assembly’, whilst the zebrafish and human gene lists showed an enrichment of the term ‘cilium assembly’.

Terms such as ‘regulation of cell projection organization’ were also enriched in the human and mouse ‘jogging ortholog’ gene lists. ‘Regulation’ terms have a ‘regulates’ relationship with the relevant processes; for example the term ‘positive regulation of cell projection organization’ has a ‘positively\_regulates’ relationship to the term ‘cell projection organization’. In GO an important benefit of building

a DAG, rather than a flat-list of controlled vocabulary terms, is that relationships can be used to make inferences from one term to another. However, the VLAD enrichment tool does not automatically create a transitive relationship between ‘regulation’ terms and the processes or functions they regulate. Consequently genes annotated to a ‘regulation’ term will not be associated with the regulated process term (unless there is an independent annotation to the process term). It is also important to recognise that it can be difficult for a curator to choose between annotating to the *biological process* itself, or to the term describing the regulation of that *biological process*, based on the published experimental data. Therefore, to get a full picture of the genes involved in a process, including the genes that regulate the process, it is necessary to combine the genes annotated to GO terms describing both the ‘process’ and the ‘regulation of the process’. For example, 10 zebrafish, 12 human and 14 mouse genes within the ‘jogging ortholog’ gene lists are annotated to either ‘cell projection organization’ or ‘regulation of cell projection organization’



**Figure 1. View of relationships between enriched terms from the cellular component organization or biogenesis ontology.** The GONUTS view of relationships between enriched terms from the cellular component organization or biogenesis ontology<sup>84</sup>. The grey arrows are used where a term has an ‘is a’ relationship to its parent term, the blue arrows indicate a ‘part of’ relationship. The bars below each GO term indicates which of these terms are enriched in the zebrafish ‘jogging’ gene list, and the human and mouse ‘jogging ortholog’ gene lists.

(or children of these terms). This represents 33%, 40% and 47% of these zebrafish, human and mouse ‘jogging’ gene lists respectively, indicating that the process of cell projection organization is an important function for this group of genes. In addition, many of the ‘jogging’ and ‘jogging ortholog’ genes annotated to ‘cell projection organization’ terms have also been annotated to the cellular component term ‘cell projection’ (7, 11, 13, genes in zebrafish, human and mouse, respectively, see [Data File](#)). The enrichment of the *biological process* term ‘cell projection organization’ and *cellular component* term ‘cell projection’ within these gene lists is consistent with the key role of cilia located in the node/Kuppfer’s vesicle to determine heart left/right asymmetry in all three species.

**Enrichment of cell migration terms.** Cell migration also plays a key role in the establishment of the heart cone, heart jogging and heart looping<sup>9,10</sup> and enrichment of the GO term ‘cell migration’ is seen in the ‘jogging’ gene lists of all three species ([Table 2](#) and [Biological\\_process\\_summary.csv](#) in [Data File](#)). Lenhart *et al.* (2013)<sup>9</sup> identified *FoxH1*, *spaw*, *Bmp4*, *Lefty2* and *Has2* as essential to the asymmetric cell migration that leads to heart jogging. However, our literature review suggests that some genes may have functions in both cilia assembly, within the Kuppfer’s vesicle, and cell migration. For example, thymocytes from *Foxj1* transgenic mice display defective migration<sup>41</sup>, whereas *Foxj1*-null mice are defective in ciliogenesis<sup>42</sup>. Similarly, in zebrafish, *Fzd2* has been shown to play a role in cilium assembly<sup>22</sup> as well as pancreatic insulin-cell migration<sup>43</sup>. Consequently, further investigations into the role of these genes in heart jogging cell migration may provide further insight into this process.

### Co-annotation of heart development associated genes

In order to investigate the contribution of individual genes in the multiple processes associated with early heart development we created human and mouse heart development gene lists and examined the associated GO *biological processes* terms. A list of 103 mouse genes with roles in early heart developmental processes was created by merging the three gene lists created using the Mouse Genome Informatics phenotype browser, the QuickGO browser as well as the ‘jogging ortholog’ gene list ([Mousegenelist.csv](#) in [Data File](#)).

GO captures a range of biological processes that a single gene is involved in. By comparing the overlap between the GO terms associated with specific gene lists it is possible to see what cellular mechanisms are likely to be contributing to the various heart developmental processes. Using the QuickGO browser, genes in the zebrafish ‘heart jogging’ gene list, which were associated with the GO terms ‘heart looping’, ‘signal transduction’, ‘cell migration’ and ‘cell projection organization’ (and all child terms, including ‘regulation’ terms), were downloaded, as well as the genes associated with these terms that were also present in the mouse ‘early heart development’ gene list ([Mousegenelist.csv](#) in [Data File](#)).

In the zebrafish ‘heart jogging’ gene list a similar proportion of the genes have the potential to play a role in cell projection organisation (10 genes), cell migration (8 genes) and signal transduction (13 genes) ([Figure 2A](#)). In the list of 103 mouse genes that are associated with early heart development, either by phenotype, annotation or homology to the zebrafish ‘heart jogging’ gene list,

82 have been annotated to the GO term heart looping. In contrast to the zebrafish ‘jogging’ gene list, signal transduction appears to play a major role in the mouse early heart development, with 27 genes associated with both signal transduction and heart looping, whereas only 18 and 9 genes, respectively, are associated with cell migration and cell projection organization ([Figure 2B](#)). These results fit well with what is known about these gene lists. The zebrafish ‘jogging’ gene list defines a group of genes whose functions are required very early in heart development, when the role of cilia in symmetry breaking initiates the heart jogging process. Whereas, in the mouse ‘early heart development’ gene list the genes included have roles in heart looping, which is developmentally later event than heart jogging. Therefore, although the initial events associated with breaking of left-right symmetry are represented within this gene list, the genes involved in the later process of ensuring the complex looping of the heart tube, through controlled signalling and cell migration, contribute to a large proportion of this list.

### Human disease phenotypes associated with the ‘jogging ortholog’ genes

While annotating the 26 human ‘jogging ortholog’ genes we noticed that almost half of these genes have not been associated with a specific disease phenotype ([Table 3](#)). However, of the 26 genes examined, mutations in 14 had been associated with a disease phenotype, a fifth of which were ciliopathies. Dextrocardia or *situs inversus totalis* (reversal or mirroring of the major visceral organs) was associated with 6 of the human ‘jogging ortholog’ genes. Location of the heart on the right side (rather than the left) is generally agreed to be the result of left-handed, instead of right-handed looping of the heart tube in early embryogenesis<sup>2</sup>. The association of these ‘jogging ortholog’ genes with heart looping defects confirm that there is conserved functional homology between at least some of these orthologous zebrafish and human genes in the very early stages of heart development, which lead to the initial heart asymmetry. All four of ciliopathy-associated ‘jogging orthologs’ were also described as associated with *situs inversus totalis*, confirming the conserved role of these genes in the cilia within the symmetry determining left-right organizer.


#### Significantly enriched GO terms in heart jogging genes in zebrafish and their human and mouse orthologs F1000Research

5 Data Files

<http://dx.doi.org/10.6084/m9.figshare.844630>

### Discussion

We have used GO to annotate the key genes involved in zebrafish heart jogging and their human and mouse orthologs. Heart jogging is not a process that is thought to occur in mammals. However, these genes are conserved between species and play essential roles in many developmental processes. The information available about these genes in several diverse species can be used to shed light on the roles of these genes and possible mechanisms in heart jogging and other heart developmental processes. Our analyses are in agreement with the well described essential role of cilia in early



**Figure 2. Venn diagrams describing the multiple roles of genes associated with heart development.** Venn diagrams showing the overlap between the GO terms associated with **A** the zebrafish 'heart jogging' gene list (30 genes) and **B** the mouse combined heart development gene list (103 genes).

development<sup>4,5,31</sup>, with a third of the zebrafish 'heart jogging' genes associated with the *biological process* 'cell projection organization' (Table 2).

However, it is also important to recognise that although there is considerable evidence for conserved mechanisms of heart development across vertebrates there are also areas of divergence<sup>44</sup>. For example, in the mouse, zebrafish and *Xenopus* the rotation of cilia is responsible for the early asymmetric gene expression pattern around the left-right organizer, whereas cilia do not play a role in symmetry breaking in the chicken or pig<sup>44</sup>.

The early phases of heart development are particularly difficult to study in mammals, however various approaches are enabling progress in this area<sup>2,29,45,46</sup> and using phenotype, annotation and orthology data we have created a list of 103 genes with a putative role in early mouse heart developmental processes. Furthermore, the phenotypes associated with experimentally generated mutant mice provide further clues to the likely role of these genes in human heart development; the genes associated with *situs inversus totalis* phenotypes are most likely to have functional roles within the node. Conversely, genes not associated with *situs inversus totalis* but associated with an abnormal direction of heart looping, dextrocardia or mesocardia are likely to be involved in the response of the embryonic heart tube to the left/right asymmetry signals. This is not a completely reliable interpretation, for example mutations in the transcription factor *Pitx2* lead to mice with *situs inversus totalis*, however, *Pitx2* is expressed in the left lateral plate and its continued asymmetric expression is necessary for asymmetric morphogenesis of most visceral organs<sup>44</sup>. The mouse knockout consortia data<sup>47</sup> will continue to help with the identification of additional early heart development genes, and informed interpretation of these phenotypes will make it possible to separate those

genes likely to be associated with the node from those with functions within the heart tube.

In humans, defects in early heart development are likely to result in spontaneous abortion and therefore many genes required for early heart development will go undetected<sup>48</sup>. Consequently, human embryos with heart defects, which develop to full term, represent the less severe end of the spectrum. Mutations in several human genes have now been identified as causative of abnormal heart looping, such as *ACVR2B*, *LEFTY2*, *GJA1* and *ZIC3*<sup>49-52</sup>, and some of the 'jogging ortholog' genes (*CCDC103*, *CCDC40*, *DNAAF1*, *LRRC6*, *NPHP3* and *PKD2*) are also associated with heart looping defects. Thus providing evidence to support an involvement of these genes in left-right asymmetry determination in the heart. Furthermore, mutations in some of the 'jogging ortholog' human genes, *FOXH1* and *PTPN11*, are associated with heart septal defects in humans, which seems to imply that in individuals with these mutations early heart developmental processes have proceeded normally, suggesting that, contrary to their role in zebrafish, these genes may not be involved in the early stages of human heart development. However, there are other possible reasons why there is a poor association of heart defects with the 'jogging ortholog' gene list. This may simply be due to the lack of detection of *situs inversus totalis*<sup>53</sup>, or reflect a redundancy in gene function, or it may be that the majority of mutations in these genes are simply not detected in humans because they are masked by first trimester spontaneous abortions, which are known to have a high level of heart defects<sup>48</sup>.

The impact of lethal mutations on detection of genes associated with heart development would suggest that mutations in these genes would only be detected in individuals with mutations with relatively minor impact on gene function. This idea is supported by the recent identification of multiple 'minor' heterozygous mutations

**Table 3. Diseases associated with the human ‘jogging ortholog’ genes.** The associated diseases are described in the listed publications.

| <b>Human gene symbol (protein ID)</b> | <b>Heart relevant phenotype</b>   | <b>Other associated phenotypes</b>   |
|---------------------------------------|---|--|
| <b>ACVRL1 (P37023)</b>                | -   | Hereditary haemorrhagic telangiectasia type 2 (HHT2) <sup>59</sup> , HHT2 with pulmonary hypertension <sup>60</sup>                |
| <b>APC (P25054)</b>                   | -   | Familial adenomatous polyposis coli-1 <sup>61</sup>  |
| <b>BMP4 (P12644)</b>                  | -   | Microphthalmia, syndromic 6 <sup>62</sup> , orofacial cleft 11 <sup>63</sup>   |
| <b>BMP7 (P18075)</b>                  | -   | -  |
| <b>BMPR2 (Q13873)</b>                 | -   | Pulmonary hypertension <sup>64</sup>   |
| <b>CAMK2A (Q9UQM7)</b>                | -   | -  |
| <b>CAMK2B (Q13554)</b>                | -   | -  |
| <b>CAMK2G (Q13555)</b>                | -   | -  |
| <b>CCDC103 (Q8IW40)</b>               | Dextrocardia, <i>situs inversus totalis</i> <sup>65</sup>                                   | Ciliary dyskinesia, primary, 17 <sup>65</sup>  |
| <b>CCDC40 (Q4G0X9)</b>                | <i>Situs inversus totalis</i> <sup>66</sup>   | Ciliary dyskinesia, primary, 15 <sup>67</sup> , Kartagener's Syndrome <sup>66</sup>  |
| <b>COBL (O75128)</b>                  | -   | -  |
| <b>DAND5 (Q8N907)</b>                 | -   | -  |
| <b>DNAAF1 (Q8NEP3)</b>                | <i>Situs inversus totalis</i> <sup>68,69</sup>  | Ciliary dyskinesia, primary, 13 <sup>68,69</sup>   |
| <b>FGFR2 (P21802)</b>                 | -   | Several craniosynostosis <sup>70,71</sup> , see OMIM for more information  |
| <b>FOXH1 (O75593)</b>                 | Ventricular septal defect <sup>72</sup> , transposition of the great arteries <sup>54</sup> | -  |
| <b>FOXJ1 (Q92949)</b>                 | -   | -  |
| <b>FZD2 (Q14332)</b>                  | -   | -  |
| <b>GSK3B (P49841)</b>                 | -   | -  |
| <b>HAS2 (Q92819)</b>                  | -   | -  |
| <b>LRR6 (Q86X45)</b>                  | <i>Situs inversus totalis</i> <sup>73</sup>   | Ciliary dyskinesia, primary, 19, Kartagener's Syndrome <sup>73</sup>   |
| <b>NIPBL (Q6KC79)</b>                 | Cardiac septal defects (not confirmed as associated with NIPBL mutations) <sup>74</sup>     | Cornelia de Lange syndrome 1 <sup>74,75</sup>  |
| <b>NKD1 (Q969G9)</b>                  | -   | Colorectal adenocarcinoma <sup>76</sup>  |
| <b>NPHP3 (Q7Z494)</b>                 | <i>Situs inversus totalis</i> <sup>77</sup>   | nephronophthisis type 3 <sup>78</sup> , Meckel syndrome type 7 <sup>77</sup> , renal-hepatic-pancreatic dysplasia <sup>77,79</sup> |
| <b>PKD2 (Q13563)</b>                  | Dextrocardia, <i>situs inversus totalis</i> <sup>53</sup>                                   | Polycystic kidney disease 2 <sup>53,80</sup>   |
| <b>PTPN11 (Q06124)</b>                | atrioventricular canal defects <sup>81</sup>  | juvenile myelomonocytic leukemia <sup>82</sup> , LEOPARD syndrome <sup>81</sup> , Noonan syndrome <sup>81,83</sup>                 |
| <b>RCSD1 (Q6JBY9)</b>                 | -   | -  |

within a functional network in three patients with transposition of the great arteries. All of these genes either participate or cooperate within the Nodal signaling pathway<sup>54</sup> and the carriers of single mutations exhibit no heart or laterality defects. The impact of ‘minor’ mutations, such as these, may explain the contribution of ‘genetic modifiers’ to congenital heart defects with variable penetrance within a family<sup>55</sup>, or may suggest a polygenic basis for some of these diseases<sup>56</sup>. This is supported by model organism data, which provides evidence of multigenic origins for congenital heart disease<sup>56</sup>. However, model organisms are rarely used to examine the impact of genetic modifiers on heart development, as the majority of model organisms are inbred and examination of mutations leading to ‘minor’ phenotypic variations is often not viewed with the same level of interest as the more extreme heart development defects.

Next Generation Sequencing (NGS) has the potential to identify many more instances of multiple mutations in genes which are functionally linked through a specific pathway. However, teasing out which gene mutations are contributing to a disease, as a genetic modifier or as the causative gene variant, and which are not involved in the disease, is likely to take considerable time. Gene Ontology, KEGG and Reactome pathways, along with protein interaction networks have the potential to inform the process of identifying genetic variants associated with heart defect risk through the identification of pathways and networks which are common to the genes associated with the risk gene variants. Consequently, interpretation of NGS data will be greatly improved with full annotation of the candidate genes involved. The identification of these risk gene variants is likely to be of considerable value to those patients seeking prenatal diagnosis. In addition, the identification of more genes

associated with heart defects will also help clarify the conserved and divergent heart development pathways that exist between humans and key model organisms.

## Conclusions

This study demonstrates that full annotation, using GO, of a set of genes known to be associated with early stages of heart development in zebrafish can be used to confirm functional conservation of the role of these genes in a variety of developmental processes. While this study supports the assertion of gene function based on orthology between genes, it also identifies that for some genes there is no direct evidence for their conserved involvement in specific developmental processes through evolution. Consequently, for evolutionary studies, manual annotation of the genome of individual species will be necessary to enable a bioinformatics approach to investigating the evolution of developmental processes.

## Author contributions

VKK conceived and designed the study, undertook curation of the prioritized human genes, analysed the datasets and drafted the manuscript. DH participated in the design of the study, undertook curation of the prioritized zebrafish genes, and drafted the manuscript. PJT and RB participated in the study design and helped to draft

the manuscript. RCL participated in the design of the study, analysed the datasets and drafted the manuscript. All authors read and approved the final manuscript.

## Competing interests

The authors declare that they have no conflict of interests. VKK is currently employed by *F1000Research*. Her role at the journal does not include any involvement in the pre-publication editorial checks, or with the refereeing process.

## Grant information

The Cardiovascular GO Annotation Initiative is funded by the British Heart Foundation (SP/07/007/23671 and RG/13/5/30112) and PJT is supported by the British Heart Foundation (RG08/014). The Zebrafish Model Organism Database is funded by the National Human Genome Research Institute (P41 HG002659) of the National Institutes of Health.

*The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

## Acknowledgements

Many thanks to Dr. Jim Hu and Dr Mary Dolan for their help with the creation of [Figure 1](#) and Dr Fotios Drenos for creating the Venn diagrams in [Figure 2](#).

## References

- Bakkers J: **Zebrafish as a model to study cardiac development and human cardiac disease.** *Cardiovasc Res.* 2011; **91**(2): 279–288.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Manner J: **The anatomy of cardiac looping: a step towards the understanding of the morphogenesis of several forms of congenital cardiac malformations.** *Clin Anat.* 2009; **22**(1): 21–35.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Schlueter J, Brand T: **Left-right axis development: examples of similar and divergent strategies to generate asymmetric morphogenesis in chick and mouse embryos.** *Cytogenet Genome Res.* 2007; **117**(1–4): 256–267.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Shiraishi I, Ichikawa H: **Human heterotaxy syndrome - from molecular genetics to clinical features, management, and prognosis.** *Circ J.* 2012; **76**(9): 2066–2075.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ahmad N, Long S, Rebagliati M: **A southpaw joins the roster: the role of the zebrafish nodal-related gene southpaw in cardiac LR asymmetry.** *Trends Cardiovasc Med.* 2004; **14**(2): 43–49.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chen JN, van Eeden FJ, Warren KS, *et al.*: **Left-right pattern of cardiac BMP4 may drive asymmetry of the heart in zebrafish.** *Development.* 1997; **124**(21): 4373–4382.  
[PubMed Abstract](#)
- Ferrante MI, Romio L, Castro S, *et al.*: **Convergent extension movements and ciliary function are mediated by *ofd1*, a zebrafish orthologue of the human oral-facial-digital type 1 syndrome gene.** *Hum Mol Genet.* 2009; **18**(2): 289–303.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Speder P, Petzoldt A, Suzanne M, *et al.*: **Strategies to establish left/right asymmetry in vertebrates and invertebrates.** *Curr Opin Genet Dev.* 2007; **17**(4): 351–358.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lenhart KF, Holtzman NG, Williams JR, *et al.*: **Integration of Nodal and BMP Signals in the Heart Requires FoxH1 to Create Left-Right Differences in Cell Migration Rates That Direct Cardiac Asymmetry.** *PLoS Genet.* 2013; **9**(1): e1003109.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Baker K, Holtzman NG, Burdine RD: **Direct and indirect roles for Nodal signaling in two axis conversions during asymmetric morphogenesis of the zebrafish heart.** *Proc Natl Acad Sci U S A.* 2008; **105**(37): 13924–13929.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rohr S, Otten C, Abdellah-Seyfried S: **Asymmetric involution of the myocardial field drives heart tube formation in zebrafish.** *Circ Res.* 2008; **102**(2): e12–19.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- de Campos-Baptista MI, Holtzman NG, Yelon D, *et al.*: **Nodal signaling promotes the speed and directional movement of cardiomyocytes in zebrafish.** *Dev Dyn.* 2008; **237**(12): 3624–3633.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Smith KA, Chocron S, von der Hardt S, *et al.*: **Rotation and asymmetric development of the zebrafish heart requires directed migration of cardiac progenitor cells.** *Dev Cell.* 2008; **14**(2): 287–297.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Francescato L, Rothschild SC, Myers AL, *et al.*: **The activation of membrane targeted CaMK-II in the zebrafish Kupffer's vesicle is required for left-right asymmetry.** *Development.* 2010; **137**(16): 2753–2762.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hashimoto H, Rebagliati M, Ahmad N, *et al.*: **The Cerberus/Dan-family protein Charon is a negative regulator of Nodal signaling during left-right patterning in zebrafish.** *Development.* 2004; **131**(8): 1741–1753.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jopling C, van Geemen D, den Hertog J: **Shp2 knockdown and Noonan/LEOPARD mutant Shp2-induced gastrulation defects.** *PLoS Genet.* 2007; **3**(12): e225.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lee HC, Tsai JN, Liao PY, *et al.*: **Glycogen synthase kinase 3 alpha and 3 beta have distinct functions during cardiogenesis of zebrafish embryo.** *BMC Dev Biol.* 2007; **7**: 93.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Liu DW, Hsu CH, Tsai SM, *et al.*: **A variant of fibroblast growth factor receptor 2 (*Fgfr2*) regulates left-right asymmetry in zebrafish.** *PLoS One.* 2011; **6**(7): e21793.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Long S, Ahmad N, Rebagliati M: **The zebrafish nodal-related gene southpaw is required for visceral and diencephalic left-right asymmetry.** *Development.* 2003; **130**(11): 2303–2316.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Monteiro R, van Dinther M, Bakkers J, *et al.*: **Two novel type II receptors mediate**

- BMP signaling and are required to establish left-right asymmetry in zebrafish.** *Dev Biol.* 2008; **315**(1): 55–71.  
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Muto A, Calof AL, Lander AD, *et al.*: **Multifactorial origins of heart and gut defects in nipbl-deficient zebrafish, a model of Cornelia de Lange Syndrome.** *PLoS Biol.* 2011; **9**(10): e1001181.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. Oishi I, Kawakami Y, Raya A, *et al.*: **Regulation of primary cilia formation and left-right patterning in zebrafish by a noncanonical Wnt signaling mediator, duboraya.** *Nat Genet.* 2006; **38**(11): 1316–1322.  
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Ravanelli AM, Klingensmith J: **The actin nucleator Cordon-bleu is required for development of motile cilia in zebrafish.** *Dev Biol.* 2011; **350**(1): 101–111.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
24. Schneider I, Schneider PN, Derry SW, *et al.*: **Zebrafish Nkd1 promotes Dvl degradation and is required for left-right patterning.** *Dev Biol.* 2010; **348**(1): 22–33.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Tian T, Zhao L, Zhang M, *et al.*: **Both foxj1a and foxj1b are implicated in left-right asymmetric development in zebrafish embryos.** *Biochem Biophys Res Commun.* 2009; **380**(3): 537–542.  
[PubMed Abstract](#) | [Publisher Full Text](#)
26. Zhou W, Dai J, Attanasio M, *et al.*: **Nephrocystin-3 is required for ciliary function in zebrafish embryos.** *Am J Physiol Renal Physiol.* 2010; **299**(1): F55–62.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. Ashburner M, Ball CA, Blake JA, *et al.*: **Gene ontology: tool for the unification of biology. The Gene Ontology Consortium.** *Nat Genet.* 2000; **25**(1): 25–29.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Biben C, Harvey RP: **Homeodomain factor Nkx2-5 controls left/right asymmetric expression of bHLH gene eHand during murine heart development.** *Genes Dev.* 1997; **11**(11): 1357–1369.  
[PubMed Abstract](#) | [Publisher Full Text](#)
29. Chen CM, Norris D, Bhattacharya S: **Transcriptional control of left-right patterning in cardiac development.** *Pediatr Cardiol.* 2010; **31**(3): 371–377.  
[PubMed Abstract](#) | [Publisher Full Text](#)
30. Ramsdell AF: **Left-right asymmetry and congenital cardiac defects: getting to the heart of the matter in vertebrate left-right axis determination.** *Dev Biol.* 2005; **288**(1): 1–20.  
[PubMed Abstract](#) | [Publisher Full Text](#)
31. Nakhleh N, Francis R, Giese RA, *et al.*: **High prevalence of respiratory ciliary dysfunction in congenital heart disease patients with heterotaxy.** *Circulation.* 2012; **125**(18): 2232–2242.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. Khodiyar VK, Hill DP, Howe D, *et al.*: **The representation of heart development in the gene ontology.** *Dev Biol.* 2011; **354**(1): 9–17.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. Deegan née Clark JI, Dimmer EC, Mungall CJ: **Formalization of taxon-based constraints to detect inconsistencies in annotation and ontology development.** *BMC Bioinformatics.* 2010; **11**(1): 530.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. Eyre TA, Wright MW, Lush MJ, *et al.*: **HCOP: a searchable database of human orthology predictions.** *Brief Bioinform.* 2007; **8**(1): 2–5.  
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Kent WJ: **BLAT—the BLAST-like alignment tool.** *Genome Res.* 2002; **12**(4): 656–664.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Magrane M, Consortium U: **UniProt Knowledgebase: a hub of integrated protein data.** *Database (Oxford).* 2011; **2011**: bar009.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. Lowe LA, Yamada S, Kuehn MR: **Genetic dissection of nodal function in patterning the mouse embryo.** *Development.* 2001; **128**(10): 1831–1843.  
[PubMed Abstract](#)
38. Khodiyar VK, Dimmer EC, Huntley RP, *et al.*: **Fundamentals of gene ontology functional annotation.** In: *Knowledge-based Bioinformatics.* Edited by Alterovitz G, Ramoni M. Boston, Massachusetts: Wiley; 2010; 171–208.  
[Publisher Full Text](#)
39. Eppig JT, Blake JA, Bult CJ, *et al.*: **The Mouse Genome Database (MGD): comprehensive resource for genetics and genomics of the laboratory mouse.** *Nucleic Acids Res.* 2012; **40**(Database issue): D881–886.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. Binns D, Dimmer E, Huntley R, *et al.*: **QuickGO: a web-based tool for Gene Ontology searching.** *Bioinformatics.* 2009; **25**(22): 3045–3046.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. Srivatsan S, Peng SL: **Cutting edge: Foxj1 protects against autoimmunity and inhibits thymocyte egress.** *J Immunol.* 2005; **175**(12): 7805–7809.  
[PubMed Abstract](#)
42. Pan J, You Y, Huang T, *et al.*: **RhoA-mediated apical actin enrichment is required for cilogenesis and promoted by Foxj1.** *J Cell Sci.* 2007; **120**(Pt 11): 1868–1876.  
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Kim HJ, Schleifarth JR, Jessurun J, *et al.*: **Wnt5 signaling in vertebrate pancreas development.** *BMC Biol.* 2005; **3**: 23.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
44. Nakamura T, Hamada H: **Left-right patterning: conserved and divergent mechanisms.** *Development.* 2012; **139**(18): 3257–3262.  
[PubMed Abstract](#) | [Publisher Full Text](#)
45. MacGrogan D, Nus M, de la Pompa JL: **Notch signaling in cardiac development and disease.** *Curr Top Dev Biol.* 2010; **92**: 333–365.  
[PubMed Abstract](#) | [Publisher Full Text](#)
46. Abu-Issa R, Kirby ML: **Heart field: from mesoderm to heart tube.** *Annu Rev Cell Dev Biol.* 2007; **23**: 45–68.  
[PubMed Abstract](#) | [Publisher Full Text](#)
47. Skarnes WC, Rosen B, West AP, *et al.*: **A conditional knockout resource for the genome-wide study of mouse gene function.** *Nature.* 2011; **474**(7351): 337–342.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
48. van der Linde D, Konings EE, Slager MA, *et al.*: **Birth prevalence of congenital heart disease worldwide: a systematic review and meta-analysis.** *J Am Coll Cardiol.* 2011; **58**(21): 2241–2247.  
[PubMed Abstract](#) | [Publisher Full Text](#)
49. Kosaki K, Bassi MT, Kosaki R, *et al.*: **Characterization and mutation analysis of human LEFTY A and LEFTY B homologues of murine genes implicated in left-right axis development.** *Am J Hum Genet.* 1999; **64**(3): 712–721.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. Kosaki R, Gebbia M, Kosaki K, *et al.*: **Left-right axis malformations associated with mutations in ACVR2B, the gene for human activin receptor type IIB.** *Am J Med Genet.* 1999; **82**(1): 70–76.  
[PubMed Abstract](#) | [Publisher Full Text](#)
51. Britz-Cunningham SH, Shah MM, Zuppan CW, *et al.*: **Mutations of the Connexin43 gap-junction gene in patients with heart malformations and defects of laterality.** *N Engl J Med.* 1995; **332**(20): 1323–1329.  
[PubMed Abstract](#) | [Publisher Full Text](#)
52. Gebbia M, Ferrero GB, Pilia G, *et al.*: **X-linked situs abnormalities result from mutations in ZIC3.** *Nat Genet.* 1997; **17**(3): 305–308.  
[PubMed Abstract](#) | [Publisher Full Text](#)
53. Bataille S, Demoulin N, Devuyst O, *et al.*: **Association of PKD2 (polycystin 2) mutations with left-right laterality defects.** *Am J Kidney Dis.* 2011; **58**(3): 456–460.  
[PubMed Abstract](#) | [Publisher Full Text](#)
54. De Luca A, Sarkozy A, Consoli F, *et al.*: **Familial transposition of the great arteries caused by multiple mutations in laterality genes.** *Heart.* 2010; **96**(9): 673–677.  
[PubMed Abstract](#) | [Publisher Full Text](#)
55. Rankin J, Auer-Grumbach M, Bagg W, *et al.*: **Extreme phenotypic diversity and nonpenetrance in families with the LMNA gene mutation R644C.** *Am J Med Genet A.* 2008; **146A**(12): 1530–1542.  
[PubMed Abstract](#) | [Publisher Full Text](#)
56. Stankunas K, Shang C, Twu KY, *et al.*: **Pbx/Meis deficiencies demonstrate multigenetic origins of congenital heart disease.** *Circ Res.* 2008; **103**(7): 702–709.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
57. Lin X, Xu X: **Distinct functions of Wnt/beta-catenin signaling in KV development and cardiac asymmetry.** *Development.* 2009; **136**(2): 207–217.  
[PubMed Abstract](#) | [Publisher Full Text](#)
58. Slagle CE, Aoki T, Burdine RD: **Nodal-dependent mesendoderm specification requires the combinatorial activities of FoxH1 and Eomesodermin.** *PLoS Genet.* 2011; **7**(5): e1002072.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
59. Johnson DW, Berg JN, Baldwin MA, *et al.*: **Mutations in the activin receptor-like kinase 1 gene in hereditary haemorrhagic telangiectasia type 2.** *Nat Genet.* 1996; **13**(2): 189–195.  
[PubMed Abstract](#) | [Publisher Full Text](#)
60. Trembath RC, Thomson JR, Machado RD, *et al.*: **Clinical and molecular genetic features of pulmonary hypertension in patients with hereditary hemorrhagic telangiectasia.** *N Engl J Med.* 2001; **345**(5): 325–334.  
[PubMed Abstract](#) | [Publisher Full Text](#)
61. Groden J, Thliveris A, Samowitz W, *et al.*: **Identification and characterization of the familial adenomatous polyposis coli gene.** *Cell.* 1991; **66**(3): 589–600.  
[PubMed Abstract](#) | [Publisher Full Text](#)
62. Bakrania P, Efthymiou M, Klein JC, *et al.*: **Mutations in BMP4 cause eye brain, and digit developmental anomalies: overlap between the BMP4 and hedgehog signaling pathways.** *Am J Hum Genet.* 2008; **82**(2): 304–319.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
63. Suzuki S, Marazita ML, Cooper ME, *et al.*: **Mutations in BMP4 are associated with subepithelial, microform, and overt cleft lip.** *Am J Hum Genet.* 2009; **84**(3): 406–411.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
64. Lane KB, Machado RD, Pauciulo MW, *et al.*: **Heterozygous germline mutations in BMP2, encoding a TGF-beta receptor, cause familial primary pulmonary hypertension.** *Nat Genet.* 2000; **26**(1): 81–84.  
[PubMed Abstract](#) | [Publisher Full Text](#)

65. Panizzi JR, Becker-Heck A, Castleman VH, *et al.*: **CCDC103 mutations cause primary ciliary dyskinesia by disrupting assembly of ciliary dynein arms.** *Nat Genet.* 2012; **44**(6): 714–719.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
66. Blanchon S, Legendre M, Copin B, *et al.*: **Delineation of CCDC39/CCDC40 mutation spectrum and associated phenotypes in primary ciliary dyskinesia.** *J Med Genet.* 2012; **49**(6): 410–416.  
[PubMed Abstract](#) | [Publisher Full Text](#)
67. Becker-Heck A, Zohn IE, Okabe N, *et al.*: **The coiled-coil domain containing protein CCDC40 is essential for motile cilia function and left-right axis formation.** *Nat Genet.* 2011; **43**(1): 79–84.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
68. Duquesnoy P, Escudier E, Vincensini L, *et al.*: **Loss-of-function mutations in the human ortholog of *Chlamydomonas reinhardtii* ODA7 disrupt dynein arm assembly and cause primary ciliary dyskinesia.** *Am J Hum Genet.* 2009; **85**(6): 890–896.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
69. Loges NT, Olbrich H, Becker-Heck A, *et al.*: **Deletions and point mutations of LRRC50 cause primary ciliary dyskinesia due to dynein arm defects.** *Am J Hum Genet.* 2009; **85**(6): 883–889.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
70. Reardon W, Winter RM, Rutland P, *et al.*: **Mutations in the fibroblast growth factor receptor 2 gene cause Crouzon syndrome.** *Nat Genet.* 1994; **8**(1): 98–103.  
[PubMed Abstract](#) | [Publisher Full Text](#)
71. Jabs EW, Li X, Scott AF, *et al.*: **Jackson-Weiss and Crouzon syndromes are allelic with mutations in fibroblast growth factor receptor 2.** *Nat Genet.* 1994; **8**(3): 275–279.  
[PubMed Abstract](#) | [Publisher Full Text](#)
72. Wang B, Yan J, Mi R, *et al.*: **Forkhead box H1 (FOXH1) sequence variants in ventricular septal defect.** *Int J Cardiol.* 2010; **145**(1): 83–85.  
[PubMed Abstract](#) | [Publisher Full Text](#)
73. Kott E, Duquesnoy P, Copin B, *et al.*: **Loss-of-function mutations in LRRC6, a gene essential for proper axonemal assembly of inner and outer dynein arms, cause primary ciliary dyskinesia.** *Am J Hum Genet.* 2012; **91**(5): 958–964.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
74. Gillis LA, McCallum J, Kaur M, *et al.*: **NIPBL mutational analysis in 120 individuals with Cornelia de Lange syndrome and evaluation of genotype-phenotype correlations.** *Am J Hum Genet.* 2004; **75**(4): 610–623.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
75. Krantz ID, McCallum J, DeScipio C, *et al.*: **Cornelia de Lange syndrome is caused by mutations in NIPBL, the human homolog of *Drosophila melanogaster* Nipped-B.** *Nat Genet.* 2004; **36**(6): 631–635.  
[PubMed Abstract](#) | [Publisher Full Text](#)
76. Guo J, Cagatay T, Zhou G, *et al.*: **Mutations in the human naked cuticle homolog NKD1 found in colorectal cancer alter Wnt/Dvl/beta-catenin signaling.** *PLoS One.* 2009; **4**(11): e7982.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
77. Bergmann C, Fliegau M, Bruchle NO, *et al.*: **Loss of nephrocystin-3 function can cause embryonic lethality, Meckel-Gruber-like syndrome, situs inversus, and renal-hepatic-pancreatic dysplasia.** *Am J Hum Genet.* 2008; **82**(4): 959–970.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
78. Olbrich H, Fliegau M, Hoefele J, *et al.*: **Mutations in a novel gene, NPHP3, cause adolescent nephronophthisis, tapeto-retinal degeneration and hepatic fibrosis.** *Nat Genet.* 2003; **34**(4): 455–459.  
[PubMed Abstract](#) | [Publisher Full Text](#)
79. Fiskerstrand T, Houge G, Sund S, *et al.*: **Identification of a gene for renal-hepatic-pancreatic dysplasia by microarray-based homozygosity mapping.** *J Mol Diagn.* 2010; **12**(1): 125–131.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
80. Mochizuki T, Wu G, Hayashi T, *et al.*: **PKD2, a gene for polycystic kidney disease that encodes an integral membrane protein.** *Science.* 1996; **272**(5266): 1339–1342.  
[PubMed Abstract](#) | [Publisher Full Text](#)
81. Digilio MC, Conti E, Sarkozy A, *et al.*: **Grouping of multiple-lentiginos/LEOPARD and Noonan syndromes on the PTPN11 gene.** *Am J Hum Genet.* 2002; **71**(2): 389–394.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
82. Tartaglia M, Niemeyer CM, Fragale A, *et al.*: **Somatic mutations in PTPN11 in juvenile myelomonocytic leukemia, myelodysplastic syndromes and acute myeloid leukemia.** *Nat Genet.* 2003; **34**(2): 148–150.  
[PubMed Abstract](#) | [Publisher Full Text](#)
83. Tartaglia M, Mehler EL, Goldberg R, *et al.*: **Mutations in PTPN11, encoding the protein tyrosine phosphatase SHP-2, cause Noonan syndrome.** *Nat Genet.* 2001; **29**(4): 465–468.  
[PubMed Abstract](#) | [Publisher Full Text](#)
84. Renfro DP, McIntosh BK, Venkatraman A, *et al.*: **GONUTS: the Gene Ontology Normal Usage Tracking System.** *Nucleic Acids Res.* 2012; **40**(Database issue): D1262–D1269.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

# Open Peer Review

Current Referee Status:



---

## Version 2

Referee Report 12 February 2014

doi:10.5256/f1000research.3844.r3593



**Jeroen Bakkers**

Cardiac Development and Genetics, Hubrecht Institute, Utrecht, Netherlands

**I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

**Competing Interests:** No competing interests were disclosed.

---

## Version 1

Referee Report 29 January 2014

doi:10.5256/f1000research.2891.r3269



**Jeroen Bakkers**

Cardiac Development and Genetics, Hubrecht Institute, Utrecht, Netherlands

The title and abstract accurately describes the content of the manuscript.

**Method:**

The analysis described here is based on the selected 30 genes from literature that are associated with the GO term 'heart jogging'. According to the description by the authors these genes are involved in zebrafish heart jogging, but a better definition would be that these genes affect heart jogging (either direct or indirect). Indeed some of the genes that were included play a role in cilia function and establishment of the left-right axis.

Many more genes have been identified in zebrafish that affect left-right patterning and thus 'heart jogging'. It remains unclear why these were not included in the search. The observation that many zebrafish left-right genes are not associated with the GO term 'cardiac jogging' could be due to annotation issues (e.g. not every authors uses the term 'jogging' for heart laterality defects in zebrafish embryos). This limitation in the design of this study should be discussed more thoroughly.

In addition the authors could make some conclusions about the usefulness of the GO term 'heart jogging'. Although heart jogging is a process specific to zebrafish it is controlled by a conserved left-right

patterning mechanism. However the GO term 'cardiac jogging' is associated with genes that control heart morphogenesis and left-right patterning.

**I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

**Competing Interests:** No competing interests were disclosed.

Author Response 19 Feb 2014

**Ruth Lovering**, University College London, UK

Dear Jeroen,

Thank you for your comments about our paper.

We have updated the methods, as suggested, to: A list of 30 zebrafish genes that affect heart jogging...

As already described in the methods we only included genes in this list where there was experimental evidence in ZFIN database which confirmed that these genes had an impact on heart jogging in zebrafish. Many genes are likely to be missing from this list because the heart jogging process is often not studied in zebrafish carrying mutations in relevant genes.

We have added the following statement to the methods:

This list does not represent all genes which play a role in heart jogging, as the process of heart jogging is not always studied in zebrafish carrying mutations in relevant genes.

We have not provided a detailed discussion about the GO term 'heart jogging' as we do not want to describe the detail of GO transitivity any further here.

Regards

Ruth

**Competing Interests:** No competing interests were disclosed.

Referee Report 29 November 2013

doi:[10.5256/f1000research.2891.r2480](https://doi.org/10.5256/f1000research.2891.r2480)



**Vincent VanBuren**

Computational Systems Biology, Texas A&M University, Temple, TX, USA

The title is fine, but it would be better if it summarized findings rather than summarized the procedures. The abstract is an adequate summary of the article.

Required revisions:

- **Methods-** Generation of the list of zebrafish jogging genes: "*The search identified a further 23 zebrafish genes, however manual review of these publications led to 5 being disregarded, as the evidence for an involvement in heart jogging was not strong enough.*" The phrase, "*was not strong enough*" does not provide sufficient detail to make construction of this list reproducible. The details should be provided.
- **Discussion-** paragraph 4: "*only a few of the 'jogging ortholog' genes, CCDC103, CCDC40, DNAAF1, LRRC6, NPHP3, and PKD2, are associated with heart looping defects, and thus provide evidence which suggests an involvement of these genes in left-right asymmetry determination in the heart.*" First, making an assertion that genes are not associated with a particular process requires that high-powered studies were performed to reach a negative conclusion. Does the literature support this? Second, while it is true that eliminating one possibility increases the probability of other explanations, it is not very solid evidence for a particular explanation. The logic of the above statement should be better supported, or the statement should be removed.
- **Discussion-** paragraph 4: "*However, there are numerous other reasons why there is a poor association of heart defects with the 'jogging ortholog' gene list.*" This should be, "...numerous other possible reasons...".

**I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

**Competing Interests:** No competing interests were disclosed.

Author Response 19 Feb 2014

**Ruth Lovering**, University College London, UK

Dear Vincent,

Thank you for your helpful comments. We also recognise that since the original search was conducted that there are now an additional 7 other zebrafish genes (Celf1, Cdc14aa, Cdc14b, Dmrt2a, Enpp2, Grem2, Ipar3) with experimental evidence supporting their role in heart jogging ( [Matsui, T. et al., 2012](#); [Clément, A. et al., 2012](#); [Lai, S.L. et al., 2012](#); [Müller, I.I. et al., 2013](#))

As this comment will remain associated with this manuscript we will not add a comment about these additional genes to the manuscript itself, but we will revise the manuscript as follows:

**Methods - Generation of the list of zebrafish jogging genes section:**

To make it clearer how this list of zebrafish jogging genes was generated we will replace part of this section with the following:

*"A list of 30 zebrafish heart jogging genes was compiled using a variety of approaches. Twelve zebrafish proteins were identified as they were already annotated to the 'heart jogging' GO terms, the remaining 18 proteins were then identified using the ZFIN (<http://zfin.org/>) Site Search, with the search phrase 'heart jogging', and filtering using the 'Expression/Phenotypes' category. This search retrieves figures from papers that have 'heart jogging' in the figure legend, and thus are likely to be describing specific zebrafish genes (and proteins) involved in this process. Many of these genes had not yet been curated with GO terms. Each of the papers identified in this way*

*were reviewed; of the 23 zebrafish genes identified in these papers five (Bmpr1aa, Tbx1, unnm\_hu119, unnm\_hu202, unnm\_hu304) were eliminated, as none of these papers provided experimental evidence for the involvement of these genes in heart jogging."*

**In Discussion - paragraph 4:**

We did not intend to make a negative conclusion here, this statement was making a positive statement that some of the 'jogging ortholog' genes, CCDC103, CCDC40, DNAAF1, LRRC6, NPHP3, and PKD2, are associated with heart looping defects. And as included in the comment below we do state that 'there are numerous other reasons why there is a poor association of heart defects with the 'jogging ortholog' gene list'. To make this statement less controversial, we will modify it as follows:

*"Mutations in several human genes have now been identified as causative of abnormal heart looping, such as ACVR2B, LEFTY2, GJA1 and ZIC3(49–52), and some of the 'jogging ortholog' genes (CCDC103, CCDC40, DNAAF1, LRRC6, NPHP3 and PKD2) are also associated with heart looping defects. Thus providing evidence to support an involvement of these genes in left-right asymmetry determination in the heart."*

**Discussion- paragraph 4 (2<sup>nd</sup> comment):**

As suggested we will revise this to "*However, there are numerous other possible reasons why there is a poor association of heart defects with the 'jogging ortholog' gene list.*"

**Competing Interests:** No competing interests were disclosed.

---