

## Developing fencing policies for dryland ecosystems

Durant, S.M., Becker, M.S., Creel, S., Bashir, S., Dickman, A.J., Beudels-Jamar, R.C., Lichtenfeld, L., Hilborn, R., Wall, J., Wittemyer, G., Badamjav L., Blake, S., Boitani, L., Breitenmoser, C., Broekhuis, F., Christianson, D., Cozzi, G., Davenport, T.R.B., Deutsch, J., Devillers, P., Dollar, L., Dolrenry, S., Douglas-Hamilton, I., Dröge, E., FitzHerbert, E., Foley, C., Hazzah, L., Hopcraft, J.G.C., Ikanda, D., Jacobson, A., Joubert, D., Kelly, M.J., Milanzi, J., Mitchell, N., M'Soka, J., Msuha, M., Mweetwa, T., Nyahongo, J., Rosenblatt, E., Schuette, P., Sillero-Zubiri, C., Sinclair, A.R.E., Stanley-Price, M. R., Zimmermann, A., Pettorelli, N.

1. Badamjav L., CMS Scientific Council & Mongolian Academy of Sciences Jukov Avenue 77 Iaanbaatar 51 MONGOLIA
2. Bashir, S., Birdlife International, Singapore
3. Becker, M.S., Zambian Carnivore Programme, Box 80, Mfuwe, Eastern Province Zambia; and Conservation Biology and Ecology Program, 310 Lewis Hall, Department of Ecology, Montana State University, Bozeman, MT 59717 USA
4. Beudels-Jamar, R.C., Royal Belgian Institute of Natural Sciences, 29 Vautier str., 1000 Bruxelles & CMS Scientific Council, UNEP/CMS, Hermann-Ehlers-Str. 10, 53113 Bonn, Germany
5. Blake, S., Max Planck Institute for Ornithology, Whitney R. Harris World Ecology Center, Washington University in St. Louis, USA
6. Boitani, L., Department of Biology and Biotechnologies, Università di Roma La Sapienza, Viale Università 32, 00185 - Roma, Italy
7. Breitenmoser, C., Co-Chair IUCN/SSC Cat Specialist Group, c/o KORA, Thunstrasse 31, 3074 Muri, Switzerland
8. Broekhuis, F., Mara Cheetah Project, Kenya Wildlife Trust, P.O. Box 86, 00502 Karen, Nairobi, Kenya & Wildlife Conservation Research Unit, Department of Zoology, University of Oxford, The Recanati-Kaplan Centre, Tubney House, Tubney OX13 5QL, UK
9. Christianson, D.A., School of Natural Resources and the Environment, University of Arizona, Tucson, AZ, 85721, USA
10. Cozzi, G., Institute of Evolutionary Biology and Environmental Studies, Zurich University, Winterthurerstrasse 190, CH – 8057 Zürich, Switzerland
11. Creel, S., Department of Ecology, Montana State University, Bozeman, MT 59717, USA.
12. Davenport, T.R.B., Wildlife Conservation Society, Tanzania Program, PO Box 922, Zanzibar, Tanzania
13. Deutsch, J., Wildlife Conservation Society, Bronx Zoo, 2300 Southern Blvd., Bronx, NY, 10460, USA
14. Devillers, P., Royal Belgian Institute of Natural Sciences, 29 Vautier str., 1000 Bruxelles & CMS Scientific Council, UNEP/CMS, Hermann-Ehlers-Str. 10, 53113 Bonn, Germany
15. Dickman, A.J., Wildlife Conservation Research Unit, Department of Zoology, University of Oxford, The Recanati-Kaplan Centre, Tubney House, Tubney OX13 5QL, UK
16. Dollar, L., Big Cats Initiative, National Geographic Society, 1145 17th Street NW, Washington D.C., 20036-4688, USA; Department of Biology; Pfeiffer University; Misenheimer, North Carolina USA; and Nicholas School of the Environment; Duke University; Durham, North Carolina USA
17. Dolrenry, S., Lion Guardians, PO Box 15550 Langata, Kenya 00509
18. Douglas-Hamilton, I., Save the Elephants, PO Box 54667, Nairobi, Kenya; Department of Zoology, University of Oxford, Oxford OX1 3PS, United Kingdom
19. Dröge, E., Zambian Carnivore Programme, Box 80, Mfuwe, Eastern Province Zambia
20. Durant, S.M., Institute of Zoology, Zoological Society of London, Regents Park, London NW1 4RY, UK & Wildlife Conservation Society, Bronx Zoo, 2300 Southern Blvd., Bronx, NY, 10460, USA

21. FitzHerbert, E., Wildlife Conservation Research Unit, Department of Zoology, University of Oxford, The Recanati-Kaplan Centre, Tubney House, Tubney OX13 5QL, UK
22. Foley, C., Wildlife Conservation Society Tanzania Program, PO Box 922, Zanzibar, Tanzania
23. Hazzah, L., Lion Guardians, PO Box 15550, Langata, Kenya 00509
24. Hilborn, R., School of Aquatic and Fishery Sciences, University of Washington, Seattle, WA 98195 USA
25. Hopcraft, J.G.C., Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow, University Avenue, G12 8QQ, UK
26. Ikanda, D., Tanzania Wildlife Research Institute, Box 661, Arusha, Tanzania
27. Jacobson, A., Institute of Zoology, Zoological Society of London, Regents Park, London NW1 4RY, UK
28. Joubert, D., Big Cats Initiative, National Geographic Society, 1145 17th Street NW, Washington D.C., 20036-4688, USA
29. Kelly, M.J., Dept. of Fish & Wildlife Conservation, Virginia Tech; 146 Cheatham Hall, Blacksburg, VA 24061-0321; USA
30. Lichtenfeld, L., African People & Wildlife Fund, PO Box 624, Bernardsville, NJ 07924 USA
31. Milanzi, J., Regional Manager, Western Region, Zambia Wildlife Authority, Private Bag 1, Chilanga, Zambia
32. Mitchell, N., Conservation Programmes, Zoological Society of London, Regents Park, London NW1 4RY, UK & Wildlife Conservation Society, Bronx Zoo, 2300 Southern Blvd., Bronx, NY, 10460, USA
33. M'Soka, J., Department of Ecology, Conservation Biology and Ecology Program, Montana State University, 310 Lewis Hall, Bozeman, MT, 59717, USA and Zambia Wildlife Authority, Private Bag 1, Chilanga, Zambia, Zambian Carnivore Programme, Box 80, Mfuwe, Eastern Province Zambia
34. Msuha, M., Tanzania Wildlife Research Institute, Box 661, Arusha, Tanzania
35. Mweetwa, T., Zambian Carnivore Programme, Box 80, Mfuwe, Eastern Province Zambia; School of Natural Resources and the Environment, University of Arizona, Tucson, AZ, 85721, USA
36. Nyahongo, J., University of Dodoma, Dodoma, Tanzania
37. Pettorelli, N., Institute of Zoology, Zoological Society of London, Regents Park, London NW1 4RY, UK
38. Rosenblatt, E., Zambian Carnivore Programme, Box 80, Mfuwe, Eastern Province Zambia; and Conservation Biology and Ecology Program, 310 Lewis Hall, Department of Ecology, Montana State University, Bozeman, MT 59717 USA
39. Schuette, P., Zambian Carnivore Programme, Box 80, Mfuwe, Eastern Province Zambia; and Conservation Biology and Ecology Program, 310 Lewis Hall, Department of Ecology, Montana State University, Bozeman, MT 59717 USA
40. Sillero-Zubiri, C., Chair, IUCN/SSC Canid Specialist Group & Wildlife Conservation Research Unit, Department of Zoology, University of Oxford, The Recanati-Kaplan Centre, Tubney House, Tubney OX13 5QL, UK,
41. Sinclair, A.R.E., Beaty Biodiversity Research Centre, University of British Columbia, 6270 University Boulevard, Vancouver V6T 1Z4, Canada
42. Stanley-Price, M. R., Wildlife Conservation Research Unit, Department of Zoology, University of Oxford, The Recanati-Kaplan Centre, Tubney House, Tubney OX13 5QL, UK
43. Wall, J., Lab for Advanced Spatial Analysis, Department of Geography, University of British Columbia, 1984 West Mall, Vancouver B.C., Canada, V6T 1Z2
44. Wittemyer, G., Colorado State University, Fish, Wildlife and Conservation Biology, Fort Collins, Colorado, USA
45. Zimmermann, A., Wildlife Conservation Research Unit, Department of Zoology, University of Oxford, The Recanati-Kaplan Centre, Tubney House, Tubney OX13 5QL, UK & Chester Zoo, Chester CH2 1LH, UK

**Running head:** Fencing policies for dryland ecosystems

**Key-words:** migration, desert, rangelands, barriers, nomadic pastoralism, biodiversity conservation, ecosystem function, transhumance, management interventions, conservation policy

## Abstract

1. In dryland ecosystems, mobility is essential for both wildlife and people to access unpredictable and spatially heterogeneous resources, particularly in the face of climate change. Fences may prevent connectivity vital for this mobility.
2. There are recent calls for large-scale barrier fencing interventions to address human–wildlife conflict and illegal resource extraction. Fencing has costs and benefits to people and wildlife. However, the evidence available for facilitating sound decision-making for fencing initiatives is limited, particularly for drylands.
3. We identify six research areas that are key to informing evaluations of fencing initiatives: economics; edge permeability; reserve design; connectivity; ecosystem services; and communities.
4. *Policy implications.* Implementing this research agenda to evaluate fencing policies for dryland ecosystems will enable better management and policy decisions. The United Nations Conventions on Migratory Species (CMS) and to Combat Desertification (UNCCD) are appropriate policy frameworks for moving this agenda forward and leading the development of policies and guidelines on fencing in drylands.

## **A resurgence in calls for large scale fencing interventions in Africa**

Fencing has been used worldwide for a variety of purposes, including protecting remnant wildlife populations from overhunting, poaching or invasive species and reducing human–wildlife conflict and human encroachment (Somers & Hayward 2012). In Africa, after a proliferation of fencing initiatives in the 1960s and 1970s, there has been a recent resurgence in calls for large-scale fencing to protect biodiversity, and to separate wildlife from people, livestock and crops. For example, Uganda intends to fence all of its national parks in a bid to stem human–wildlife conflict (Government of Uganda 2012). The Rwandan authorities recently erected a 120-km fence around the Akagera National Park at a capital cost of \$2.5 million in a bid to eliminate human–wildlife conflict (Hall 2013). Meanwhile the government of Malawi has stated a wish to protect all parks in the country with electric fences (Kafemveka 2013).

In stark contrast, elsewhere in Africa, authorities are removing fences to restore wildlife populations and migratory movements and to promote wildlife-based economies for conservancies and local communities. The Southern African Development Community (SADC) in the *Phakalane Declaration* has recently recommended strategic realignment of veterinary cordon fences (erected for wildlife disease control) to counteract the harmful impacts of fences on wildlife populations (SADC 2012). In addition, the non-governmental organisation (NGO)-led Transfrontier Conservation Area and privately-led conservancy movements across Africa are encouraging the widespread removal of fencing to re-establish large scale animal movements (WCS 2008; Lindsey, Romañach & Davies-Mostert 2009; Van Aarde & Jackson 2007). Their aim is to support or restore wide-ranging species whose populations are no longer viable in small reserves.

Scientific opinion on the topic of fencing is equally divided. A recent analysis of African lion *Panthera leo* densities and growth rates from fenced and unfenced populations concluded that fencing was a cost-effective conservation strategy for this species, and recommended fencing as a primary conservation tool for lions (Packer *et al.* 2013). However, Creel *et al.* (2013) demonstrated

that the studied populations differed in key aspects other than fencing, with fenced populations having markedly higher budgets for substantially smaller areas, which often held intensively managed populations maintained well above carrying capacity. In concert these factors confounded analyses and a reanalysis with the same data using population size, rather than carrying capacity, found the opposite result: that many more lions are conserved per dollar invested in unfenced ecosystems, while also avoiding the ecological and economic costs of fencing (Creel *et al.* 2013). This debate prompted a subsequent article in *Science* that highlighted the problems associated with large-scale fencing, and concluded that, as climate change increases the importance of wildlife mobility and landscape connectivity, fencing of wildlife should become an action of last resort (Woodroffe, Hedges & Durant 2014).

To reconcile such widely divergent opinions and contradictory policies, we review and identify key information needs for conservation policy-makers and practitioners for better assessment of costs and benefits of proposed fencing interventions. Critical evaluation of fencing initiatives is most urgent in the world's dryland ecosystems<sup>1</sup> where mobility is essential for both wildlife and people to access temporally variable and spatially heterogeneous resources (Notenbaert *et al.* 2012). In such landscapes, the erection of large-scale impermeable barriers may reduce connectivity and lead to significant ecological and economic impacts (Okin *et al.* 2009).

Drylands cover 41% of the land's surface and are home to an estimated 64% of all bird, 55% of mammal and 25% of amphibian species (Davies *et al.* 2012). They support some of the world's largest populations of terrestrial megafauna and significant wildlife migratory systems (Harris *et al.* 2009; Milner-Gulland, Fryxell & Sinclair 2011). Moreover, two billion people live in drylands, including some of the most vulnerable and marginalized communities in the world (Middleton *et al.* 2011). Survival of wildlife and people in these arid lands has depended on adapting to a harsh and

---

<sup>1</sup> We define drylands as those areas with an aridity index value of less than 0.65, in accordance with the Millennium Ecosystem Assessment (Safriel *et al.* 2005; Fig. 1), the United Nations Environmental Programme (UNEP), the International Union for the Conservation of Nature (IUCN) and the Convention on Biological Diversity (CBD) (Davies *et al.* 2012).

highly variable environment, characterized by short growing seasons and low, unpredictable rainfall that are not conducive to agriculture. Historically, people living in drylands depended on nomadic or semi-nomadic pastoralism, a strategy that allows the most efficient use of highly variable and localized rainfall (McCabe 2004; Homewood 2009). Thus, in unpredictable dryland environments, mobility is critical to access transient forage and water resources for both wildlife and people (Notenbaert *et al.* 2012).

### **Costs and benefits of fencing in dryland ecosystems**

Fences are free-standing structures aimed at restricting or preventing movement across boundaries (Hayward & Kerley 2009). Fences are usually erected to reduce threats to wildlife from direct human activities (such as ecosystem degradation, harvesting, persecution and disturbance); reduce conflict between people and wildlife (Lindsey *et al.* 2012); and reduce disease transmission between wildlife and domestic animals, most notably the extensive veterinary barrier fences stretching across southern Africa (Gadd 2012). Fencing is widely used in Australian drylands to exclude invasive non-native species from wildlife areas, though the maintenance and construction costs incurred in building fences able to exclude small invasive predators generally keep such fenced areas relatively small (Dickman 2012).

While we can relatively easily identify the potential benefits, the negative consequences of large-scale fencing interventions may be less obvious. Large-scale fencing can disrupt migration pathways and reduce access to key areas within drylands, such as seasonal foraging areas (Harris *et al.* 2009), and wetland refuges (Davies *et al.* 2012). This can lead to severe reductions in migratory or nomadic ungulate populations and may prompt wider impacts on non-migratory species (Harris *et al.* 2009; Gadd 2012). Some impacts may occur over a long time, which makes them particularly difficult to detect (Norrdahl *et al.* 2002). Fencing also restricts ranging of keystone species, such as African elephants *Loxodonta Africana*, which significantly influence ecosystem structure and function (Shrader, Pimm & Van Aarde 2010; Asner & Levick 2012). The potentially damaging habitat

impacts arising from 'compressing' elephants within protected areas have been well documented (Western & Gichohi 1993; Douglas-Hamilton, Krink & Vollrath 2005; Loarie, Van Aarde & Pimm 2009). Similarly, fencing may also cause disruption at high trophic levels, such as altering the population dynamics or restricting movement of top predators, which is likely to lead to cascading impacts, loss of ecosystem function and impoverished biodiversity (Estes *et al.* 2011). Implementing intensive and expensive management to mitigate against such effects, such as translocation or anthropogenic control of population size (e.g. see discussion of lions in Packer *et al.* 2013; Creel *et al.* 2013), may not be feasible or cost effective, particularly for multiple species, and is unlikely to provide an adequate replacement for naturally regulated and connected ecosystems.

### **Keeping wildlife in or people out?**

A well-constructed, well-maintained fence can be wildlife-proof, but can never be human-proof. Even the most heavily-fortified fences have not prevented the illegal killing of white rhinoceros *Ceratotherium simum* and black rhinoceros *Diceros bicornis* in South Africa over recent years (Woodroffe, Hedges & Durant 2014). People are likely to be able to circumvent any fence, but they may also destroy fences in order to gain access to useful resources on the other side of the fence, such as bushmeat, ivory, honey, medicinal plants and grazing. In doing so, they may make the fence permeable to wildlife, and sometimes wildlife may not be able to find their way back through the fence. The fence itself may also serve as a readily available source of snare wire, rendering a fence erected to protect wildlife from bushmeat extraction counterproductive (Lindsey *et al.* 2011; Lindsey *et al.* 2013; Becker *et al.* 2013). Alternative fencing materials, such as kinked mesh wire, can reduce this risk, but they are not well known to local management agencies, difficult to source and more expensive. Thus they are less likely to be adopted, particularly in government fencing programmes that may be focused more on protecting people than wildlife.

A fence can reduce human–wildlife conflict, but may also prevent people from accessing benefits from nature and adversely impact the development of community-based incentives for

wildlife conservation (East *et al.* 2012; Gadd 2012). Moreover, a fence may also contribute to the loss of coping strategies that have enabled communities to coexist with wildlife. Thus, if a fence, after erection, is lost or breached, human–wildlife conflict may reach levels much higher than those that existed prior to the establishment of the fence (Gadd 2012). Hence, it is critical that, once erected, a fence is maintained as an impermeable barrier. Wildlife often inflict small breaches in a fence, necessitating frequent and costly ongoing maintenance to sustain its effectiveness as a barrier (Kesch, Bauer & Loveridge 2013; Lindsey *et al.* 2009). Thus, the initial capital construction costs are only a small part of the investment required.

### **Developing an evidence-base to evaluate dryland fencing interventions**

Scientific understanding of the costs and benefits of fences is still in its infancy (Somers & Hayward 2012), and is currently inadequate to support sound policy-making. Here we identify six research areas where incomplete or poor information hinders the wise use of fencing (Table 1). For the purposes of this discussion, we consider perimeter fencing of reserves, but our analysis is relevant for other large-scale fencing interventions, such as the increasing use of fencing to safeguard oil or gas pipelines and transport networks.

- (1) *Economics*. Economic costs form the basis for many conservation policies, but we still know very little about the ability of different conservation interventions, including fencing, to deliver conservation success for a given cost (McCreless *et al.* 2013). This makes it very difficult to assess the relative expenditure to benefit ratio of fencing against other alternative interventions (Possingham *et al.* 2001). Yet the economic assessment of fencing is fundamental to sound policy decisions since limited conservation resources must be spent wisely to deliver sustainable solutions and maximize conservation impact. The only economic analyses conducted on the efficacy of fencing do not control for the apportioning of the overall budget to other reserve management activities (i.e. Packer *et al.* 2013; Creel *et al.* 2013), and only the most well-financed reserves are able to afford fencing interventions. Thus,



it is not possible to disentangle the benefits of fencing from those of other investments such as anti-poaching efforts, community engagement, infrastructure investment and other activities that potentially confound the effect of fences on the effectiveness of a reserve and the density of a focal species. Without such an analysis, it is impossible to ascertain whether a budget increase, which allows fencing interventions and subsequent management, would deliver better outcomes for conservation and communities compared with investing the same funds in other reserve management strategies, such as community engagement and anti-poaching, without any fencing. A proper comparison of alternative strategies using long-term data and metrics of conservation success must include short-term capital costs, which can be considerable for fencing, as well as recurring maintenance costs.

- (2) *Edge permeability.* Fencing an already existing abrupt transition (i.e. “hard edge”) between a reserve and the surrounding anthropogenically-modified landscape can be part of the justification for fencing interventions. Fencing of such habitat edges prevents the movement of wildlife beyond the reserve, where they might forage in crops or kill livestock. A presumed “hard edge” suggests that negative impacts on wildlife from the fence due to restriction in movement will be minimal since the surrounding modified landscape is often viewed as comprising marginal habitat. Yet, the actual permeability of the edge will be species- and system-specific, as well as context-specific (Ries & Sisk 2010). Understanding what constitutes a hard edge for different species in the context of overall conservation and management objectives of fencing interventions is necessary to assess whether a “hard edge” justification is appropriate.
- (3) *Reserve design.* A landscape perspective on fencing implementation is critical as the impacts of a fence on wildlife, ecosystems and communities depends on its location relative to the broader ecological context (Soule & Terborgh 1999). Dryland protected areas often have boundaries delineated by key resources that are often shared by wildlife and humans, such as

major rivers that are accessed by both humans and wildlife. Fencing interventions to separate people and wildlife may result in a barrier preventing access to the resource for wildlife, or preventing access to human communities, or both. Productive agricultural land or key habitats, such as wetlands, may also border reserves (Watson *et al.* 2013) and play key roles for ecosystems and local communities. The impacts of fencing in relation to the design of reserve boundaries, and how to mitigate these impacts, need to be better understood.

(4) *Connectivity.* Connectivity is fundamental to the long-term viability of many wildlife populations, particularly migratory and nomadic species common to dryland systems. Dryland reserves often do not cover the entire extent of an animal's range, and may often be placed in either dry or wet season ranges for migratory or nomadic species (Fynn & Bonyongo 2011). In such situations, perimeter fencing of reserves, preventing access to critical seasonal resources, can lead to collapse in the populations of these species (Gadd 2012). Moreover, access to ephemeral resources may also be critical to the long-term survival of some species. For example, foraging or water resources in key areas outside a reserve may be important for the survival of long-lived species, such as elephants, during extreme climatic events (Foley, Pettoirelli & Foley 2008). Identification of those species that are most vulnerable to reserve isolation and developing a clear understanding as to what constitutes connectivity for such species is key to evaluating the ecological impacts of fencing interventions.

(5) *Ecosystem services.* Beyond the specifics of the reserve site and design, there is also a need to better understand how the delivery of ecosystem services (e.g. soil and watershed protection, timber, plant and animal harvesting, etc.) is compromised or enhanced by fencing initiatives. Given the large-scale ecological processes that characterize dryland systems and the dependence of people and wildlife on them, it is unlikely that fencing will have no impact on ecosystem service delivery and access. Indeed, studies show that simply subdividing land in drylands can substantially reduce overall grazing carrying capacity (Boone *et al.* 2005). Soil-

based ecosystem services, such as nutrient recycling and water capture, are particularly vulnerable to degradation in drylands (Parr *et al.* 1990), yet there is no information on how these services may be impacted by fencing. An understanding is needed as to how fencing might hinder or help meet a reserve's overall biodiversity conservation goals and the continued delivery of ecosystem services, as well as how this may be modified by climate change.

- (6) *Human communities.* Many protected areas permit some limited access for local communities, and some of the poorest and most marginalized members of communities may be particularly dependent on natural resources from these areas (Loibooki *et al.* 2002; Brashares *et al.* 2011). Fencing interventions are likely to make legitimate access more difficult, and risk marginalizing these individuals still further. Local communities are heterogeneous; some individuals may suffer the costs of wildlife, in the form of crop and livestock depredation for example, while others may benefit from wildlife through tourism and hunting revenue or associated ecosystem services (Thompson & Homewood 2002), and hence the costs and benefits of fencing interventions are likely to be unevenly distributed between households. While it is important that conservation interventions maintain the integrity of reserves, they should avoid contributing to or exacerbating existing inequities within communities. A better understanding of the socioeconomic impacts of fencing is needed to avoid such unintended consequences on local communities.

The information from these six major research areas is key to a proper evaluation of fencing interventions. Evaluations need to be carefully undertaken in the context of the aims of the proposed fencing intervention. For example, fences designed to keep wildlife in versus those meant to keep people out, are two substantially different objectives, which in turn will likely have variable success and impacts. Any evaluation also needs to be undertaken in the context of the overall

management goals for each reserve; rarely, for example, are such goals focused on a single species as per the analyses of Packer *et al.* (2013).

While the information required for these evaluations may appear extensive, in reality, many of these research areas can be addressed by collating and analysing existing information, or by implementing targeted monitoring and evaluation of new fencing interventions (Table 1). For example, most protected areas have documented expenditure reports; measures of reserve design are available from the protected area database; and remote sensing data can be used to delineate edge permeability and monitor the delivery of some key ecosystem services (Avanu *et al.* 2012). There are, however, some areas where additional information is required. For example, while research areas such as ecosystem service delivery may be measured using remote sensing data, there are others, such as wildlife abundance or species diversity, which require direct sampling. There is also a need for improvements in our understanding of movement patterns, and what constitutes barriers to movement, for many wide-ranging wildlife species. Such information could be provided through fitting satellite or GPS collars to target species. Regardless of the availability of ecological data, a clear information gap is the socioeconomic impacts of reserves on local human communities, and there is a clear need for detailed socioeconomic studies on people living close to fenced and unfenced wildlife areas.

### **Towards policy guidelines on large-scale fencing interventions for drylands**

It is clear that fences erected to protect wildlife or people can be a useful conservation tool, but can also be counterproductive. Guidelines, which take into account species-specific requirements, ecological conditions and human communities would help conservation practitioners better evaluate large-scale fencing interventions. The United Nations Convention on Migratory Species (CMS) is ideally suited to lead such guideline development, given the CMS's focus on wide-ranging species, experience with fencing as a management tool, and recognized expertise in conservation action for arid areas (e.g. CMS 2011).

The United Nations Convention to Combat Desertification (UNCCD), because of its mandate for sustainable management of drylands, is also well placed to engage with the breadth of the proposed research agenda. The UNCCD is one of three treaties developed from the United Nations Earth Summit in 1992 and aims to prevent and reverse land degradation and to mitigate the effects of drought, and is particularly relevant to developing countries, where most drylands are located. As well as CMS and UNCCD, the Food and Agriculture Organisation of the United Nations (FAO) also has an important role to play in the sustainable management of drylands.

Neither the CMS or UNCCD currently provide general policy guidelines as to the use of large-scale fencing, nor does the FAO. Better understanding of the impacts of fencing interventions would facilitate the development of appropriate policies to help communities and governments to improve sustainable management of drylands. Developing policies and guidelines for assessing when, where, and the type of fencing that should, or should not, be used in drylands, would help to prevent a repeat of the past harm done by fences to people, wildlife and ecosystems. Preventing further degradation is likely to require solutions within an integrated landscape approach to conservation that acknowledges local communities as part of the ecosystems (IIED 2013).

Many large-scale fencing interventions are likely to impact multiple countries, hence it may also be useful to make use of regional economic structures, such as the Southern African Development Community (SADC), East African Community (EAC), West African Economic and Monetary Union (UEMOA) and South Asian Association for Regional Cooperation (SAARC), and target bilateral and multilateral donors, to enforce guidelines and to help promote the need for full environmental impact assessments (EIA). These structures could also be used to ensure that all large-scale fencing interventions have a practical and achievable long-term maintenance and financing plan to guarantee the long-term integrity of the barrier once established. We recommend active engagement of these organisations in contributing to the improvement of knowledge of the impacts of fencing in drylands and in the development and implementation of policy guidelines.

Despite the high capital costs, fencing can initially appear to be an easy solution. Yet, unless fencing strategies have local community support and a financing plan to meet the expensive long-term costs of fence maintenance, there is a danger that they may generate more problems than they solve. The research agenda proposed will generate information necessary for better evaluation of fencing interventions that take into account the full range of likely impacts in dryland systems. Ultimately there is a need for funding agencies to increase support for these areas and their marginalized peoples, and develop better management strategies to sustain dryland ecosystems (Mortimore *et al.* 2009). The CMS and UNCCD could help to prevent further degradation of these important systems by leading global efforts to develop an understanding of the impacts of large-scale fencing interventions in drylands and establishing guidelines to regulate their use.

**Acknowledgements:** We thank Stuart Pimm and John Fryxell who provided constructive comments on the manuscript.

#### **Data Accessibility**

Data have not been archived because this article does not contain data.

## References

- Asner, G.P. & Levick, S.R. (2012) Landscape-scale effects of herbivores on treefall in African savannas. *Ecology Letters*, **15**, 1211-1217.
- Ayanu, Y.Z., Conrad, C., Nauss, T., Wegmann, M. & Koellner, T. (2012) Quantifying and Mapping Ecosystem Services Supplies and Demands: A Review of Remote Sensing Applications. *Environmental Science & Technology* **46**, 8529-8541.
- Becker, M., McRobb, R., Watson, F., Droge, E., Kamyembo, B., Murdoch, J., & Kakumbi, C. (2013) Evaluating wire-snare poaching trends and the impacts of by-catch on elephants and large carnivores. *Biological Conservation*, **158**, 26-36.
- Boone, R.B., BurnSilver, S.B., Thornton, P.K., Worden, J.S. & Galvin, K.A. (2005) Quantifying declines in livestock due to land subdivision. *Rangeland Ecology & Management*, **58**, 523-532.
- Brashares, J.S., Golden, C.D., Weinbaum, K.Z., Barrett, C.B. & Okello, G. V. (2011) Economic and geographic drivers of wildlife consumption in rural Africa. *Proceedings of the National Academy of Sciences of the United States of America*, **108**, 13931–13936.
- CMS (2011) Barriers to migration: case study in Mongolia. Ulaanbaator, available from [http://www.cms.int/sites/default/files/publication/Inf\\_23\\_Migration\\_Barriers\\_WWF\\_Mongolia\\_E.pdf](http://www.cms.int/sites/default/files/publication/Inf_23_Migration_Barriers_WWF_Mongolia_E.pdf).
- Creel, S., Becker, M.S., Durant, S.M., M'Soka, J., Matandiko, W., Dickman, A.J., Christianson, D., Dröge, E., Mweetwa, T., Pettorelli, N., Rosenblatt, E., Schuette, P., Woodroffe, R., Bashir, S., Beudels-Jamar, R.C., Blake, S., Borner, M., Breitenmoser, C., Broekhuis, F., Cozzi, G., Davenport, T.R.B., Deutsch, J., Dollar, L., Dolrenry, S., Douglas-Hamilton, I., Fitzherbert, E., Foley, C., Hazzah, L., Henschel, P., Hilborn, R., Hopcraft, J.G.C., Ikanda, D., Jacobson, A., Joubert, B., Joubert, D., Kelly, M.S., Lichtenfeld, L., Mace, G.M., Milanzi, J., Mitchell, N., Msuha, M., Muir, R., Nyahongo, J., Pimm, S., Purchase, G., Schenck, C., Sillero-Zubiri, C., Sinclair, A.R.E., Songorwa, A.N., Stanley-Price, M., Tehou, C.A., Trout, C., Wall, J., Wittemyer,

- G. & Zimmermann, A. (2013) Conserving large populations of lions – the argument for fences has holes. *Ecology Letters*, **16**, 1413-e3.
- Davies, J., Poulsen, L., Schulte-Herbrüggen, B., Mackinnon, K., Crawhall, N., Henwood, W.D., Dudley, N., Smith, J. & Gudka, M. (2012) Conserving Dryland Biodiversity. IUCN (International Union for the Conservation of Nature) [www.iucn.org/publications](http://www.iucn.org/publications).
- Dickman, C.R. (2012) Fences or ferals? Benefits and costs of conservation fencing in Australia. *Fencing for Conservation: Restriction of Evolutionary Potential or a Riposte to Threatening Processes?* (eds M.J. Somers & M.W. Hayward), pp. 43-64. Springer, New York.
- Douglas-Hamilton, I., Krink, T. & Vollrath, F. (2005) Movements and corridors of African elephants in relation to protected areas. *Naturwissenschaften* **92**, 158-163.
- East, M.L., Nyahongo, J.W., Goller, K.V. & Hofer, H. (2012) Does the vastness of the Serengeti limit human-wildlife conflicts? *Fencing for Conservation: Restriction of Evolutionary Potential or a Riposte to Threatening Processes?* (eds M.J. Somers & M.W. Hayward), pp. 125-151. Springer, New York.
- Estes, J. A, Terborgh, J., Brashares, J.S., Power, M.E., Berger, J., Bond, W.J., Carpenter, S.R., Essington, T.E., Holt, R.D., Jackson, J.B.C., Marquis, R.J., Oksanen, L., Oksanen, T., Paine, R.T., Pickett, E.K., Ripple, W.J., Sandin, S. a, Scheffer, M., Schoener, T.W., Shurin, J.B., Sinclair, A.R.E., Soulé, M.E., Virtanen, R. & Wardle, D. A. (2011) Trophic downgrading of planet Earth. *Science*, **333**, 301–306.
- Foley, C., Pettorelli, N. & Foley, L. (2008) Severe drought and calf survival in elephants. *Biology Letters*, **4**, 541-544.
- Fynn, R.W.S. & Bonyongo, M.C. (2011) Functional conservation areas and the future of Africa's wildlife. *African Journal of Ecology*, **49**, 175-188.
- Gadd, M.E. (2012) Barriers, the beef industry and unnatural selection: A review of the impact of veterinary fencing on mammals in southern Africa. *Fencing for Conservation: Restriction of*



- Evolutionary Potential or a Riposte to Threatening Processes?* (eds M.J. Somers & M.W. Hayward), pp. 153-186. Springer, New York.
- Government of Uganda (2012) UWA to fence off protected areas.  
<http://www.visituganda.com/information-centre/media/news/archive/?id=51>
- Hall, S. (2013) Launch of the Akagera fence by the Rwanda Development Board. African Park Latest News. African Parks, [www.african-parks.org](http://www.african-parks.org).
- Harris, G., Thirgood, S., Hopcraft, J.G.C., Crooms, J.P.G.M. & Berger, J. (2009) Global decline in aggregated migrations of large terrestrial mammals. *Endangered Species Research*, **7**, 55-76.
- Hayward, M.W. & Kerley, G.I.H. (2009) Fencing for conservation: Restriction of evolutionary potential or a riposte to threatening processes? *Biological Conservation*, **142**, 1-13.
- Homewood, K. (2009) *Ecology of African pastoralist societies*. Ohio University Press, Athens.
- IIED (2013) Global public policy narratives on the drylands and pastoralism. International Institute for the Environment and Development. <http://pubs.iied.org/pdfs/10040IIED.pdf>
- Kafemveka, M. (2013) Malawi government to hand over Liwonde, Nkhotakota Game Reserves. Malawi Voice, 29<sup>th</sup> September, <http://www.malawivoice.com/2013/09/29/malawi-govt-to-hand-over-liwonde-nkhotakota-game-reserves/>
- Kesch, K.M., Bauer, D.T. & Loveridge, A.J. (2013) Undermining game fences: who is digging holes in Kalahari sands? *African Journal of Ecology*, **52**, 144-150.
- Lindsey, P.A., Romañach, S.S. & Davies-Mostert, H.T. (2009) The importance of conservancies for enhancing the value of game ranch land for large mammal conservation in southern Africa. *Journal of Zoology*, **277**, 99-105.
- Lindsey, P.A., Romañach, S.S., Matema, S., Matema, C., Mupamhadzi, I. & Muvengwi, J. (2011) Dynamics and underlying causes of illegal bushmeat trade in Zimbabwe. *Oryx*, **45**, 84-95.
- Lindsey, P.A., Masterson, C.L., Beck, A.L. & Romanach, S. (2012) Ecological, social and financial issues related to fencing as a conservation tool in Africa. *Fencing for Conservation: Restriction of*

- Evolutionary Potential or a Riposte to Threatening Processes?* (eds M.J. Somers & M.W. Hayward), pp. 215-234. Springer, New York.
- Lindsey, P.A., Balme, G., Becker, M., Begg, C., Bento, C., Bocchino, C., Dickman, A., Diggle, R.W., Eves, H., Henschel, P., Lewis, D., Marnewick, K., Mattheus, J., Weldon McNutt, J., McRobb, R., Midlane, N., Milanzi, J., Morley, R., Murphree, M., Opyene, V., Phadima, J., Purchase, G., Rentsch, D., Roche, C., Shaw, J., Westhuizen, H.v.d., Vliet, N.V. & Zisadza-Gandiwa, P. (2013) The bushmeat trade in African savannas: Impacts, drivers, and possible solutions. *Biological Conservation*, **160**, 80-96.
- Loarie, S.R., Van Aarde, R.J. & Pimm, S.L. (2009) Fences and artificial water affect African savannah elephant movement patterns. *Biological Conservation*, **142**, 3086-3098.
- Loibooki, M., Hofer, H., Campbell, K.L.I. & East, M.L. (2002) Bushmeat hunting by communities adjacent to the Serengeti National Park, Tanzania: the importance of livestock ownership and alternative sources of protein and income. *Environmental Conservation*, **29**, 391-398.
- McCabe, J.T. (2004) *Cattle Bring Us to Our Enemies: Turkana Ecology, Politics, and Raiding in a Disequilibrium System*. University of Michigan Press, Ann Arbor.
- McCreless E., Visconti P., Carwardine J. & Wilcox C, Smith R.J. (2013) Cheap and nasty? The potential perils of using management costs to identify global conservation priorities. *PLoS ONE* **8**, e80893
- Middleton, N., Stringer, L., Goudie, A. & Thomas, D. (2011) The forgotten billion: MDG Achievement in the Drylands. United Nations Office at Nairobi, Publishing Services Section, ISO 14001:2004.
- Milner-Gulland, E.J., Fryxell, J.M. & Sinclair, A.R.E. (2011) *Animal migration: A synthesis*. Oxford University Press, Oxford.
- Mortimore, M., Anderson, S., Cotula, L., Davies, J., Faccar, K., Hesse, C., Morton, J., Nyangena, W., Skinner, J. & Wolfangel, C. (2009) Dryland Opportunities: A new paradigm for people,

ecosystems and development. IUCN, Gland, Switzerland; IIED, London, UK and UNDP/DDC, Nairobi, Kenya.

Notenbaert, An M.O., Davies, J, De Leeuw, J, Said, M, Herrero, M, Manzano, P, Waithaka, M, Aboud, A, & Omondi, S (2012) Policies in support of pastoralism and biodiversity in the heterogeneous drylands of East Africa. *Pastoralism: Research Policy and Practice* **2**, 14. (doi:10.1186/2041-7136-2-14).

Norrdahl, K., Klemola T., Korpimäki, E. & Koivula, M. (2002) Strong seasonality may attenuate trophic cascades: vertebrate predator exclusion in boreal grassland. *Oikos* **99**, 419–430.

Okin, G. S., Parsons, A.J., Wainwright, J., Herrick, J.E., Bestelmeyer, B.T., Peters, D.C., & Fredrickson, E. L. (2009) Do Changes in Connectivity Explain Desertification? *BioScience* **59**, 237-244.

Packer, C., Loveridge, A., Canney, S., Caro, T., Garnett, S.T., Pfeifer, M., Zander, K.K., Swanson, A., Macnulty, D., Balme, G., Bauer, H., Begg, C.M., Begg, K.S., Bhalla, S., Bissett, C., Bodasing, T., Brink, H., Burger, A., Burton, A.C., Clegg, B., Dell, S., Delsink, A., Dickerson, T., Dloniak, S.M., Druce, D., Frank, L., Funston, P., Gichohi, N., Groom, R., Hanekom, C., Heath, B., Hunter, L., Deiongh, H.H., Joubert, C.J., Kasiki, S.M., Kissui, B., Knocker, W., Leathem, B., Lindsey, P.A., MacLennan, S.D., McNutt, J.W., Miller, S.M., Naylor, S., Nel, P., Ng'weno, C., Nicholls, K., Ogutu, J.O., Okot-Omoya, E., Patterson, B.D., Plumptre, A., Salerno, J., Skinner, K., Slotow, R., Sogbohossou, E.A., Stratford, K.J., Winterbach, C., Winterbach, H. & Polasky, S. (2013) Conserving large carnivores: dollars and fence. *Ecology Letters* **16**, 635-641.

Parr, J.F., Stewart B.A., Hornick, S.B. & Singh, R.P. (1990) Improving the Sustainability of Dryland Farming Systems: A Global Perspective. *Advances in Soil Science* (eds R.P. Singh, J.F. Parr & B.A. Stewart), pp. 1-8. Springer, New York.

Possingham, H.P., Andelman, S., Noon, B.R., Trombulak, S. & Pulliam, H.R. (2001) Making smart conservation decisions. *Conservation biology: research priorities for the next decade* (eds M.E. Soulé & G. Orians), pp. 225-244. Island Press, Washington D.C..

- Ries, L. & Sisk, T.D. (2010) What is an edge species? The implications of sensitivity to habitat edges. *Oikos*, **119**, 1636–1642.
- SADC (2012) The Phakalane Declaration. [http://www.wcs-ahead.org/phakalane\\_declaration.html](http://www.wcs-ahead.org/phakalane_declaration.html)
- Safriel, U., Adeel, Z., Niemeijer, D., Puigdefabres, J., White, R., Lal, R., Winslow, M., Ziedler, J., Prince, S., Archer, E., King, C., (2005) Dryland systems. *Ecosystems and human well-being: Current state and trends, vol. 1.* (eds R. Hassan, R., Scholes & N. Ash), pp. 623–662. Island Press, Washington, Covelo, London.
- Shrader, A.M., Pimm, S.L. & Van Aarde, R.J. (2010) Elephant survival, rainfall and the confounding effects of water provision and fences. *Biodiversity and Conservation*, **19**, 2235-2245.
- Somers, M.J. & Hayward, M.W. (2012) Fencing for Conservation: Restriction of Evolutionary Potential or a Riposte to Threatening Processes? (eds), Springer, New York.
- Soulé M.E. & Terborgh J. (1999) Continental conservation: scientific foundations of regional reserve networks. Island Press, Washington D. C
- Thompson, M. & Homewood K. (2002) Entrepreneurs, elites, and exclusion in maasailand: Trends in wildlife conservation and pastoralist development. *Human Ecology* **30**, 107-138.
- Van Aarde, R.J. & Jackson T.P. (2007) Megaparks for metapopulations: Addressing the causes of locally high elephant numbers in southern Africa. *Biological Conservation*, **134**, 289-297.
- Watson, F., Becker, M.S., McRobb, R. & Kanyembo B. (2013) Spatial Patterns of Wire-Snare Poaching: Implications for Community Conservation in National Park Buffer Zones. *Biological Conservation*, **168**, 1-9.
- WCS (2008) As the fences come down. Available from <http://www.wcs-ahead.org/documents/asthefencescomedown.pdf>. Accessed 18 June 2013.
- Western, D. & Gichohi, H. (1993) Segregation effects and the impoverishment of savanna parks: the case for ecosystem viability analysis. *African Journal of Ecology*, **31**, 269-281.
- Woodroffe, R., Hedges, S. & Durant, S.M. (2014) To fence or not to fence. *Science*, **344**, 46–8.

**Table 1** Evidence needs and potential data that can be used to evaluate fencing interventions in drylands

<b>Research issue</b>	<b>Question</b>	<b>Evidence needs</b>	<b>Data</b>
Economics	Is fencing a cost effective and sustainable approach to deliver conservation success?	How do different conservation activities compare with fencing under similar operating budgets, bearing in mind the substantial capital and maintenance costs required for fencing?  Is fencing economically more or less sustainable than other options?	Reserve expenditure reports broken down by management activity  Data on fence integrity  Measures of conservation success
Edge permeability	Does the boundary to be fenced constitute a hard edge?	What is the definition of a hard edge and how does this vary between species and ecosystems?	Remote sensing data  Wildlife movement and distribution data on the edge of protected areas
Reserve design	How does the reserve's design impact the costs and benefits of fencing?	What is the impact of reserve shape?  If reserve boundaries lie on key landscape features and resources, such as rivers, then how will this mediate the balance between costs and benefits of fencing for wildlife and people?	Protected area database  Remote sensing data  Geographic information system (GIS) layers  Wildlife movement and distribution data in relation to landscape and resources  Contribution of resources to local livelihoods
Connectivity	How important is connectivity to the overall goals of the reserve and ecosystem function?	How important are wildlife movements into and out of reserves to their population viability?  How does a fence affect wildlife movements and does it prevent wildlife from accessing key resources?  Which species are most vulnerable to reserve isolation?  What constitutes connectivity for these species?  What are the impacts from the restriction of wildlife movement due to fencing on ecosystem function?	Remote sensing data  GIS layers  Wildlife movement data in fenced and unfenced areas  Wildlife habitat and resource use data inside and outside the reserve  Map of potential barriers to movement  Map of potential areas of connectivity  Measures of immigration and emigration for wide-ranging wildlife  Life history and survivorship data for wide-ranging and dispersing species
Ecosystem services	How does the establishment of a fence impact delivery of ecosystem services?	What is the relationship between habitat subdivision and carrying capacity? How does fencing affect delivery of ecosystem services?  If the fence is to entirely enclose a reserve – how will this affect the viability of low density and wide ranging species within the reserve? (If such species require intensive management then this should be included in the economic costing of the fencing intervention)  How is climate change likely to affect ecosystem resilience and how is this likely to be impacted by fencing?  How does fencing affect the interactions between ecosystem service delivery and rainfall and productivity?	Remote sensing data  GIS layers  Protected area database  Wildlife surveys  Demographic data and population viability modelling  Climate data and climate change predictions
Communities	What are the benefits and costs to local communities of fencing and how are these distributed between individuals?	What legal and illegal benefits do the communities derive from the presence of the reserve?  How are these benefits distributed within the community?  What are the costs to communities from the presence of the reserve?  How are these costs distributed within the community?  How will fencing affect these costs and benefits?  Who is likely to benefit from fencing and by how much, and who is likely to pay the costs and by how much?	Game scout and ranger patrol reports  Resource extraction data  Socioeconomic data from local communities around fenced and unfenced reserves including:  Wealth and livelihoods  Distribution of resources  Costs and benefits from wildlife

Fig. 1 The world's dryland zones based on an aridity index  $<0.65$  (Safriel *et al.* 2005).

