

## Image of the month: Bilateral gigantic earlobe keloids

**Authors:** Filipe Brogueira Rodrigues,<sup>A</sup> Catarina Bekerman,<sup>B</sup> João Sousa,<sup>C</sup> Joana Vieira<sup>D</sup> and Vitor Ramalhinho<sup>E</sup>



**Fig 1. (a) Right and (b) left earlobes.**

A 90-year-old woman of African descent was admitted to our department. Stiff, painless, multilobulated, earlobe masses were noted bilaterally (Fig 1). The lesions appeared after ear piercing during childhood and progressed further after a surgical resection attempt 20 years ago. Biopsy wasn't attempted due to risk of progression. The clinical diagnosis of keloids was established and the patient was offered a dermatology follow-up. ■

**Authors:** <sup>A</sup>MD, Department of Medicine, Hospital de Santa Maria, Lisbon, Portugal; <sup>B</sup>MD, Department of Medicine, Hospital de Santa Maria, Lisbon, Portugal; <sup>C</sup>MD, Department of Medicine, Hospital de Santa Maria, Lisbon, Portugal; <sup>D</sup>MD, Department of Medicine, Hospital de Santa Maria, Lisbon, Portugal; <sup>E</sup>MD, Department of Medicine, Hospital de Santa Maria, Lisbon, Portugal

**Address for correspondence:** Dr FB Rodrigues, Department of Medicine, Hospital de Santa Maria, Centro Hospitalar de Lisboa Norte, Av Professor Egas Moniz, 1649-035 Lisboa, Portugal.

**Email:** [filipebrodrigues@gmail.com](mailto:filipebrodrigues@gmail.com)