



Mitochondrial DNA Part B Resources

ISSN: (Print) 2380-2359 (Online) Journal homepage: <http://www.tandfonline.com/loi/tmdn20>

The complete mitochondrial genome of *Epomophorus gambianus* (Chiroptera: Pteropodidae) and its phylogenetic analysis

Silke A. Riesle-Sbarbaro, Stefan P. W. de Vries, Samuel Stubbs, Kofi Amponsah-Mensah, Andrew A. Cunningham, James L. N. Wood & David R. Sargan

To cite this article: Silke A. Riesle-Sbarbaro, Stefan P. W. de Vries, Samuel Stubbs, Kofi Amponsah-Mensah, Andrew A. Cunningham, James L. N. Wood & David R. Sargan (2016) The complete mitochondrial genome of *Epomophorus gambianus* (Chiroptera: Pteropodidae) and its phylogenetic analysis, *Mitochondrial DNA Part B*, 1:1, 447-449, DOI: [10.1080/23802359.2016.1181993](https://doi.org/10.1080/23802359.2016.1181993)

To link to this article: <http://dx.doi.org/10.1080/23802359.2016.1181993>



© 2016 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 08 Jul 2016.



Submit your article to this journal [↗](#)



Article views: 37



View related articles [↗](#)



View Crossmark data [↗](#)

Full Terms & Conditions of access and use can be found at
<http://www.tandfonline.com/action/journalInformation?journalCode=tmdn20>

The complete mitochondrial genome of *Epomophorus gambianus* (Chiroptera: Pteropodidae) and its phylogenetic analysis

Silke A. Riesle-Sbarbaro^{a,b}, Stefan P. W. de Vries^a, Samuel Stubbs^a, Kofi Amponsah-Mensah^c, Andrew A. Cunningham^b, James L. N. Wood^a and David R. Sargan^a

^aDepartment of Veterinary Medicine, University of Cambridge, Cambridge, UK; ^bZoological Society of London, Institute of Zoology, London, UK;

^cDepartment of African Center for Wetlands Conservation, University of Legon, Accra, Ghana

ABSTRACT

The Gambian epauletted fruit bat, *Epomophorus gambianus*, is widely distributed across sub-Saharan Africa. Its assembled and annotated mitochondrial genome (GenBank accession no. KT963027) is 16,702 bases in length, containing 13 protein-coding genes, 22 transfer RNA genes, two ribosomal RNA genes and two non-coding regions: the control region (D-loop) and the origin of light-strand replication (O_L). The average base composition is 32.2% A; 27.6% C; 14% G; and 26.1% T. The mitogenome presented a structural composition greatly conserved between members of the Pteropodidae family.

ARTICLE HISTORY

Received 11 March 2016
Revised 18 April 2016
Accepted 20 April 2016

KEYWORDS

Chiroptera; *Epomophorus gambianus*; mitogenome; phylogeny

Epomophorus gambianus is an old world fruit bat, part of one of the most diversified taxa: the family Pteropodidae (Simmons 2005). Throughout its distribution, *Epomophorus* co-roost with several Pteropodid bats; including members of Rousettini, Myonycterini and Epomorphorini tribes (Nesi et al. 2013). With increasing number of Chiropteran studies exploring their role as disease reservoirs (Calisher et al. 2006; Baker et al. 2012), it is imperative to resolve their evolutionary relationships accurately.



Using the DNeasy Blood and Tissue Kit (Qiagen, Valencia, CA), DNA was extracted from wing membrane biopsies from five *E. gambianus* captured in Ghana (6°59'32.12"N; 0°25'32.12"E) and stored in the University of Cambridge, UK. A DNA sequencing library was constructed and sequenced on an Illumina MiSeq (Supplementary methods). A total of 27,143,463 paired-end reads were mapped with CLC Genomics Workbench v7.5.1 software (CLC Bio, Aarhus, Denmark) to *Rousettus aegyptiacus* (GenBank AB205183.1). The resulting assembly (16,624 bp) was used as reference for *de novo* assembly, presenting 98% coverage against the reference mitochondrion. Decreasing coverage was observed downstream of bp position 15,900. The lowest coverage (< 7x) was found within D-loop, in a region that overlapped with a perfect repeat (CATACACGTACG)₂₃. Sanger sequencing


was used for gap closure (Supplementary Table 1); the Sanger sequences were mapped to the assembly using GENEious v8.1 (CLC Bio, Aarhus, Denmark). The resulting assembly was annotated by referencing nine closely related mt-genomes (Supplementary Table 2).

Most of the genes are encoded on the heavy strand of the mitogenome with the exception of eight tRNA and ND6 protein-coding gene that are on the light strand. Except for NADH dehydrogenase 1, 2, 3 and 5, ATG was the starting codon. TAA was the stop codon except for gene ND2 ending with TAG, and *Cyt b* with AGG. ND1, COX III and ND4 presented incomplete stop codons (Supplementary Table 3).

To evaluate the evolutionary relationship of *E. gambianus* to other Chiroptera, eighteen mitogenomes were aligned using Mega6 (Tamura et al. 2013). Three methods were used to produce phylogenetic trees and the resulting topologies were visualized with FigTree v1.4.2 (CLC Bio, Aarhus, Denmark). Node supports were assembled in a consensus tree with TreeGraph 2.4.0-456 beta (Stover & Muller 2010) and manually verified and merged in Adobe Illustrator (Figure 1).

The presented assembly is in agreement with previous classification of the species and corroborate the Yinpterochiroptera/Yangochiroptera division (Teeling et al.

CONTACT Ms Silke A. Riesle-Sbarbaro  sasr3@cam.ac.uk  University of Cambridge, Department of Veterinary Medicine, Madingley road, Cambridge CB3 0ES, United Kingdom of Great Britain and Northern Ireland; Zoological Society of London, Institute of Zoology, London NW1 4RY, UK

 Supplemental data for this article can be accessed [here](#).

© 2016 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Phylogenetic tree of complete bat mitogenomes. Three methods are shown: maximum parsimony (MP) created using a Min-mini heuristic algorithm and 1000 bootstrap replicates; maximum-likelihood (ML) constructed in PAUP 4.0 (Swofford 2011) with bootstrap percentages computed after 1000 replicates using PhyML software (Guindon et al. 2010); and Bayesian Inference performed in MrBayes 3.1.2 (Huelsenbeck & Ronquist 2001) using MCMC algorithm, running six chains for 300,000 generations and discarding the first 25% of trees. The best-fit nucleotide substitution model for ML and Bayesian methods were selected with JmodelTest 2.1.7 (Darriba et al. 2012) and PAUP 4.0 beta software. Bayesian posterior probabilities appear above and left to the node it supports. Bootstrap values appear below in bold text for ML and in yellow for MP, values inside square brackets refer to conflicting topology. *Epomophorus gambianus* is denoted with a bat illustration. *Canis lupus* complete mt-genome was assigned as an out-group.

2005; Nesi et al. 2013). *Epomophorus gambianus* mitogenome could prove useful for resolving introgressive hybridizations, a phenomenon that has been recorded between this species and the sympatric *Micropteropus pusillus* (Nesi et al. 2011).

Disclosure statement

The authors report that they have no conflicts of interest.

Funding information

This study was supported by a Becas-Chile (CONICYT) scholarship; and the European Union FP7 project ANTIGONE under Grant (number 278976).

References

- Baker KS, Todd S, Marsh G, Fernandez-Loras A, Suu-Ire R, Wood JL, Wang LF, Murcia PR, Cunningham AA. 2012. Co-circulation of diverse paramyxoviruses in an urban African fruit bat population. *J Gen Virol*. 93:850–856.
- Calisher CH, Childs JE, Field HE, Holmes KV, Schountz T. 2006. Bats: important reservoir hosts of emerging viruses. *Clin Microbiol Rev*. 19:531–545.
- Darriba D, Taboada GL, Doallo R, Posada D. 2012. Jmodeltest 2: more models, new heuristics and parallel computing. *Nat Methods*. 9:772
- Guindon SDJF, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of phylml 3.0. *Syst Biol*. 59:307–321.
- Huelsenbeck JP, Ronquist F. 2001. Mrbayes: Bayesian inference of phylogenetic trees. *Bioinformatics*. 17:754–755.
- Nesi N, Kadjo B, Pourrut X, Leroy E, Pongombo Shongo C, Craud C, Hassanin A. 2013. Molecular systematics and phylogeography of the tribe myonycterini (mammalia, pteropodidae) inferred from mitochondrial and nuclear markers. *Mol. Phylogenet Evol*. 66:126–137.
- Nesi N, Nakoune E, Craud C, Hassanin A. 2011. DNA barcoding of african fruit bats (mammalia, pteropodidae). The mitochondrial genome does not provide a reliable discrimination between *Epomophorus gambianus* and *Micropteropus pusillus*. *C R Biol*. 334:544–554.
- Simmons NB. 2005. Order chiroptera. In: Wilson DE, Reeder DM, editors. *Mammal species of the world: a taxonomic and geographic reference*. 3rd ed. US: John Hopkins University Press. p. 312–529.

- Stover BC, Muller KF. 2010. Treegraph 2: combining and visualizing evidence from different phylogenetic analyses. *BMC Bioinform.* 11:7.
- Swofford DL. 2011. Paup*: phylogenetic analysis using parsimony, version 4.0b10.
- Tamura K, Stecher G, Peterson D, Filipksi A, Kumar S. 2013. Mega6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol.* 30:2725–2729.
- Teeling EC, Springer MS, Madsen O, Bates P, O'Brien SJ, Murphy WJ. 2005. A molecular phylogeny for bats illuminates biogeography and the fossil record. *Science.* 307:580–584.