

The vestibular body: vestibular contributions to bodily representations

Journal:	<i>Cognitive Neuropsychology</i>
Manuscript ID	CN-RV 84.15.R2
Manuscript Type:	Review
Keywords:	Vestibular System, Multisensory Integration, Body Representation

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Review article for Cognitive Neuropsychology
Special Issue on Body Representations

**The vestibular body:
vestibular contributions to bodily representations**

For Peer Review Only

Abstract

Uniquely among the sensory modalities, vestibular inputs do not project to any primary unimodal cortex. Rather vestibular signals are integrated with signals from other sensory modalities, such as vision, touch and proprioception. This convergence could reflect an important mechanism for maintaining the perception of the body. Neuropsychological, electrophysiological and psychophysical studies show support for this hypothesis. Here we review the current literature in order to develop a framework for understanding how the vestibular system contributes to body representation. According to recent models, we distinguish between three processes for body representation, and we look at whether vestibular signals might influence each process. These are (i) *somatosensation*, the primary sensory processing of somatic stimuli, (ii) *somatoperception*, the processes of constructing percepts and experiences of somatic objects and events and (iii) *somatorepresentation*, the knowledge about the body as a physical object in the world. Vestibular signals appear to contribute to all three levels in this model of body processing. Thus, the traditional view of the vestibular system as a low-level, dedicated orienting module tends to underestimate the pervasive role of vestibular input in bodily self-awareness.

Abbreviations:

OP: Parietal Operculum

PIVC: Parieto-Insular-Vestibular Cortex

SEPs: Somatosensory Evoked Potentials

LEPs: Laser Evoked Potentials

BIID: Body Identity Integrity Disorder

Keywords:

Vestibular system; multisensory integration; body representation.

Introduction

Our body is the object we know best. Not only **do we** have more information on our body compared to any other objects, but we also have a privileged access to our own bodily sensations that we cannot have for any other bodies or objects (Eilan et al., 1998). It might not be surprising **therefore** that few issues in cognitive neuropsychology and experimental psychology have encountered such interest as how the brain represents our body. Current research has accumulated a large body of evidence showing that coherent body representations are generated by the successful integration of multimodal sensory signals (Blanke and Metzinger, 2009; Blanke, 2012). Strong emphasis has been traditionally put on the two classical signals which tell us about the body: vision and touch. **Only recently have studies highlighted the essential role played by our sixth sense, the vestibular system, in the perception and representation of the body.**

The vestibular system provides continuous information to the brain to update body position and to maintain orientation in the surrounding space (Berthoz, 1996). The semicircular canals detect rotational movements of the head in three-dimensional space (i.e., pitch, yaw and roll) and the otolith organs (utricle and saccule) code translational acceleration, including the orientation of the head relative to the gravitational vertical. These signals are highly relevant to several interactions between the organism and the environment. In particular, dynamic vestibular inputs from the canals are associated with low-level visuo-vestibular interactions to control gaze and eye fixation, while static gravitational inputs from the otolith organs contribute to path integration and navigation. Accordingly, vestibular peripheral dysfunction leads to disintegration in the normal relation between personal and extra-personal space (Blanke et al., 2004), confirming a vestibular contribution to the brain's *Global Positioning System*.

Vestibular signals are *somatic* in two distinct senses. First, they continuously detect movement and acceleration of the head and the body. Second, vestibular signals may also provide a more general experience of our body. Vestibular receptors are directly sensitive to gravitational accelerations, and they have been shown to play a crucial role in sensing the

1
2
3
4 vertical direction of gravity (Böhmer and Mast, 1999; Bronstein, 1999; Mittelstaedt, 1991;
5 Mittelstaedt, 1992; Mittelstaedt, 1999; Snyder, 1999; Barnett-Cowan et al., 2015). The
6 perception of verticality is based on the integration of somatic information generated by
7 vestibular receptors in the inner ear with signals from the muscles, joints, skin, and viscera
8 (Lackner, 1988; Lackner and DiZio, 2005; Mittelstaedt, 1992; see Lacquaniti et al. 2015 for a
9 review). The well-known *Aubert effect* is a clear example of such a vestibular-somatic relation
10 (Anastasopoulos et al., 1999; Yardley, 1990): when a person is roll-tilted, and deprived of
11 relevant visual cues about their orientation, the subjective perception of gravitational vertical is
12 biased towards the body axis. That is, the online perception of verticality seems to include a
13 strong prior that the body is upright, as well as sensory afferent information. A stronger bias
14 towards the body midline could be taken as evidence for a weaker sensory verticality signal
15 input, so the contributions of priors and sensory evidence to verticality perception can be
16 measured. Several studies have shown that the online sensory evidence in the Aubert situation
17 is not only vestibular, but also somatosensory. For instance, case reports showed that patients
18 with entire sensory loss and patients suffering from hemisensory loss lying on their impaired
19 side did not exhibit the normal Aubert effect when tilted in the way that normally induces such
20 effect (Anastasopoulos et al., 1999; Yardley, 1990). This suggests that vestibular inputs are
21 strongly combined with other somatic signals even for the processing of percepts, such as
22 verticality, that are classically considered purely vestibular.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 Uniquely among the sensory modalities, vestibular inputs do not project to any
44 unimodal cortex. Multimodal convergence has been described in almost all vestibular relays,
45 including the vestibular nuclei, the thalamus and several areas in the cerebral cortex (zu
46 Eulenburg et al., 2012; Lopez et al., 2012a). Electrophysiological studies have identified a
47 widespread vestibular network whose core area is the Parieto-Insular-Vestibular Cortex (PIVC).
48 This area lies in the posterior parietal operculum extending into the posterior insular lobe
49 (Guldin and Grüsser, 1998). The human homologue of the primate PIVC is a distributed set of
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 regions including the posterior and anterior insula, temporo-parietal junction, superior temporal
5 gyrus, inferior parietal lobule, and somatosensory cortices (see Lopez et al., 2012a for a review).
6
7
8 This network can be divided in three main functional components (Balaban et al., 2011). First,
9
10 a *sensorimotor component* which includes pathways for the sensory integration and perceptual
11 responses of vestibular, visual, proprioceptive and somatosensory information and for both
12 somatic and visceral motor response mechanisms (Balaban and Yates, 2004). Second, a
13
14 *cognitive component* which includes pathways for regulation of affect, decision making,
15 attention and higher cognitive behaviours. Third, an *interoceptive component* which includes
16 pathways for the integration of information regarding ongoing sensory processes relative to the
17
18 current physiological condition of the body (Balaban et al., 2011).
19
20
21
22
23

24
25 Several neuropsychological observations have further suggested that vestibular
26 information underpins a strong link between the sense of self and awareness of one's own body.
27
28 Individuals with peripheral vestibular disorders frequently report misperception of the shape,
29
30 size, and location of their body parts (Sang et al., 2006; Lopez et al., 2008). Altered bodily
31 representations have been described following artificial vestibular stimulation in neurological
32 patients. For instance, artificial vestibular stimulation induces transient remission of
33
34 disownership for body parts (*somatoparaphrenia*, Bisiach et al., 1991), somatoagnosia (Rode et
35 al., 2012) and anosognosia (Cappa et al., 1987) caused by brain damage. Additionally, artificial
36
37 vestibular stimulation modifies phantom limb sensations in both paraplegic (Le Chapelain et al.,
38
39 2001) and amputee patients (André et al., 2001). More dramatically, vestibular dysfunction
40
41 may contribute to autoscopic phenomena such as the impression of seeing one's own body in
42
43 extrapersonal space (Blanke et al., 2005).
44
45
46
47
48

49
50 Here we review both neuropsychological observations and experimental results in healthy
51
52 volunteers to argue that vestibular signals are fundamental for achieving a coherent
53
54 representation of the body. Following Longo et al. (2010), we distinguished between three
55
56 hierarchically-arranged levels of body representation, and we looked at whether vestibular
57
58 signals might contribute to each of them (Figure 1). These are (i) somatosensation: "primary
59
60

1
2
3
4 sensory processing of somatic stimuli”, (ii) somatoperception: “*the processes of constructing*
5
6 *percepts and experiences of somatic objects and events, and of one's own body*” and (iii)
7
8 somatopresentation: knowledge about “*the body's character as a physical object in the*
9
10 *external world*” (Longo et al., 2010). Thus, somatosensation refers to the primary processing of
11
12 bodily sensations, while somatoperception involves an online percept of the state of the body, or
13
14 of an object interacting with the body, based on the integration of multisensory inputs.
15
16 Somatopresentation refers to both explicit and implicit knowledge about the body, which may
17
18 be less directly related to immediate input. The organisation of the components of this model is
19
20 hierarchical: somatosensation is a necessary condition for somatoperception, because perceiving
21
22 the current state of the body always involves integrating a range of somatosensory afferent
23
24 signals. In contrast, somatopresentation can persist in the absence of any particular
25
26 somatosensory input: one knows what one’s body is like even in the absence of any strong
27
28 sensory input, for example after prolonged immobility, or while floating in water.

29
30 Longo et al.’s (2010) model highlights a duality in two classes of higher-order
31
32 representations about the body. On the one hand, somatoperception refers to the *perceptual*
33
34 process of constructing perceptual representations of the body. On the other,
35
36 somatopresentation refers to the *cognitive* process of constructing abstract knowledge about
37
38 the body as a physical object (Longo et al., 2010). These processes differ in timescale as well as
39
40 in their functions: somatoperception is online and short-lasting, while representations of the
41
42 body may be retained over the long-term, potentially up to lifespan durations: many
43
44 autobiographies demonstrate compelling long-term retention of first-person accounts of body
45
46 knowledge. Merleau-Ponty, for example, alludes to this enduring quality of body knowledge:
47
48 “We have found underneath the objective and detached knowledge of the body that other
49
50 knowledge which we have of it in virtue of its always being with us and of the fact that we are
51
52 our body” (Merleau-Ponty, 1962, p. 206).

53
54
55 However, interactions between body representations might occur. Longo et al. (2010)
56
57 suggested that these interactions were primarily horizontal, between distinct types of
58
59
60

1
2
3
4 information at each level. For instance, tactile localization and body posture represent distinct
5
6 and dissociable somatoperceptual representations (see below), yet there is evidence supporting
7
8 functional interactions between them (Ho and Spence, 2007; Medina and Rapp, 2008). Vertical
9
10 interactions between somatoperception and somatopresentation are less likely to occur: for
11
12 example, the ability to recognise one's own arm as belonging to one's own body
13
14 (somatopresentation) should be independent of the current posture of the arm
15
16 (somatoperception). However, as neuropsychological evidence highlights, some between-level
17
18 interactions exist. Nevertheless, it is plausible to hypothesise that vertical interactions and
19
20 horizontal interactions require different computational principles. For instance, horizontal
21
22 within-level interactions might be based on multisensory processes. Conversely, between-level
23
24 vertical interactions might reflect cognitive hierarchical processes. Critically, Longo et al.'s
25
26 (2010) model allows us to consider how vestibular signals *independently* contribute to low-level
27
28 signal processing and to higher-order representations of the body. We suggest that each level of
29
30 this somatic hierarchy can be influenced by vestibular signals (Figure 1).
31
32
33

34 **Vestibular contributions to somatosensation**

35
36 Somatosensation refers to the primary sensory processing of somatic stimuli, which
37
38 principally occurs in the primary somatosensory cortices (Kaas, 1983; Mountcastle, 2005;
39
40 Iwamura, 1998; Johnson and Hsiao, 1992; Romo and Salinas, 2001). For instance, it involves
41
42 the ability to consciously detect and recognise somatosensory stimuli on the skin surface.
43
44

45
46 The influence of the vestibular system on somatosensation was first described by Vallar
47
48 and colleagues in 1990 (Vallar et al., 1990), who investigated the effects of caloric vestibular
49
50 stimulation on visuo-spatial hemineglect and associated somatosensory deficits. Caloric
51
52 vestibular stimulation (CVS) is a routine diagnostic technique in the assessment of vestibular
53
54 peripheral functionality (see Grabherr et al., 2015 for a review). This stimulation requires the
55
56 irrigation of cold or warm water into the external ear canal (hereafter we refer to it as cold water
57
58 irrigation, unless otherwise indicated). CVS elicits a movement of the semicircular canal fluid,
59
60

1
2
3
4 stimulating the afferent vestibular nerve which transmits the signals to the vestibular nuclei and
5
6 activates the contralateral cortical and subcortical structures. Vallar et al. (1990) reported a
7
8 transient amelioration of somatosensory deficits (*hemianaesthesia*) and many manifestations of
9
10 the syndrome of visuo-spatial hemineglect in three right brain-damaged patients immediately
11
12 after CVS (Vallar et al., 1990). This observation was replicated in a group study including both
13
14 right brain-damaged and left brain-damaged patients (Vallar et al., 1993). Critically, the mirror-
15
16 reversed paradigm, i.e. right ear CVS in right hemianaesthesia was unsuccessful: recovery of
17
18 left hemianaesthesia was observed only in right brain-damaged patients following left ear CVS.
19
20 The stimulation was generally ineffective in left brain-damaged patients, though it was effective
21
22 in left brain-damaged patients presenting with visuo-spatial hemineglect (Vallar et al., 1993).
23
24 This evidence suggested that CVS was only effective in restoring tactile perception because
25
26 tactile deficits were associated with visuo-spatial neglect. Hemianaesthesia was therefore a
27
28 manifestation of inattention for the neglected side of space (Vallar et al., 1990; Vallar et al.,
29
30 1993). The temporary remission caused by CVS might have been induced by the indirect
31
32 activation of an attentional orientation mechanism: CVS caused a shift of attention toward the
33
34 neglected side of the body, re-activating normal somatosensation (Vallar et al., 1990; Vallar et
35
36 al., 1993).

37
38
39 Similar remissions of somatosensory deficits have been observed by applying a
40
41 different technique, namely galvanic vestibular stimulation. Galvanic vestibular stimulation
42
43 (GVS) is a non-invasive technique which involves passing a weak direct current between
44
45 surface electrodes placed behind the ear (see Fitzpatrick and Day, 2004 for a review). It is
46
47 thought to stimulate the entire vestibular nerve, including both otolith and semicircular canal
48
49 afferents (Stephan et al., 2005). In the traditional bilateral bipolar GVS configuration, an anode
50
51 and cathode are placed on the left and right mastoid, or vice versa. Perilymphatic cathodal
52
53 currents are thought to depolarize the trigger site and lead to excitation, whereas anodal currents
54
55 hyperpolarize it resulting in inhibition (Goldberg et al., 1984). GVS evokes a sensation of head
56
57 motion in space and disturbances of equilibrium (Fitzpatrick and Day, 2004). Crucially, the
58
59
60

1
2
3
4 polarity of stimulation can be reversed as part of the experimental procedure, producing
5
6 opposite effects on firing rate in the two vestibular organs, and thus reversing of direction of the
7
8 apparent head motion. GVS has been found to influence the inability to process or attend to the
9
10 contralesional stimulus when two stimuli are simultaneously presented, in right brain-damaged
11
12 patients (i.e. tactile extinction) (Kerkhoff et al., 2011; Schmidt et al., 2013a). This remission of
13
14 tactile extinction is polarity-specific: left anodal and right cathodal GVS reduces tactile
15
16 extinction in right brain-damaged patients. Importantly, a repeated number of GVS sessions
17
18 can induce significant changes in tactile extinction that remains stable for several weeks
19
20 (Kerkhoff et al., 2011; Schmidt et al., 2013a).
21

22 Although these clinical observations favour the idea that vestibular-induced remission
23
24 of primary somatosensory deficits may be an attentional phenomenon (Utz et al., 2010; Utz et
25
26 al., 2011; Miller et al., 2008), some evidence suggests that the influence of vestibular signals on
27
28 somatosensation may be direct. First, the vestibular-induced remission of somatosensory
29
30 deficits is dissociated from visuo-spatial hemineglect (Vallar et al., 1993). Although the
31
32 majority of right brain-damaged patients with somatosensory deficits who were sensitive to
33
34 vestibular stimulation also showed visuo-spatial hemineglect, Vallar et al. (1993) additionally
35
36 described vestibular-induced remission of hemianaesthesia in two patients without visuo-spatial
37
38 hemineglect. These observations suggest that vestibular stimulation might directly affect
39
40 somatosensory processing, independently from its modulation of visuo-spatial attention.
41
42 Importantly, it is known that primary somatosensory deficits and visuo-spatial neglect are
43
44 double-dissociated disorders (Bisiach and Vallar, 1988). Second, it has been demonstrated that
45
46 contralesional hemianaesthesia was ameliorated with comparable effects in both right brain-
47
48 damaged and left brain-damaged patients who were administered with left ear CVS in order to
49
50 selectively activate the right hemisphere vestibular projections (Bottini et al., 2005; see Bottini
51
52 and Gandola 2015 for a review). Third, remission of somatosensory deficits has been described
53
54 even in a patient affected by a lesion directly involving the primary somatosensory cortex
55
56 (Bottini et al., 1995). Finally, the neural correlates of the vestibular-induced remission of left
57
58
59
60

1
2
3
4 hemianesthesia included activations in the right hemisphere involving the somatosensory
5
6 cortices, insula, right putamen, inferior frontal gyrus in the premotor cortex (Bottini et al.,
7
8 1995). Thus an undamaged subset of “sensory body representations” (cf. Bottini et al., 1995)
9
10 may be able to mediate somatosensation when an appropriate physiological manipulation is
11
12 introduced.

13
14 We recently hypothesised that vestibular signals might directly modulate the gain of the
15
16 somatosensory afferent pathway, [influencing somatosensation independently of spatial attention](#)
17
18 (Ferrè et al., 2011b; Ferrè et al., 2012; Ferrè et al., 2013b). Since this hypothesis proposes
19
20 modality-specific changes in [somatosensory](#) processing, it is possible to distinguish it from non-
21
22 specific attentional or spatial effects (Ferrè et al., 2011a; Ferrè et al., 2011b; Ferrè et al., 2013a).
23
24 To test this hypothesis, [CVS](#) was administered in healthy participants (Ferrè et al., 2011b). The
25
26 sensitivity to detect faint [tactile](#) stimuli applied on the index finger of both hands was estimated
27
28 [using signal detection](#). An enhancement of perceptual sensitivity was induced by [left ear CVS](#).
29
30 This effect was found for detection of somatosensory stimuli on both hands, i.e., ipsilateral and
31
32 contralateral to the side of [the vestibular stimulation](#). A visual detection task was administered
33
34 in the same group of participants during the same testing session to control for [arousal or](#)
35
36 [attentional effects](#): no vestibular-induced [modulation](#) of visual sensitivity was found.

37
38
39 A similar experimental design combined with [GVS](#) confirmed such vestibular-induced
40
41 enhancement. Left anodal and right cathodal [GVS](#) improved the ability to detect faint [tactile](#)
42
43 stimuli (Ferrè et al., 2013a). The vestibular-induced enhancement was found for stimuli on both
44
45 left and right hands, i.e. ipsilateral and contralateral to left anodal and right cathodal [GVS](#).
46
47 Right anodal and left cathodal [GVS](#) had no significant effects on somatosensory detection.
48
49 [Such polarity-specific influence on touch could reflect altered somatosensory processing in the](#)
50
51 [right hemisphere](#). [GVS produces polarity-dependent differences in a percept described as a](#)
52
53 [virtual rotation vector](#) (Day and Fitzpatrick, 2005) as well as sensory and cognitive functions in
54
55 both healthy volunteers and brain-damaged patients (Utz et al., 2010). [Neuroimaging studies](#)
56
57 [have revealed asymmetrical cortical vestibular projections, suggesting that the core region of the](#)
58
59
60

1
2
3
4 vestibular network is primarily located in the non-dominant right hemisphere in right-handed
5 subjects (Bense et al., 2001; Suzuki et al., 2001; Dieterich et al., 2003; Janzen et al., 2008).
6
7
8 Fink et al. (2003) used fMRI to study the effects of GVS. They found that left anodal and right
9
10 cathodal GVS caused unilateral activation of the right hemisphere vestibular projections only.
11
12 In contrast, the reversed polarity, right anodal and left cathodal stimulation, activated both left
13
14 and right hemispheres. Thus, the right hemisphere vestibular projections are activated by both
15
16 left anodal and right cathodal GVS and right anodal and left cathodal GVS (Eickhoff et al.,
17
18 2006; Lopez et al., 2012a; Zu Eulenburg et al., 2011), yet the enhancement in somatosensory
19
20 sensitivity was found only for left anodal and right cathodal GVS. Two alternative mechanisms
21
22 could explain this vestibular modulation of somatosensation. First, GVS might diffusely
23
24 activate an entire cerebral hemisphere by vestibular input, producing non-specific increases in
25
26 cortical excitability (Utz et al., 2010). If the two hemispheres further differ in their
27
28 contributions to somatosensation, then this might account for the described effects.
29
30 Alternatively, vestibular input might specifically project to cortical areas within each
31
32 hemisphere that are involved in somatosensation, such as the primary and secondary
33
34 somatosensory cortices. However, fMRI studies identify a relatively stronger activation of the
35
36 right hemisphere during left anodal and right cathodal GVS compared to the inverse polarity
37
38 (Fink et al., 2003). Thus, it is not possible to exclude the possibility that the right-hemispheric
39
40 activation during right anodal and left cathodal GVS was too weak to modulate the
41
42 somatosensory processing.
43
44

45 Electrophysiological results also support the hypothesis of vestibular contributions to
46
47 somatosensation. Somatosensory evoked potentials (SEPs) elicited by left median nerve
48
49 showed a vestibular-induced modulation in the N80 component over both ipsilateral and
50
51 contralateral somatosensory areas (Ferrè et al., 2012). The N80 component has been localised
52
53 in the parietal operculum (area OP 1; Jung et al., 2009; Eickhoff et al., 2010), which
54
55 functionally corresponds to the secondary somatosensory cortex (Eickhoff et al., 2010). The
56
57 cortical area from which N80 is assumed to arise is immediately adjacent to the
58
59
60

1
2
3
4 neuroanatomical site of vestibular-somatosensory convergence in the human homologue of the
5
6 monkey PIVC, identified as OP 2 (zu Eulenberg et al., 2012).
7

8 The multiply-connected nature of vestibular cortical anatomy suggests that vestibular
9
10 signals might also participate in a multi-way interaction with several somatic signals (Ferrè et
11
12 al., 2015a). For example, visual and vestibular signals might independently modulate
13
14 somatosensation. Alternatively, vestibular signals might interact with the effects of vision on
15
16 somatosensation, by facilitating or suppressing a visual signal that in turn influences
17
18 somatosensation. A marker of the former arrangement would be that the effects of combined
19
20 visual and vestibular stimulation should be predictable from independent effects of visual and of
21
22 vestibular stimulation on somatosensation. Conversely, a marker of the latter arrangement
23
24 would be that results of trimodal stimulation differ from simple superposition of vestibular-
25
26 somatosensory and visual-somatosensory response patterns. This hypothesis has been recently
27
28 investigated (Ferrè et al., 2015a). Participants were instructed to detect faint somatosensory
29
30 stimuli delivered to the left index finger. A visual signal occurred close to the finger in some
31
32 trials. A near infra-red caloric vestibular stimulus was used to artificially activate the vestibular
33
34 organs. Near infra-red CVS is a recently-developed technique that artificially stimulates the
35
36 vestibular system by a gradual non-contact thermal stimulation of the horizontal semicircular
37
38 canals via the external auditory canal (Ferrè et al., 2015a). Both visual and vestibular events
39
40 increased somatosensory sensitivity. Critically, no evidence for supra-additive multisensory
41
42 enhancement emerged when both visual and vestibular signals were administered together.
43
44 Visual and vestibular inputs have therefore facilitatory influences on somatosensory processing,
45
46 while vestibular inputs additionally suppress visual processing, consistently with known
47
48 neuroanatomical interactions between vestibular, visual and somatosensory systems (Ferrè et
49
50 al., 2015a).
51

52
53 Vestibular stimulation has dramatically different effects *within* the somatosensory
54
55 system. A reduction of chronic pain has been demonstrated in patients affected by right brain-
56
57 damage (McGeoch et al., 2008a; Ramachandran et al., 2007; McGeoch et al., 2008b). At least
58
59
60

1
2
3
4 two alternative mechanisms have been suggested to explain these effects (McGeoch et al.,
5
6 2008a; Ramachandran et al., 2007; McGeoch et al., 2008b). First, pain relief may be caused by
7
8 activation of the therosensory cortex in the dorsal posterior insula adjacent to PIVC stimulated
9
10 by the vestibular stimulation. Alternatively, the PIVC itself may be part of the interoceptive
11
12 system and have a direct role in pain control. In a recent study, psychophysical thresholds for
13
14 touch and pain were estimated in healthy participants immediately before and after CVS (Ferrè
15
16 et al., 2013b). Results confirmed a vestibular-induced enhancement of touch, but revealed a
17
18 reduction in levels of pain (A δ nociceptors) (Ferrè et al., 2013b). To clarify the neural
19
20 mechanisms underlying such vestibular-induced *analgesia*, CVS has been combined with
21
22 electrocortical responses elicited by nociceptive-specific laser stimulation (laser-evoked
23
24 potentials, LEPs) (Ferrè et al., 2015b). CVS resulted in a significant reduction of the amplitude
25
26 of all LEP components, including the early N1 wave which reflects the first arrival of
27
28 nociceptive input to primary somatosensory cortex. The analgesic effect induced by CVS could
29
30 be mediated by either subcortical gating of the ascending nociceptive input, or by direct
31
32 modulation of the primary somatosensory cortex. A striking feature of the vestibular
33
34 contribution to somatosensation, therefore, is the independent modulation of distinct
35
36 somatosensory submodalities. Enhancement in tactile system demonstrates an up-regulation of
37
38 tactile processing, while reduction of the level of pain demonstrates a down-regulation of
39
40 nociceptive processing. The vestibular system modulates connections with different
41
42 somatosensory submodalities, regulating the activity in multiple sensory systems independently.
43
44
45
46

47 **Vestibular contributions to somatoperception**

48
49 *Somatoperception* refers to the processes of constructing percepts and experiences of
50
51 somatic objects and events, and of one's own body (Longo et al., 2010). The body is the vehicle
52
53 for perception: percepts of stimuli applied on the skin surface must be referenced to pre-existing
54
55 representations of the body. However, the body can also be a direct object of perception. This
56
57 is seen in the process of constructing explicit models of body form (*body image*, a percept of
58
59
60

1
2
3
4 what my body is), and the configuration of body parts (*postural schema*, a percept of where my
5
6 body is). Vestibular signals contribute to cases in which the body is a vehicle for perception,
7
8 and to cases in which it is itself the object of perception.
9

10 11 12 *Localisation of touch on body surface*

13
14 The ability to localise touches on the skin surface (*topognosis*) is essential to
15
16 successfully interact with the surrounding environment. Localising the stimulus within a
17
18 somatotopic map is not by itself sufficient to identify its position on the body surface. Rather, a
19
20 two-stage process is required (Longo et al., 2010). First, the stimulus is localised within the
21
22 somatotopic map. Second, the somatotopic location must be mapped onto a corresponding
23
24 bodily location. This second step requires a body model, relating the receptor surface of the
25
26 skin to locations on the body, and thus involving an implicit somatoperception. Consistent with
27
28 this model for somatoperception, localising a tactile stimulus requires both the somatotopic
29
30 maps in the primary somatosensory cortex (Seyal et al., 1997), and also additional
31
32 somatosensory processes in parietal areas posterior to the primary somatosensory cortex (Porro
33
34 et al., 2007; Van Boven et al., 2005).
35

36
37 A recent study investigated the effect of GVS on the localisation of stimuli on the skin
38
39 of the hand (Ferrè et al., 2013c). Vestibular inputs interfered with the localisation of touches on
40
41 the dorsum of the hand. This bias was selective for the proximal-distal axis: stimuli on the hand
42
43 were shifted toward the wrist. Importantly, this shift has been showed to follow a clear spatial
44
45 gradient, with a bias in the proximal direction for touches located on the dorsum, and toward the
46
47 distal direction for touches located on the fingers. Specific polarities of GVS had differential
48
49 effects on localisation of touch. In particular, the proximal shift in tactile localisation was
50
51 stronger during right anodal and left cathodal GVS than left anodal and right cathodal GVS.
52
53 Such hemisphere-specific effects might be due to the potential hemispheric lateralisation of the
54
55 ability to localise touch. Indeed, the observed vestibular-induced effects on tactile localisation
56
57 are reminiscent of those described in patients following left brain-damage. Clinical
58
59
60

1
2
3
4 observations described patients affected by left parietal lesions showing errors of tactile
5
6 localisation in the contralesional hand (Rapp et al., 2002). These localisation errors preserved
7
8 the somatotopic arrangement of the hand, but systematically misplaced each point toward the
9
10 proximal direction (Rapp et al., 2002). Rapp et al. (2002) reported two patients showing a clear
11
12 dissociation between an intact ability to detect faint tactile stimuli compared to an impaired
13
14 localisation of touches on the affected hand. In particular, the authors commented “the post-
15
16 lesion [tactile] mislocalisations may preserve the relative locations of the pre-lesion topography,
17
18 resulting in systematically shifted and distorted somatosensory experiences”.

19
20 Neuroimaging studies reveal activation of the right temporo-parietal junction during
21
22 tactile localisation (Van Boven et al., 2005). This area is also a core region of the human
23
24 vestibular cortical network (Lobel et al., 1999; Fasold et al., 2002). The anatomical overlap
25
26 between vestibular cortical areas and areas involved in tactile localisation suggests that
27
28 vestibular inputs might influence the process of registering somatotopic skin location with
29
30 specific locations on the body, although this should be tested in further studies, possibly
31
32 investigating tactile localisations in patients with peripheral vestibular disease.

33 34 35 36 37 *Proprioceptive localisation of body parts in external space*

38
39 Proprioceptive information from mechanoreceptors in joints, muscles, tendons, and
40
41 stretch-sensitive receptors in the skin describe the configuration of the body in terms of the
42
43 degree of flexion or extension of each joint (Burgess et al., 1982). The localisation of body
44
45 parts in space requires a combination of afferent information and stored representations of the
46
47 body. Head and Holmes (1911) introduced the notion of the *postural schema* to refer to the
48
49 continuously updated representation of the posture of the body.

50
51 Artificial vestibular stimulation has been showed to ameliorate proprioceptive deficits
52
53 in right brain-damaged patients. Specifically, left anodal and right cathodal GVS improved the
54
55 accuracy in detecting the position of the left arm in seven visuo-spatial hemineglect patients:
56
57 these effects were observed during online stimulation and also after 20 minutes from the
58
59
60

1
2
3
4 stimulation (Schmidt et al., 2013b). Vestibular signals might have provided additional sensory
5
6 input to help in localising the correct position of the arm. It has been suggested that such
7
8 facilitatory vestibular effects may rely on the activation of intact structures that are involved in
9
10 the recalibration of body and limb position in space. Importantly, neither right anodal and left
11
12 cathodal GVS nor sham stimulation influenced the proprioceptive localisation of the arm, ruling
13
14 out attentional or unspecific effects. The differential effects of left anodal and right cathodal
15
16 GVS compared with right anodal and left cathodal GVS might be, once again, related with the
17
18 asymmetry of the cortical vestibular network (Dieterich et al., 2003).
19

20
21 Vestibular effects on the localisation of body parts in space have been observed in
22
23 healthy participants (Schmidt et al., 2013c). Left anodal and right cathodal GVS caused a
24
25 transient disruption of left arm localisation in right-handed participants. These results have
26
27 been interpreted as a vestibular-induced interference with arm positioning which is particular
28
29 salient in right-handers due to the anatomical asymmetry of the vestibular system (Schmidt et
30
31 al., 2013c; Dieterich et al., 2003). Similarly, it has been reported that GVS disrupted reaching
32
33 movements towards a visual target by interfering with the spatial estimate of where the hands
34
35 are in the surrounding space (Bresciani et al., 2002).
36
37
38

39 **Vestibular contributions to somatopresentation**

40
41 *Somatopresentation* involves the on-line construction of higher-level percepts related
42
43 to the body. This representation is not merely an abstract conceptualisation, but actually
44
45 underlies all bodily experience, for the reasons that we describe below.
46
47
48

49 *Model of body shape and size*

50
51 No peripheral receptors are directly informative about shape and size of body parts, and
52
53 therefore the knowledge about these features are linked to an internal stored model of the body's
54
55 metric properties (Longo and Haggard, 2010). It has been demonstrated that this *body model* is
56
57 strongly distorted (Longo and Haggard, 2010). At least for the hand, these distortions included
58
59
60

1
2
3
4 a radial–ulnar gradient of magnification of the digits and shrinkage toward the proximo–distal
5 axis. These internal models appear to be independent of any specific somatosensory input for
6 two reasons. First, people can judge the configuration and location of the body, including its
7 metric properties, in the absence of specific somatosensory stimulation (Longo and Haggard,
8 2010). Second, these metric properties are accessible even for congenital phantoms (Longo et
9 al., 2012). For this reason, the internal body model involves a level of *somatorepresentation*,
10 independent of any particular somatosensory event or stimulus.
11
12
13
14
15
16
17

18 A contribution of vestibular signals to the metric body model is evident from clinical
19 observations in patients affected by peripheral vestibular disorders. Bonnier (1905) described
20 distortions of the representation of body shape and size, such as “[the patient] felt his head
21 become enormous, immense, losing itself in the air” and “her vertigo gave her the sensation that
22 she no longer exists as a body” (Bonnier, 1905).
23
24
25
26
27

28 The relation between vestibular signals and the metric body model has been
29 investigated using CVS in psychophysical paradigm in healthy participants. Lopez et al.
30 (2012b) asked participants to point to different anatomical landmarks on the left hand (knuckles
31 and tip of the fingers) without seeing them, immediately before and after CVS. The perceived
32 length and width of the hand were estimated based on these judgments. CVS significantly
33 increased both the perceived length and width of the hand. These data are comparable with
34 clinical observations of vestibular peripheral patients (“extremities had become larger”,
35 Schilder, 1935; see Lopez 2015 for a review). However, no such effects were found in a similar
36 task combined with low-intensity GVS (Ferrè et al., 2013c). This may be due to differences in
37 the vestibular stimulation technique used. For instance, GVS effects are relatively weak
38 compared to the CVS. Further studies need to clarify those differences.
39
40
41
42
43
44
45
46
47
48
49
50

51 52 53 *Ownership of body parts*

54 The sense of body ownership is often described as a feeling of *mineness*, that one's body
55 belongs to oneself (Metzinger, 2003). The most convincing evidence for such a concept comes
56
57
58
59
60

1
2
3
4 from disordered body representation in neuropsychological patients. For example, right brain-
5
6 damaged patients with *somatoparaphrenia* lack the normal sense of the body as a coherent
7
8 entity, as to the feeling that one's body belongs to oneself (Gallagher, 2000). For example, they
9
10 may have delusional beliefs that their own limbs belong to someone else (Vallar and Ronchi,
11
12 2009). Traditionally, a strong emphasis has been put on the role of visual and somatosensory
13
14 signals in body ownership. This is because ownership of body parts strongly relies on the visual
15
16 feedback from one's own actions and the somatosensory feedback from this action (Botvinick
17
18 and Cohen, 1998; Tsakiris and Haggard, 2005). However, other sensory modalities potentially
19
20 contribute to body ownership.
21

22
23 Studies of individual cases described temporary remissions of somatoparaphrenia
24
25 following artificial vestibular stimulation (Bisiach et al., 1991; Rode et al., 1992; Schiff and
26
27 Pulver, 1999). As described in Bisiach et al. (1991), "A.R. would unhesitatingly attribute her
28
29 left upper limb to her mother, [but] under the effect of vestibular stimulation she unhesitatingly
30
31 recognizes that same arm as her own". Recently, CVS was reported to reduce
32
33 somatoparaphrenia for the right hand also in a left brain-damaged patient (Ronchi et al., 2013).
34
35 Injection of cold water in the right auditory canal transitory ameliorated the delusional belief
36
37 that the patient's right hand did not belong to her.
38

39
40 The hypothesis that vestibular inputs contribute to the sense of ownership has been
41
42 explored in healthy participants using the Rubber Hand Illusion paradigm (RHI, Botvinick and
43
44 Cohen, 1998). A rubber hand viewed in peripersonal space is experienced as part of one's body
45
46 if it is touched in synchrony with the participant's own unseen hand. RHI involves a
47
48 transposition of the sense of body from one object to another, rather than addition of a new body
49
50 part (Longo et al., 2008; Moseley et al., 2008; Barnsley et al., 2011). That is, *dis-ownership* of
51
52 one's own hand accompanies the increased sense of ownership over the rubber hand.
53
54 Consistent with this account, participants then mislocate their own hand as closer to the fake
55
56 hand than it actually is. This *proprioceptive drift* increases with the increasing strength of the
57
58 illusion (Botvinick and Cohen, 1998), and provides a quantitative proxy for assessing bodily
59
60

1
2
3
4 awareness (Longo et al., 2008). Combining RHI with GVS revealed once again a polarity
5
6 dependent vestibular modulation of the strength of the RHI illusion (Ferrè et al., 2015c). Left
7
8 anodal and right cathodal GVS, which predominantly activates the vestibular projections in the
9
10 right hemisphere, produced a smaller proprioceptive shift toward the rubber hand compared
11
12 with the opposite polarity (Ferrè et al., 2015c). The right hemisphere vestibular network
13
14 therefore increases the salience of intrinsic somatosensory and proprioceptive signals about
15
16 hand position, and decreases the salience of visual information responsible for visual capture
17
18 during the RHI. However, Lopez et al. (2010) found a vestibular induced enhancement of the
19
20 RHI as measured by questionnaires, but no reliable effects on proprioceptive drift. While it is
21
22 difficult to reconcile results from these studies, it is notable that proprioceptive drift (Ferrè et
23
24 al., 2015) and questionnaires (Lopez et al., 2010) are two different and independent aspects of
25
26 the RHI (Rohde et al., 2011).
27

28
29 Vestibular contributions to other disorders involving abnormal forms of body
30
31 ownership have been proposed. Ramachandran and McGeoch (2007) hypothesised that
32
33 vestibular stimulation could be used to treat patients with a desire for amputation of healthy
34
35 body parts, a disorder known as Body Identity Integrity Disorder (BIID). On one hypothesis,
36
37 BIID reflects a mismatch between a neural representation of the actual anatomical body, and a
38
39 second (unidentified) representation of the desired body form (Ramachandran and McGeoch,
40
41 2007). Blanke et al. (2009) suggested that BIID can be considered a chronic form of
42
43 asomatognosia, essentially a negative phantom limb. A recent study tested Ramachandran's
44
45 hypothesis in a group of 13 BIID patients (Lenggenhager et al., 2014). Neither left nor right ear
46
47 CVS reduced their desire for amputation. The underlying mechanisms of BIID may be just
48
49 different compared somatoparaphrenia: while BIID is a long-term condition that has often been
50
51 consolidated since childhood (Brugger et al., 2013), somatoparaphrenia can be observed during
52
53 the acute phase after brain injury and may be more prone to modulations by artificial vestibular
54
55 stimulation.
56
57
58
59
60

First-person perspective

An important aspect of the representation of our body is the consistent first-person perspective on the external world. Indeed, almost all our bodily experiences are tagged by a first-person perspective. Vestibular damage can often produce psychiatric symptoms of depersonalisation, defined as the subjective experience of unreality and detachment from the self, and derealisation, the experience of the external world appearing strange or unreal (Sang et al., 2006; Jauregui-Renaud et al., 2008; see Jauregui-Renaud, 2015 for a review). Similar feelings of unreality have also been found in healthy participants undergoing artificial vestibular stimulation (Sang et al., 2006).

Out-of-body experiences, in which neurological patients localise the self outside their own body and experience seeing their body from this disembodied location, represents a failure in the first person perspective and were attributed to conflicting visual, somatosensory and vestibular information (Brugger et al., 1997). Interestingly, out-of-body experiences are frequently associated with vestibular sensations, such as the feeling of floating in the room, levitation and lightness (Blanke and Mohr, 2005; Blanke et al., 2004; Heydrich and Blanke, 2011). The vestibular contribution to these phenomena is further supported by the association between out-of-body experience and temporo-parietal junction and insula (Blanke et al. 2004), which are considered the two main regions receiving vestibular projections (Heydrich and Blanke, 2013; Ionta et al., 2011, Mazzola et al., 2014; Pfeiffer et al., 2014).

One can therefore hypothesize that a fundamental function of the vestibular input might be to provide the normal first-person perspective on the world. Ferrè et al. (2014) investigated whether vestibular signals influence the perspective people take (first-person perspective vs third-person perspective) in interpreting ambiguous stimuli applied on the skin surface (e.g. graphesthesia task, Natsoulas and Dubanoski, 1964). For example, when an experimenter draws the letter *b* on the forehead of a blindfolded participant, the participant may perceive the stimulus either from an external third-person perspective (reading it as “b”), or from an internal first-person perspective (reading it as “d”). Prevalence of one judgement over the other

1
2
3
4 provides an implicit measure of whether the perceiver's perspective on the stimulus originates
5 from within his body, or from outside it. Low intensity GVS inputs promote first-person
6 perspective judgements, augmenting a natural vestibular contribution to embodiment.
7
8 Disembodied experiences, as out-of-body phenomena, may require different stimulation
9 patterns, stronger intensities, direct stimulation of vestibular cortical areas rather than the
10 vestibular periphery, or combined with additional visual and somatosensory mismatches
11
12 (Blanke et al., 2002; Blanke et al., 2012).
13
14
15
16
17

18 Critically, disruptive vestibular signals have been showed to interfere with the so-called
19 *self-advantage* effect: when people are asked to detect asynchrony between self-generated
20 movement and the visual feedback concerning that movement, they are more sensitive when the
21 feedback is presented from a first-person perspective (Hoover and Harris, 2012; Hoover and
22 Harris, 2015). Hoover and Harris (2015) recently investigated whether artificially disruptive
23 vestibular stimulation affects the self-advantage effect. Participants performed movements
24 while viewing their hand in a first-person or third-person perspective. They judged which of
25 two experiences involved a delayed visual feedback, while disruptive GVS was applied.
26 Disruptive GVS abolished the self-advantage, so there was no longer any difference in
27 performance between first-person and third-person perspectives. These data further support the
28 idea that functional vestibular information is essential in distinguishing between self and other,
29 and in maintaining a coherent sense of ownership over body parts.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 Outlook

46 Textbooks still consider the vestibular system as an organ for balance and orienting eye
47 movements. However, signals from the vestibular system make a crucial contribution to most
48 everyday adaptive behaviours. Accordingly, vestibular information interacts widely with
49 sensory and cognitive processes (Balaban et al., 2011).
50
51
52
53
54

55 Clinical observations and experimental evidence reviewed here suggests a key
56 contribution of vestibular signals to bodily representations. The present review article aimed at
57
58
59
60

1
2
3
4 organising the existing knowledge to identify *whether* and *how* vestibular signals influence
5 body representations. As summarised above, the vestibular system strongly contributes to the
6 somatosensational, somatoperceptual and somatorepresentational components of bodily
7 awareness (Figure 1). Specifically, vestibular signals *independently* contribute to low-level
8 signal processing and also to higher-order representations of the body. Vestibular information
9 directly modulates the primary processing of somatic sensations, but also the on-line percept of
10 the state of the body. There is evidence that it influences processing within other individual
11 sensory channels, rather than simply adding an extra ingredient to the multisensory mix. Our
12 review also highlights an independent vestibular modulation in two classes of higher-order
13 representations about the body. On the one hand, vestibular information influences
14 somatoperception (the *perceptual* process of constructing perceptual representations of the
15 body). On the other, vestibular information contributes to somatorepresentation (the *cognitive*
16 process of constructing abstract knowledge about the body) (Longo et al., 2010). Importantly,
17 several lines of evidence rule out explanations based on changes in arousal, and spatial
18 attention. Rather, each level of the somatic hierarchy can be influenced by vestibular signals.
19 However, more work is needed to improve our understanding of the neurophysiological
20 mechanisms supporting the described vestibular effects.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

38 Balaban and colleagues (2011) have recently proposed a model in which vestibular
39 signals are tightly interwoven with three different biochemical pathways to serve sensorimotor
40 processing, cognition and interoception. Critically, this arrangement implies that alterations
41 within one pathway may elicit functional changes in the others (cf. the well-known comorbidity
42 of balance, anxiety and migraine; Yardley, 2000). Vestibular-induced changes in body
43 representations, be these through experimental or clinical phenomena, may therefore modulate
44 processing within allied cognitive and interoceptive domains. **Some evidence already supports**
45 **this neural influence, for example deficits in somatosensation are often present in association**
46 **with cognitive, attentional, impairments.** Here, we have proposed a multicomponent account
47 for bodily awareness and identified multiple independent vestibular contributions to it. We
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 hope that this account will open new avenues of research in the field and propose that future
5
6 neuroscientific investigations should focus on how vestibular-induced changes in body
7
8 representations may interact with other neural systems.
9

10 Under normal circumstances, every movement of the head implies a new relation
11
12 between the body-self and the external world. We suggest that vestibular contributions to body
13
14 representations may arise from a neural circuit that optimizes the balance between gathering
15
16 information from personal/peripersonal space immediately surrounding the body, and farther
17
18 environmental space. Evidence showed strong vestibular modulation of somatic sensation,
19
20 perception and representation. Interestingly, these effects could be interpreted as enhancement
21
22 of the bodily self: vestibular signals rebalance sensory events away from the world and towards
23
24 the self.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- Anastasopoulos, D., Bronstein, A., Haslwanter, T., Fetter, M., & Dichgans, J. (1999). The role of somatosensory input for the perception of verticality. *Annals of the New York Academy of Sciences*, 871(1), 379-383.
- André, J. M., Martinet, N., Paysant, J., Beis, J. M., & Le Chapelain, L. (2001). Temporary phantom limbs evoked by vestibular caloric stimulation in amputees. *Cognitive and Behavioral Neurology*, 14(3), 190-196.
- Balaban, C. D., Jacob, R. G., & Furman, J. M. (2011). Neurologic bases for comorbidity of balance disorders, anxiety disorders and migraine: neurotherapeutic implications. *Expert review of neurotherapeutics*, 11(3), 379-394.
- Balaban, C. D., & Yates, B. J. (2004). Vestibuloautonomic interactions: A teleologic perspective. In *The Vestibular System* (pp. 286-342). Springer New York.
- Barnett-Cowan, M., Snow, J. C., & Culham, J. C. (2015). Contribution of Bodily and Gravitational Orientation Cues to Face and Letter Recognition. *Multisensory research*, 28(5-6), 427-442.
- Barnsley, N., McAuley, J. H., Mohan, R., Dey, A., Thomas, P., & Moseley, G. L. (2011). The rubber hand illusion increases histamine reactivity in the real arm. *Current Biology*, 21(23), R945-R946.
- Bense, S., Stephan, T., Yousry, T. A., Brandt, T., & Dieterich, M. (2001). Multisensory cortical signal increases and decreases during vestibular galvanic stimulation (fMRI). *Journal of Neurophysiology*, 85(2), 886-899.
- Berthoz, A. (1996). How does the cerebral cortex process and utilize vestibular signals. *Disorders of the vestibular system*. Oxford University Press, Oxford, 113-125.
- Bisiach E., Rusconi M. L., & Vallar G. (1991). Remission of somatoparaphrenic delusion through vestibular stimulation. *Neuropsychologia* 29, 1029–1031.
- Bisiach, E., & Vallar, G. (1988). Hemineglect in humans. In F. Boller & J. Grafman (Eds.), *Handbook of neuropsychology*. Amsterdam: Elsevier Science.

- 1
2
3
4 Blanke, O. (2012). Multisensory brain mechanisms of bodily self-consciousness. *Nature*
5
6 *Reviews Neuroscience*, 13(8), 556-571.
7
8 Blanke, O., & Arzy, S. (2005) The out-of-body experience: disturbed self-processing at the
9
10 temporo-parietal junction. *Neuroscientist*. 11(1):16-24.
11
12 Blanke, O., & Metzinger, T. (2009). Full-body illusions and minimal phenomenal selfhood.
13
14 *Trends in cognitive sciences*, 13(1), 7-13.
15
16 Blanke, O., Landis, T., Spinelli, L., & Seeck, M. (2004). Out - of - body experience and
17
18 autoscapy of neurological origin. *Brain*, 127(2), 243-258.
19
20
21 Blanke, O., Mohr, C., Michel, C. M., Pascual-Leone, A., Brugger, P., Seeck, M., ... & Thut, G.
22
23 (2005). Linking out-of-body experience and self processing to mental own-body
24
25 imagery at the temporoparietal junction. *Journal of Neuroscience*. 25(3), 550-557.
26
27 Blanke, O., Morgenthaler, F. D., Brugger, P., & Overney, L. S. (2009). Preliminary evidence for
28
29 a fronto - parietal dysfunction in able - bodied participants with a desire for limb
30
31 amputation. *Journal of Neuropsychology*, 3(2), 181-200.
32
33 Böhmer, A., & Mast, F. (1999). Assessing otolith function by the subjective visual vertical.
34
35 *Annals of the New York Academy of Sciences*, 871, 221-231.
36
37 Bonnier, P., 1905. L'Aschématie. *Rev Neurol Paris* 12, 605-609.
38
39 Bottini, G., & Gandola, M. (2015). Beyond the Non-Specific Attentional Effect of Caloric
40
41 Vestibular Stimulation: Evidence from Healthy Subjects and Patients. *Multisensory*
42
43 *research*, 28(5-6), 591-612.
44
45 Bottini, G., Paulesu, E., Gandola, M., Loffredo, S., Scarpa, P., Sterzi, R., ... & Vallar, G. (2005).
46
47 Left caloric vestibular stimulation ameliorates right hemianesthesia. *Neurology* 65(8),
48
49 1278-1283.
50
51 Bottini, G., Paulesu, E., Sterzi, R., Warburton, E., Wise, R. J. S., Vallar, G., ... & Frith, C. D.
52
53 (1995). Modulation of conscious experience by peripheral sensory stimuli. *Nature*
54
55 376(6543), 778-781.
56
57
58
59
60

- 1
2
3
4 Botvinick, M., & Cohen, J. (1998). Rubber hands 'feel' touch that eyes see. *Nature*, 391(6669),
5
6 756-756.
7
- 8 Bresciani, J. P., Blouin, J., Popov, K., Bourdin, C., Sarlegna, F., Vercher, J. L., & Gauthier, G.
9
10 M. (2002). Galvanic vestibular stimulation in humans produces online arm movement
11
12 deviations when reaching towards memorized visual targets. *Neuroscience letters*,
13
14 318(1), 34-38.
15
- 16 Bronstein, A. M. (1999). The interaction of otolith and proprioceptive information in the
17
18 perception of verticality The effects of labyrinthine and CNS disease. *Annals of the*
19
20 New York Academy of Sciences, 871, 324–333.
21
- 22 Brugger, P., Lenggenhager, B., & Giummarra, M. J. (2013). Xenomelia: a social neuroscience
23
24 view of altered bodily self-consciousness. *Frontiers in Psychology*, 48, 23-77.
25
- 26 Brugger, P., Regard, M., & Landis, T. (1997). Illusory reduplication of one's own body:
27
28 phenomenology and classification of autoscopic phenomena. *Cognitive*
29
30 *Neuropsychiatry*, 2(1), 19-38.
31
- 32 Burgess, P. R., Wei, J. Y., Clark, F. J., & Simon, J. (1982) Signaling of kinesthetic information
33
34 by peripheral sensory receptors. *Annual review of neuroscience*, 5(1), 171-188.
35
- 36 Cappa S., Sterzi R., Vallar G., & Bisiach E. (1987). Remission of hemineglect and anosognosia
37
38 during vestibular stimulation. *Neuropsychologia* 25, 775–782.
39
- 40 Day, B. L., & Fitzpatrick, R. C. (2005). The vestibular system. *Current biology*, 15(15), R583-
41
42 R586.
43
- 44 Dieterich, M., Bense, S., Lutz, S., Drzezga, A., Stephan, T., Bartenstein, P., & Brandt, T.
45
46 (2003). Dominance for vestibular cortical function in the non-dominant hemisphere.
47
48 *Cerebral Cortex*, 13(9), 994-1007.
49
- 50 Eickhoff, S. B., Jbabdi, S., Caspers, S., Laird, A. R., Fox, P. T., Zilles, K., & Behrens, T. E.
51
52 (2010). Anatomical and functional connectivity of cytoarchitectonic areas within the
53
54 human parietal operculum. *Journal of Neuroscience*. 30(18), 6409-6421.
55
56
57
58
59
60

- 1
2
3
4 Eickhoff, S. B., Weiss, P. H., Amunts, K., Fink, G. R., & Zilles, K. (2006). Identifying human
5
6 parieto - insular vestibular cortex using fMRI and cytoarchitectonic mapping. *Human*
7
8 *brain mapping*, 27(7), 611-621.
- 9
10 Eilan, N., Bermúdez, J. L., & Marcel, A. (Eds.). (1998). *The body and the self*. Massachusetts:
11
12 MIT Press.
- 13
14 Fasold, O., von Brevern, M., Kuhberg, M., Ploner, C.J., Villringer, A., Lempert, T., & Wenzel,
15
16 R. (2000) Human vestibular cortex as identified with caloric stimulation in functional
17
18 magnetic resonance imaging. *Neuroimage*. 17(3): 1384-93.
- 19
20 Ferrè, E. R., Berlot, E., & Haggard, P. (2015c). Vestibular contributions to a right-hemisphere
21
22 network for bodily awareness: Combining galvanic vestibular stimulation and the
23
24 “Rubber Hand Illusion”. *Neuropsychologia*, 69, 140-147.
- 25
26 Ferrè, E. R., Bottini, G., & Haggard, P. (2011b). Vestibular modulation of somatosensory
27
28 perception. *European Journal of Neuroscience*. 34(8), 1337-1344.
- 29
30 Ferrè, E. R., Bottini, G., & Haggard, P. (2012). Vestibular inputs modulate somatosensory
31
32 cortical processing. *Brain Structure and Function*. 217(4), 859-864.
- 33
34 Ferrè, E. R., Bottini, G., Iannetti, G. D., & Haggard, P. (2013b). The balance of feelings:
35
36 vestibular modulation of bodily sensations. *Cortex* 49(3), 748-758.
- 37
38 Ferrè, E. R., Day, B. L., Bottini, G., & Haggard, P. (2013a). How the vestibular system interacts
39
40 with somatosensory perception: A sham-controlled study with galvanic vestibular
41
42 stimulation. *Neuroscience Letters*. 550, 35-40.
- 43
44 Ferrè, E. R., Haggard, P., Bottini, G., & Iannetti, G. D. (2015b). Caloric vestibular stimulation
45
46 modulates nociceptive evoked potentials. *Experimental brain research*, 233(12), 3393-
47
48 3401.
- 49
50 Ferrè, E. R., Lopez, C., & Haggard, P. (2014). Anchoring the Self to the Body Vestibular
51
52 Contribution to the Sense of Self. *Psychological science*, 25(11), 2106-2108.
- 53
54
55
56
57
58
59
60

- 1
2
3
4 Ferrè, E. R., Sedda, A., Gandola, M., & Bottini, G. (2011a). How the vestibular system
5 modulates tactile perception in normal subjects: a behavioural and physiological study.
6 *Experimental Brain Research*, 208(1), 29-38.
7
8
9
10 Ferrè, E. R., Vagnoni, E., & Haggard, P. (2013c). Vestibular contributions to bodily awareness.
11 *Neuropsychologia*, 51(8), 1445-1452.
12
13
14 Ferrè, E. R., Walther, L. E., & Haggard, P. (2015a). Multisensory Interactions between
15 *Vestibular, Visual and Somatosensory Signals*. *PloS one*, 10(4), e0124573.
16
17
18 Fink, G. R., Marshall, J. C., Weiss, P. H., Stephan, T., Grefkes, C., Shah, N. J., ... & Dieterich,
19 M. (2003). Performing allocentric visuospatial judgments with induced distortion of the
20 egocentric reference frame: an fMRI study with clinical implications. *Neuroimage*,
21 20(3), 1505-1517.
22
23
24
25
26 Fitzpatrick, R. C., & Day, B. L. (2004). Probing the human vestibular system with galvanic
27 stimulation. *Journal of Applied Physiology*. 96(6), 2301-2316.
28
29
30
31 Gallagher, S. (2000). Philosophical concepts of the self: Implications for cognitive sciences.
32 *Trends in Cognitive Sciences*, 4, 14–21.
33
34
35 Goldberg J. M., Smith C. E., & Fernandez C. (1984). Relation between discharge regularity and
36 responses to externally applied galvanic currents in vestibular nerve afferents of the
37 squirrel monkey. *Journal of Neurophysiology*. 51, 1236–1256.
38
39
40
41 Grabherr, L., Macaudo, G., & Lenggenhager, B. (2015). The Moving History of Vestibular
42 *Stimulation as a Therapeutic Intervention*. *Multisensory research*, 28(5-6), 653-687.
43
44
45
46
47 Guldin, W. O., & Grüsser, O. J. (1998). Is there a vestibular cortex?. *Trends Neurosci*. 21(6),
48 254-259.
49
50
51
52
53
54
55
56
57
58
59
60
- Head, H., & Holmes, G. (1911) Sensory disturbances from cerebral lesions. *Brain*, 34, 102–254.
- Heydrich, L., & Blanke, O. (2013). Distinct illusory own-body perceptions caused by damage to
posterior insula and extrastriate cortex. *Brain*, 364.
- Ho, C., & Spence, C. (2007). Head orientation biases tactile localization. *Brain Research*, 1144,
136-141.

- 1
2
3
4 Hoover, A. E., & Harris, L. R. (2012). Detecting delay in visual feedback of an action as a
5
6 monitor of self recognition. *Experimental brain research*, 222(4), 389-397.
7
8 Hoover, A. E., & Harris, L. R. (2015). Disrupting Vestibular Activity Disrupts Body
9
10 Ownership. *Multisensory research*, 28(5-6), 581-590.
11
12 Iwamura, Y. (1998). Hierarchical somatosensory processing. *Current opinion in neurobiology*,
13
14 8(4), 522-528.
15
16 Ionta, S., Heydrich, L., Lenggenhager, B., Mouthon, M., Fornari, E., Chapuis, D., ... & Blanke,
17
18 O. (2011). Multisensory mechanisms in temporo-parietal cortex support self-location
19
20 and first-person perspective. *Neuron*, 70(2), 363-374.
21
22 Janzen, J., Schlindwein, P., Bense, S., Bauermann, T., Vucurevic, G., Stoeter, P., & Dieterich,
23
24 M. (2008). Neural correlates of hemispheric dominance and ipsilaterality within the
25
26 vestibular system. *Neuroimage*, 42(4), 1508-1518.
27
28 Jáuregui-Renaud, K. (2015). Vestibular Function and Depersonalization/Derealization
29
30 Symptoms. *Multisensory research*, 28(5-6), 637-651.
31
32 Jáuregui-Renaud, K., Sang, F. Y. P., Gresty, M. A., Green, D. A., & Bronstein, A. M. (2008).
33
34 Depersonalisation/derealisation symptoms and updating orientation in patients with
35
36 vestibular disease. *Journal of Neurology, Neurosurgery & Psychiatry*, 79(3), 276-283.
37
38 Johnson, K. O., & Hsiao, S. S. (1992). Neural mechanisms of tactual form and texture
39
40 perception. *Annual review of neuroscience*, 15(1), 227-250.
41
42 Jung, P., Baumgärtner, U., Stoeter, P., & Treede, R. D. (2009). Structural and functional
43
44 asymmetry in the human parietal opercular cortex. *Journal Neurophysiology*. 101(6),
45
46 3246-3257.
47
48 Kaas, J. H. (1983). What, if anything, is SI? Organization of first somatosensory area of cortex.
49
50 *Physiological Review*, 63(1), 206-231.
51
52 Kerkhoff, G., Hildebrandt, H., Reinhart, S., Kardinal, M., Dimova, V., & Utz, K. S. (2011). A
53
54 long-lasting improvement of tactile extinction after galvanic vestibular stimulation: two
55
56 Sham-stimulation controlled case studies. *Neuropsychologia* 49(2), 186-195.
57
58
59
60

- 1
2
3
4 Lackner, J. R. (1988). Some proprioceptive influences on the perceptual representation of body
5 shape and orientation. *Brain*, 111, 281–297.
6
7
8 Lackner, J. R. DiZio, P. (2005). Vestibular, proprioceptive, and haptic contributions to spatial
9 orientation. *Annual Review of Psychology*, 56, 115–147.
10
11
12 **Lacquaniti, F., Bosco, G., Gravano, S., Indovina, I., La Scaleia, B., Maffei, V., & Zago, M.**
13 **(2015). Gravity in the brain as a reference for space and time perception. *Multisensory***
14 ***research*, 28(5-6), 397-426.**
15
16
17
18 Le Chapelain, L., Beis, J. M., Paysant, J., & André, J. M. (2001). Vestibular caloric stimulation
19 evokes phantom limb illusions in patients with paraplegia. *Spinal Cord*, 39(2), 85-87.
20
21
22 Lenggenhager, B., Hilti, L., Palla, A., Macaudo, G., & Brugger, P. (2014). Vestibular
23 stimulation does not diminish the desire for amputation. *Cortex*, 54, 210-212.
24
25
26 Lobel, E., Kleine, J.F., Leroy-Willig, A., Van de Moortele, P.F., Le Bihan, D., Grüsser, O.J., &
27 Berthoz, A. (1999) Cortical areas activated by bilateral galvanic vestibular stimulation.
28 *Annals of the New York Academy of Sciences* 871:313-23.
29
30
31
32 Longo, M. R., & Haggard, P. (2010). An implicit body representation underlying human
33 position sense. *Proceedings of the National Academy of Sciences*, 107(26), 11727-
34 11732.
35
36
37
38 Longo, M. R., Azañón, E., & Haggard, P. (2010). More than skin deep: body representation
39 beyond primary somatosensory cortex. *Neuropsychologia*, 48(3), 655-668.
40
41
42 Longo, M. R., Schüür, F., Kammers, M. P., Tsakiris, M., & Haggard, P. (2008). What is
43 embodiment? A psychometric approach. *Cognition*, 107(3), 978-998.
44
45
46 Longo, M.R., Long, C., & Haggard, P. (2012) Mapping the invisible hand: a body model of a
47 phantom limb. *Psychological Science*. 23(7):740-2.
48
49
50
51 **Lopez, C. (2015). Making sense of the body: The role of vestibular signals. *Multisensory***
52 ***research*, 28(5-6), 525-557.**
53
54
55
56
57
58
59
60

- 1
2
3
4 Lopez, C., Blanke, O., & Mast, F. W. (2012a). The human vestibular cortex revealed by
5 coordinate-based activation likelihood estimation meta-analysis. *Neuroscience*. 212,
6 159-179.
7
8
9
10 Lopez, C., Halje, P., & Blanke, O. (2008). Body ownership and embodiment: vestibular and
11 multisensory mechanisms. *Neurophysiologie Clinique/Clinical Neurophysiology*, 38(3),
12 149-161.
13
14
15
16 Lopez, C., Lenggenhager, B., & Blanke, O. (2010). How vestibular stimulation interacts with
17 illusory hand ownership. *Consciousness and cognition*, 19(1), 33-47.
18
19
20 Lopez, C., Schreyer, H.M., Preuss, N., & Mast, F.W. (2012b) Vestibular stimulation modifies
21 the body schema. *Neuropsychologia*. 50(8):1830-7.
22
23
24 Mazzola, L., Lopez, C., Faillenot, I., Chouchou, F., Mauguière, F., & Isnard, J. (2014).
25 Vestibular responses to direct stimulation of the human insular cortex. *Ann. Neurol.* 76,
26 609–619.
27
28
29
30 Merleau-Ponty. (1962). *The phenomenology of perception* (C. Smith, Trans.). New York: The
31 Humanities Press.
32
33
34 McGeoch, P. D., & Ramachandran, V. S. (2008a). Vestibular stimulation can relieve central
35 pain of spinal origin. *Spinal cord*. 46(11), 756-757.
36
37
38 McGeoch, P. D., Williams, L. E., Lee, R. R., & Ramachandran, V. S. (2008b). Behavioural
39 evidence for vestibular stimulation as a treatment for central post-stroke pain. *J Neurol*
40 *Neurosurg Psychiatry*. 79;1298-1301.
41
42
43
44 Medina, J., & Rapp, B. (2008). Phantom tactile sensations modulated by body position. *Current*
45 *Biology*, 18(24), 1937-1942.
46
47
48 Metzinger, T. (2003). Phenomenal transparency and cognitive self-reference. *Phenomenology*
49 *and the Cognitive Sciences*, 2(4), 353-393.
50
51
52 Miller, S. M., & Ngo, T. T. (2007). Studies of caloric vestibular stimulation: implications for
53 the cognitive neurosciences, the clinical neurosciences and neurophilosophy. *Acta*
54 *Neuropsychiatr.* 19(3), 183-203.
55
56
57
58
59
60

- 1
2
3
4 Mittelstaedt, H. (1991). The role of the otoliths in the perception of the orientation of self and
5 world to the vertical. *Zoologische Jahrbücher Abteilung Physiologie*, 95, 418–425.
6
7
8 Mittelstaedt, H. (1992). Somatic versus vestibular gravity reception in man. *Annals of the New*
9 *York Academy of Sciences*, 656, 124–139.
10
11
12 Mittelstaedt, H. (1999). The role of the otoliths in perception of the vertical and in path
13 integration. *Annals of the New York Academy of Sciences*, 871, 334–344.
14
15
16 Moseley, G. L., Olthof, N., Venema, A., Don, S., Wijers, M., Gallace, A., & Spence, C. (2008).
17 Psychologically induced cooling of a specific body part caused by the illusory
18 ownership of an artificial counterpart. *Proceedings of the National Academy of*
19 *Sciences*, 105(35), 13169-13173.
20
21
22
23
24 Mountcastle, V. B. (2005). *The sensory hand: neural mechanisms of somatic sensation*. Harvard
25 University Press.
26
27
28 Natsoulas, T., & Dubanoski, R. A. (1964). Inferring the locus and orientation of the perceiver
29 from responses to stimulation of the skin. *The American journal of psychology*, 281-
30 285.
31
32
33
34 Pfeiffer, C., Lopez, C., Schmutz, V., Duenas, J.A., Martuzzi, R., & Blanke, O. (2013).
35 Multisensory origin of the subjective first-person perspective: visual, tactile, and
36 vestibular mechanisms. *PLoS One* 8, e61751.
37
38
39
40 Porro C.A., Martinig, M., Facchin, P., Maieron, M. Jones, A.K.P., & Fadiga L. (2007) Parietal
41 cortex involvement in the localisation of tactile and noxious mechanical stimuli: A
42 transcranial magnetic stimulation study. *Behavioural Brain Research*, 178:183–189.
43
44
45
46 Ramachandran, V. S., & McGeoch, P. (2007). Can vestibular caloric stimulation be used to treat
47 apotemnophilia?. *Medical hypotheses*, 69(2), 250-252.
48
49
50
51 Ramachandran, V. S., McGeoch, P. D., Williams, L., & Arcilla, G. (2007). Rapid relief of
52 thalamic pain syndrome induced by vestibular caloric stimulation. *Neurocase*. 13(3),
53 185-188.
54
55
56
57
58
59
60

- 1
2
3
4 Rapp, B., Hendel, S.K., & Medina J. (2002) Remodeling of somatosensory hand representations
5 following cerebral lesions in humans. *NeuroReport*, 13:207–211.
6
7
8 Rode, G., Charles, N., Perenin, M. T., Vighetto, A., Trillet, M., & Aimard, G. (1992). Partial
9 remission of hemiplegia and somatoparaphrenia through vestibular stimulation in a case
10 of unilateral neglect. *Cortex*, 28(2), 203-208.
11
12
13
14 Rode G., & Perenin M. T. (1994). Temporary remission of representational hemineglect
15 through vestibular stimulation. *Neuroreport*. 5, 869–872.
16
17
18 Rohde, M., Di Luca, M., & Ernst, M. O. (2011). The rubber hand illusion: feeling of ownership
19 and proprioceptive drift do not go hand in hand. *PloS one*, 6(6), e21659.
20
21
22 Romo, R., & Salinas, E. (2001). Touch and go: decision-making mechanisms in
23 somatosensation. *Annual review of neuroscience*, 24(1), 107-137.
24
25
26 Ronchi, R., Rode, G., Cotton, F., Farnè, A., Rossetti, Y., & Jacquin-Courtois, S. (2013).
27 Remission of anosognosia for right hemiplegia and neglect after caloric vestibular
28 stimulation. *Restor Neurol Neurosci*, 31(1), 19-24.
29
30
31
32 Sang, F. Y. P., Jauregui-Renaud, K., Green, D. A., Bronstein, A. M., & Gresty, M. A. (2006).
33 Depersonalisation/derealisation symptoms in vestibular disease. *Journal of Neurology,*
34 *Neurosurgery & Psychiatry*, 77(6), 760-766.
35
36
37
38 Schiff, N.D., & Pulver, M. (1999). Does vestibular stimulation activate thalamocortical
39 mechanisms that reintegrate impaired cortical regions? *Proc R Soc Lond B Biol Sci*
40 266, 421–3.
41
42
43
44 Schilder, P., 1935. *The image and appearance of the human body*. International Universities
45 Press, New York.
46
47
48 Schmidt, L., Artinger, F., Stumpf, O., & Kerkhoff, G. (2013b). Differential effects of galvanic
49 vestibular stimulation on arm position sense in right-vs. left-handers.
50 *Neuropsychologia*, 51(5), 893-899.
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 Schmidt, L., Keller, I., Utz, K. S., Artinger, F., Stumpf, O., & Kerkhoff, G. (2013c). Galvanic
5
6 Vestibular Stimulation Improves Arm Position Sense in Spatial Neglect A Sham-
7
8 Stimulation-Controlled Study. *Neurorehabilitation and neural repair*, 27(6), 497-506.
9
- 10 Schmidt, L., Utz, K. S., Depper, L., Adams, M., Schaadt, A. K., Reinhart, S., & Kerkhoff, G.
11
12 (2013a). Now you feel both: galvanic vestibular stimulation induces lasting
13
14 improvements in the rehabilitation of chronic tactile extinction. *Frontiers Human*
15
16 *Neuroscience*. 7.
17
- 18 Seyal, M., Siddiqui, I., & Hundal, N.S. (1997) Suppression of spatial localisation of a cutaneous
19
20 stimulus following transcranial magnetic pulse stimulation of the sensorimotor cortex.
21
22 *Electroencephalography and Clinical Neurophysiology*, 105:24–28.
23
- 24 Snyder, L. (1999). This way up: Illusions and internal models in the vestibular system. *Nature*
25
26 *Neuroscience*, 2, 396–398.
27
- 28 Stephan, T., Deutschländer, A., Nolte, A., Schneider, E., Wiesmann, M., Brandt, T., &
29
30 Dieterich, M. (2005). Functional MRI of galvanic vestibular stimulation with
31
32 alternating currents at different frequencies. *Neuroimage*, 26(3), 721-732.
33
- 34 Suzuki, M., Kitano, H., Ito, R., Kitanishi, T., Yazawa, Y., Ogawa, T., ... & Kitajima, K. (2001).
35
36 Cortical and subcortical vestibular response to caloric stimulation detected by
37
38 functional magnetic resonance imaging. *Cognitive Brain Research*, 12(3), 441-449.
39
- 40 Tsakiris, M., & Haggard, P. (2005). The rubber hand illusion revisited: visuotactile integration
41
42 and self-attribution. *Journal of Experimental Psychology: Human Perception and*
43
44 *Performance*, 31(1), 80-91.
45
- 46 Utz, K. S., Dimova, V., Oppenländer, K., & Kerkhoff, G. (2010). Electrified minds: transcranial
47
48 direct current stimulation (tDCS) and galvanic vestibular stimulation (GVS) as methods
49
50 of non-invasive brain stimulation in neuropsychology—a review of current data and
51
52 future implications. *Neuropsychologia*. 48(10), 2789-2810.
53
54
55
56
57
58
59
60

- 1
2
3
4 Utz, K. S., Keller, I., Kardinal, M., & Kerkhoff, G. (2011). Galvanic vestibular stimulation
5 reduces the pathological rightward line bisection error in neglect—a sham stimulation-
6 controlled study. *Neuropsychologia*, 49(5), 1219-1225.
7
8
9
10 Vallar, G., & Ronchi, R. (2009). Somatoparaphrenia: a body delusion. A review of the
11 neuropsychological literature. *Experimental Brain Research*, 192(3), 533-551.
12
13
14 Vallar, G., Bottini, G., Rusconi, M. L., & Sterzi, R. (1993). Exploring somatosensory
15 hemineglect by vestibular stimulation. *Brain*, 116(1), 71-86.
16
17
18 Vallar, G., Sterzi, R., Bottini, G., Cappa, S., & Rusconi, M. L. (1990). Temporary remission of
19 left hemianesthesia after vestibular stimulation. A sensory neglect phenomenon. *Cortex*.
20 26(1), 123-131.
21
22
23
24 Van Boven, R.W., Ingeholm, J.E., Beauchamp, M.S., Bikle, P.C., & Ungerleider L.G. (2005)
25 Tactile form and location processing in the human brain. *Proceedings of the National*
26 *Academy of Sciences of the United States of America*, 102:12600–12605.
27
28
29
30 Yardley, L. (1990). Contribution of somatosensory information to perception of the visual
31 vertical with body tilt and rotating visual field. *Perception & Psychophysics*, 48(2),
32 131-134.
33
34
35
36
37 Yardley, L. (2000). Overview of psychologic effects of chronic dizziness and balance disorders.
38 *Otolaryngologic Clinics of North America*, 33(3), 603-616.
39
40
41 Zu Eulenburg, P., Caspers, S., Roski, C., & Eickhoff, S. B. (2012). Meta-analytical definition
42 and functional connectivity of the human vestibular cortex. *Neuroimage*. 60(1), 162-
43 169.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 **Figure caption**
5
6
7

8 **Figure 1. Vestibular contributions to bodily representations.**
9

10 A schematic of the vestibular contributions to bodily representations. According with
11 Longo at al.'s (2010) model, we distinguish between three hierarchical levels of body
12 representation. These are (i) *somatosensation*, the primary sensory processing of somatic
13 stimuli, (ii) *somatoperception*, the processes of constructing percepts and experiences of
14 somatic objects and events and (iii) *somatorepresentation*, the knowledge about the body as a
15 physical object in the world. Vestibular signals independently contribute to low-level signal
16 processing and also to higher-order representations of the body.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Vestibular contributions to bodily representations. A schematic of the vestibular contributions to bodily representations. According with Longo et al.'s (2010) model, we distinguish between three hierarchical levels of body representation. These are (i) somatosensation, the primary sensory processing of somatic stimuli, (ii) somatopercception, the processes of constructing percepts and experiences of somatic objects and events and (iii) somatopercception, the knowledge about the body as a physical object in the world. Vestibular signals independently contribute to low-level signal processing and also to higher-order representations of the body.