

Comparison of Rectus Femoris Cross-sectional Area and Rectus Femoris-Vastus Intermedius Muscle Layer Thickness as Markers of Muscle Wasting and Weakness During Early Critical Illness

Puthuchearu ZA, PhD^{1,2,3}, McNelly AS, PhD^{2,4}, Rawal J, MRCS⁴, Connolly B, PhD^{5,6,8}, Sidhu PS FRCR⁷, Rowlerson A PhD⁸, Moxham J, MD⁹, and Harridge SD PhD⁸, Hart N, PhD^{6,9}, Montgomery HE, MD^{2,4}.

¹Division of Critical Care, University College London Hospitals NHS Foundation Trust, London, UK; ²NIHR Comprehensive Biomedical Research Centre, University College London Hospitals, London, UK; ³Division of Respiratory and Critical Care, University Medicine Cluster, National University Health System Singapore; ⁴Institute of Sport, Exercise and Health, UCL/University College London Hospitals, London, UK; ⁵NIHR Biomedical Research Centre, Guy's & St Thomas' NHS Foundation Trust and King's College London, London, UK; ⁶Lane Fox Clinical Respiratory Physiology Research Centre, Guy's & St Thomas' NHS Foundation Trust, London, UK; ⁷Department of Radiology, King's College Hospital NHS Foundation Trust, London, UK; ⁸Centre of Human and Aerospace Physiological Sciences, King's College London, London, UK; ⁹Department of Asthma, Allergy and Lung Biology, King's College London, London, UK.

Word Count 988

Corresponding author details:

Dr Zudin Puthuchearu, PhD

Institute of Human Health and Performance, University College London

1st Floor, Institute for Sport, Exercise and Health, 170 Tottenham Court Road,

London, W1T 7HA.

Tel: +44 203 447 2843/ +44 7767 357983; Fax: +44 203 447 2898

zudin.puthuchearu.09@ucl.ac.uk

Reprints will not be available.

Author Contributions:

Concept and design: ZP, PS, JM, SH, NH, HM. Data Collection: ZP, AM, JR, BC, AR. Analysis and Interpretation: ZP, AM, BC, PS, AR, JM, SH, NH, HM. Manuscript drafting and revision: ZP, AM, BC, PS, AR, JM, SH, NH, HM.

Conflicts of Interest and Source of Funding

The UK National Institute of Health Research (NIHR) funded ZP. AM was supported by the Batchworth Charitable Trust, the Moulton Charitable Foundation and the NIHR University College London Hospitals Biomedical Research Centre (UCLH BRC). SH received support from the Research Councils UK. NH and BC received funding from the NIHR Clinical Research Facility and BRC at Guy's and St Thomas' NHS Foundation Trust (GSST) and King's College London. HM was funded by University College London and UCLH BRC. The NIHR doctorate fellowship (£479,000; 2010–2014) underpinned the core patient population study on which this follow-up was built; Batchworth Charitable Trust (£15,000; January 2013 to July 2014); Moulton Foundation (£45, 000; August 2014 to June 2017); and European Society of intensive Care Medicine (£18,000; Single donation 2010). Additional funding was received from the Whittington Hospital NHS Trust and the European Society of Intensive Care Medicine. The authors do not recognise any conflicts of interest.

Key Words: Intensive Care; Critical Illness; Ultrasound; Muscle Wasting; Outcome Assessment

Muscle wasting during critical illness has been suggested to contribute to survivor functional disability(1). Two B-mode ultrasound measures measurements have been reported that quantify wasting(2, 3): (i) combined thickness of the Rectus Femoris (RF) and Vastus Intermedius muscles ('Muscle Layer Thickness', henceforth referred to as 'Thickness') (4, 5) and (ii) RF cross-sectional area (RF_{CSA}) which correlates with lower limb strength in other clinical circumstances (6). The degree to which either of these ultrasound measures reflect muscle weakness in the critically ill is unclear (7). We hypothesised that like change in RF_{CSA} (ΔRF_{CSA}), change in Thickness (Δ Thickness) would underestimate loss of muscle size as measured by the histological gold standard (myofibre thickness) and the biochemical gold standard of protein: DNA ratio measured in skeletal muscle biopsies. Secondly we hypothesised that ΔRF_{CSA} and Δ Thickness would both be related to muscle weakness.

Subjects were patients of the Musculoskeletal Ultrasound in Critical Illness: Longitudinal Evaluation study (NCT01106300) (8), the original study having been approved by University College London Ethics Committee A. All patients were recruited within 24 hours of admission to a university hospital and a community hospital (August 2009-April 2011), and were expected to survive intensive care unit (ICU) admission after being invasively ventilated for > 48 hours and in the ICU >7 days. Excluded were those with pregnancy, lower limb amputation, primary neuromuscular pathology or disseminated cancer. Next-of-kin assent and retrospective patient consent were obtained.

Images were acquired on ICU days 1, 7 and 10. ICU RF_{CSA} assessment and reliability have been previously described (8). Thickness was measured at the midpoint of Rectus Femoris between the two fascial lines. Images were excluded where the femur was not visible.

Δ Thickness and ΔRF_{CSA} were compared with change in myofibre cross-sectional area (Δ fibre $_{CSA}$) and protein:DNA in sequential Vastus Lateralis muscle biopsies acquired on days 1 and 7 as described previously (8).

Manual Muscle Testing was performed (9) on day 10 if patients could follow ≥ 3 of De Jonghe's 5-command criteria and the knee extension component score of $\leq 4/5$ used to define lower limb weakness (10).

Bland-Altman comparisons were used to establish i) inter-rater reliability of Thickness measurements and ii) longitudinal bias between Δ Thickness and Δ RF_{CSA} over the study period. Normality was assessed using D'Agostino and Pearson omnibus normality tests, and data were analysed using two-tailed Student's t-test or Mann Whitney U test as appropriate. Differential longitudinal change in muscle size (Δ Thickness vs. Δ RF_{CSA}) was compared using 2-way repeated measures of variance (ANOVA). A bivariable logistical regression was performed with knee extensor weakness as the dependent variable and ultrasound measurements as the independent variable.

Of the initial cohort of 62 patients with serial muscle ultrasounds, 8 had incomplete or missing electronic scan records. Of the remaining 54, 11 had ≥ 1 scan where the femur was not visualized. Two assessors analysed images at 21 time-points to establish inter-rater reliability. Thickness measurements were highly correlated between observers (AM and ZP: Pearson $r=0.98$) with an intra-class co-efficient of 0.986 (95%CI 0.965-0.994). A Bland Altman plot demonstrated minimal bias of -0.07 ± 0.2 cm (95%CI -0.46-0.32 cm).

Nineteen patients had Thickness, RF_{CSA}, fibre_{CSA} and protein/DNA ratio measured on Day 1 and Day 7. Δ Thickness significantly underestimated Δ fibre_{CSA} (-4.6% (95%CI -14.19-4.95) vs. -16.4% (95%CI -32.0—0.74); $p=0.025$) and change in protein/DNA ratio ((-4.6% (95%CI -14.19-4.95) vs. -30.9% (95%CI -51.2—10.6); $p=0.019$). We have previously shown Δ RF_{CSA} to underestimate change in protein/DNA ratio (10.3% (95%CI 6.1-14.5) vs. 29.5% (95%CI 13.4-45.6%; $p=0.03$) but not Δ fibre_{CSA} (10.3% (95%CI 6.1-14.5) vs. 17.5% (95%CI 5.8-29.3); $p=0.31$) (8).

Δ Thickness and Δ RF_{CSA} correlated ($r^2=0.22$, $p=0.049$) but a Bland Altman comparison between Δ Thickness and Δ RF_{CSA} over 10 days revealed a bias of $-8.3\% \pm 19.7\%$ (95% CI-46.7-30.7) for Thickness resulting in significant underestimation of muscle wasting at days 7 and 10 (Figure 1A and table 1).

Of the 63 patients, 40 were able to obey commands and underwent volitional strength testing on Day 10, amongst whom Thickness was available in 27.

Δ RF_{CSA} was greater in those with knee extensor weakness than those without (20.7% (95%CI 13.7-27.7) vs. 8.4% (95%CI 2.5-14.3) respectively $p=0.012$). Δ Thickness did not differ between these groups (12.6% (95%CI 0.94-24.2) vs. 12.1 (95%CI 2.7-21.5)

respectively, $p=0.95$) (Figure 1B). In a bivariable logistical regression, ΔRF_{CSA} was associated with knee extensor weakness (OR 1.101 (95%CI 1.011-1.199); $p=0.027$), but $\Delta Thickness$ was not (OR 1.001 (95%CI 0.960-1.044); $p=0.947$).

All other things being equal, muscle strength and size are proportional - the latter acting as a proxy for the former in ICU, where non-volitional objective measures of strength are logistically challenging. Our results suggest that ΔRF_{CSA} reflects knee extensor weakness and muscle loss better than $\Delta Thickness$. $\Delta Thickness$ also underestimated ΔRF_{CSA} (a -8% bias on Bland Altman plot being relevant, given that a 10% change in RF_{CSA} is considered sufficient to affect function(11))- in part, perhaps, because it is a unidimensional measure when compared to (2D) muscle area or (3D) volume. The specific relationship of tissue edema to ultrasound measures remains unclear (3, 8), though edema may also affect $Fibre_{CSA}$ (12).

Although these data are derived from the largest cohort available for longitudinal radiopathological correlation, our study is limited by its size. The cohort size was further limited by a third of patients not being able to perform volitional strength testing, albeit this being in keeping with published rates (13). Finally, measurement of Thickness was not an original primary goal of image analysis, a fact which might account for the lack of femoral image availability in one third. Although considered unlikely to have impacted on the observations made, non-random bias cannot be excluded.

We have previously shown RF_{CSA} studies to indicate muscle quality (3) and not to underestimate muscle $fibre_{CSA}$. We now show that Thickness measurements significantly underestimate ICU muscle wasting compared to RF_{CSA} . In addition RF_{CSA} is a more reliable proxy for muscle strength in a setting where volitional and non-volitional muscle strength measurements are challenging. We suggest measurement of ΔRF_{CSA} as a biomarker for proximal lower limb muscle loss and knee extensor weakness during early critical illness.

REFERENCES

1. Herridge MS, Cheung AM, Tansey CM, Matte-Martyn A, Diaz-Granados N, Al-Saidi F, Cooper AB, Guest CB, Mazer CD, Mehta S, Stewart TE, Barr A, Cook D,

- Slutsky AS. One-year outcomes in survivors of the acute respiratory distress syndrome. *N Engl J Med* 2003; 348: 683-693.
2. Parry SM, El-Ansary D, Cartwright MS, Sarwal A, Berney S, Koopman R, Annoni R, Puthuchery Z, Gordon IR, Morris PE, Denehy L. Ultrasonography in the intensive care setting can be used to detect changes in the quality and quantity of muscle and is related to muscle strength and function. *J Crit Care* 2015; 30: 1151 e1159-1151 e1114.
 3. Puthuchery ZA, Phadke R, Rawal J, McPhail MJ, Sidhu PS, Rowleron A, Moxham J, Harridge S, Hart N, Montgomery HE. Qualitative ultrasound in acute critical illness muscle wasting. *Crit Care Med* 2015; 43: 1603-1611.
 4. Reid CL, Campbell IT, Little RA. Muscle wasting and energy balance in critical illness. *Clin Nutr* 2004; 23: 273-280.
 5. Gruther W BT, Zorn C, Paternostro-Sluga T, Quittan M, Fialka-Moser V, Spiss C, Kainberger F, Crevenna R. Muscle wasting in intensive care patients: Ultrasound observation of the m. Quadriceps femoris muscle layer. *J Rehabil Med* 2008; 40: 185-189.
 6. Seymour JM, Ward K, Sidhu P, Puthuchery Z, Steier J, Jolley C, Rafferty G, Polkey MI, Moxham J. Ultrasound measurement of rectus femoris cross-sectional area and the relationship to quadriceps strength in chronic obstructive pulmonary disease. *Thorax* 2009; 64: 418.
 7. Connolly B, MacBean V, Crowley C, Lunt A, Moxham J, Rafferty GF, Hart N. Ultrasound for the assessment of peripheral skeletal muscle architecture in critical illness: A systematic review. *Crit Care Med* 2015; 43: 897-905.
 8. Puthuchery ZA, Rawal J, McPhail M, Connolly B, Ratnayake G, Chan P, Hopkinson NS, Padhke R, Dew T, Sidhu PS, Velloso C, Seymour J, Agle CC, Selby A, Limb M, Edwards LM, Smith K, Rowleron A, Rennie MJ, Moxham J, Harridge SD, Hart N, Montgomery HE. Acute skeletal muscle wasting in critical illness. *JAMA* 2013; 310: 1591-1600.
 9. Kleyweg RP, van der Meche FG, Schmitz PI. Interobserver agreement in the assessment of muscle strength and functional abilities in guillain-barre syndrome. *Muscle Nerve* 1991; 14: 1103-1109.
 10. Connolly BA, Jones GD, Curtis AA, Murphy PB, Douiri A, Hopkinson NS, Polkey MI, Moxham J, Hart N. Clinical predictive value of manual muscle strength testing during critical illness: An observational cohort study. *Crit Care* 2013; 17: R229.
 11. Seymour JM, Spruit MA, Hopkinson NS, Natanek SA, Man WD, Jackson A, Gosker HR, Schols AM, Moxham J, Polkey MI, Wouters EF. The prevalence of quadriceps weakness in copd and the relationship with disease severity. *Eur Respir J* 2010; 36: 81-88.
 12. Hauptmann S, Klosterhalfen B, Weis J, Mittermayer C, Kirkpatrick CJ. Skeletal muscle oedema and muscle fibre necrosis during septic shock. Observations with a porcine septic shock model. *Virchows Arch* 1994; 424: 653-659.
 13. Hough C, Lieu B, Caldwell E. Manual muscle strength testing of critically ill patients: Feasibility and interobserver agreement. *Crit Care* 2011; 15: R43.

Figures and Tables




Figure 1AB: (A) Change in Rectus Femoris Cross Section Area (RF_{CSA}) and Muscle Layer Thickness (Thickness) over 10 days of critical illness. * Represents $p<0.05$ and ** $p<0.01$ using Two-way repeated measures Analysis of Variance (ANOVA). (B) Knee Extensor Medical Research Council (MRC) Strength Score and loss of muscle size as measured by Rectus Femoris Cross Sectional Area (RF_{CSA}) and Muscle Layer Thickness (Thickness) ($n=27$). *Represents $p<0.05$ using 2-tailed unpaired Student's T-test.

	Δ THICKNESS	Δ RF_{CSA}	
Day 7	-5.88% (-11.69—0.06%)	-13.0%(-16.52—9.48%)	0.031*
Day 10	-9.36% (-15.43—3.84%)	-17.72 (-21.15—14.29)	0.004*

Table 1: Comparison of change in Muscle Limb Thickness (Δ THICKNESS) and Rectus Femoris Cross Sectional Area (Δ RF_{CSA}) at days 7 and 10 of critical illness. *Represents $p<0.05$ using 2-way repeated measures Analysis of Variance (ANOVA).