

High-resolution absorption measurements of NH₃ at high temperatures: 2100 - 5500 cm⁻¹

Emma J. Barton, Sergei. N. Yurchenko, Jonathan Tennyson

Department of Physics and Astronomy, University College London, London, WC1E 6BT, UK

Sønnik Clausen, Alexander Fateev

Technical University of Denmark, Department of Chemical and Biochemical Engineering, Frederiksborgvej 399, 4000 Roskilde, Denmark

Abstract

High-resolution absorption spectra of NH₃ in the region 2100 - 5500 cm⁻¹ at 1027 °C and approximately atmospheric pressure (1045 ± 3 mbar) are measured. An NH₃ concentration of 10% in volume fraction is used in the measurements. Spectra are recorded in a high-temperature gas-flow cell using a Fourier Transform Infrared (FTIR) spectrometer at a nominal resolution of 0.09 cm⁻¹. The spectra are analysed by comparison to a variational line list, BYTe, and experimental energy levels determined using the MARVEL procedure. 2308 lines have been assigned to 45 different bands, of which 1755 and 15 have been assigned or observed for the first time in this work.

Key words:

High temperature, Ammonia, Absorption, FTIR spectroscopy,
High-temperature flow gas cell, BYTe, line assignments

1. Introduction

NH₃ spectra can be used to extract physical information from spectroscopic observations of a range of hot and cold environments. On Earth NH₃ is an important component in several industrial processes as for example gasification and NO_x reduction in combustion [1]. Such processes can be monitored and optimised with the help of *in situ* measurement of gas temperature and composition [2]. In space NH₃ is ubiquitous and used to probe, for example, circumstellar envelopes [3], star-forming regions [4], dense molecular clouds [5], the atmospheres of cool stars [6], brown dwarfs [7] and giant solar system planets [8]. Recent work includes the first detection of gas-phase ammonia in a planet-forming disk [9].

Many experimental studies have focused on the NH₃ molecule providing, for example, high temperature experimental line lists [10, 11], ro-vibrational assignments [12, 13, 14, 15] and experimentally derived energies [10, 11, 12, 13, 14, 16]. A comprehensive compilation of measured NH₃ rotational and ro-vibrational spectra can be found in a recent MARVEL study [17]. The MARVEL (measured active rotation-vibration energy levels) algorithm [18, 19] simultaneously analyses all available assigned and labelled experimental lines, thus yielding the associated energy levels. The recent study for NH₃ analysed 29,450 measured transitions and yielded 4961 accurately-determined energy levels which mostly lie below 7000 cm⁻¹ [17]. Very recently Sung *et al* [20] have significantly improved the spectral coverage for ammonia in the far infrared.

The broad temperature and spectral range of applications can be difficult to cover exhaustively in the lab because of NH₃ thermal decomposition either

in the gas phase or on the walls of a gas cell (heterophase). To help fill in the gaps a number of theoretical line lists have been computed for NH_3 [21, 22, 23]. In the present work a variationally computed line list for hot NH_3 , BYTe [22], is employed. This line list covers the spectral range 0 - 12,000 cm^{-1} and is expected to be fairly accurate for all temperatures up to 1500 K (1226 °C). It comprises of 1,138,323,251 transitions constructed from 1,373,897 energy levels lying below 18 000 cm^{-1} . It was computed using the NH3-2010 potential energy surface [24], the TROVE ro-vibrational computer program [25] and an *ab initio* dipole moment surface [21]. However a new line list currently being constructed as part of the ExoMol project [26, 27] as BYTe is known to have some problems reproducing experimental intensities [13, 14] and is less accurate for higher wavenumber transitions [14, 28, 29, 30]. Assigned high resolution laboratory spectra are needed to refine and validate theoretical line positions and intensities.

In our previous study [13] we extended work by Zobov *et al.* [12] by analysing new hot absorption spectra in the region 500 - 2100 cm^{-1} . In the current work we present and analyse new hot absorption spectra in the region 2100 - 5500 cm^{-1} . It should be noted that high temperature (up to 1400 °C) experimental line lists for the region 2100 - 4000 cm^{-1} are available due to Hargreaves *et al.* [11] based on their observed emission spectra.

This article has the following structure. The experimental set-up used for the measurements is described in Section 2. Section 3 gives an overview the assignment procedure and the method used to calculate experimental and theoretical absorbance spectra. The accuracy of BYTe is assessed by a direct comparison with the experimental spectra in Section 4.1 and summary of all

assignments is given in Section 4.2. Finally our conclusions are presented in Section 5.

2. Experimental Details

The experimental setup is described in our previous work [13], the main points are summarised below.

An Agilent 660 FTIR spectrometer, linearised Mercury-Cadmium Telluride (MCT) detector, ceramic high-temperature gas-flow cell (c-HGC) (see Figure 1) and an external IR light source, which is Blackbody-like (BB) at 1800 K were used in the measurements. The optical setup is illustrated in Figure 2.

The c-HGC operates at temperatures up to 1873 K (1600 °C) [31] and has also been used by the DTU group [32, 33, 34] to study for example hot CO, CO₂, CH₄ and H₂O. This cell has a fully-heated, temperature-uniform central part and two partially-heated buffer parts with interchangeable optical (KBr) windows at the ends. The buffer parts are purged with N₂ or dry air taken from a purge generator while the sample gas (e.g. N₂ + NH₃) is preheated and fed into the central part of the cell. Laminar flow sheets (flow windows) are established between the central and buffer parts where the purge and sample gases meet, meaning the sample gas can not reach, or react with or form deposits on, the optical windows [35]. To minimise reactions with the internal surface of the gas cell the inner part of the c-HGC is made from high quality pure ceramic (Al₂O₃(99.5%)). The absorption path length, defined by the flow windows, has a value of 53.3 cm at room temperature. At higher temperatures the length changes a little due to thermal expansion, see [33].

Bottles of premixed gas mixture of $\text{N}_2 + (98.3 \pm 2.0\%) \text{NH}_3$ were made using N_2 and NH_3 from Air Liquide, and 'pure' NH_3 were obtained from Linde Gas. The purity of NH_3 in the gas bottles was 99.98% with air or H_2O being the main impurity. High purity N_2 (99.998%) has been used in reference measurements. NH_3 (10%) was obtained through mixture of N_2 (99.998%) and NH_3 (99.98%) flows controlled by high-end mass flow controllers (BRONKHORST).

Further details on the c-HGC, its performance and a comparison with the other HGCs in the laboratory will be presented elsewhere [31]. For now the reader is referred to Ref. [34].

Single beam (SB) spectra from measured interferograms at a nominal spectral resolution 0.09 cm^{-1} are calculated using Agilent Resolution Pro software (supplied with the FTIR spectrometer) using inverse fast Fourier transform (FFT) and boxcar and triangular apodization functions. Mertz phase correction is applied. Triangular apodization results in less noise in the final spectra while boxcar apodization gives narrower peaks. To ensure consistent results both sets of calculated SB spectra were used in the final analysis.

Measured wavenumbers were multiplied by a factor of 1.000059 to account for the linear wavenumber shift caused by beam divergence, in accordance with the discussion in [34]. The experimental uncertainties on absorbance measurements are estimated to be within 0.5% [34].




Figure 1: High temperature ceramic gas flow cell (c-HGC) used in the experiments. Black arrows indicate flow direction. Reproduced from [34].




Figure 2: Experimental setup for the high-resolution measurements of NH_3 at high temperatures. Adapted from [34].

3. Data Analysis

This study used the BYTe [22] variational line list and experimental energies determined using the MARVEL procedure [17].

3.1. Calculating Experimental Absorption Spectra

Experimental transmission spectra $\tau_{\text{exp}}(\nu, T)$ at a temperature T [K] and a line position ν [cm^{-1}] are calculated from four SB spectra, two reference (N_2 in the central part of the cell) measurements $\mathbf{I}_{\text{ref+BB}}$ and \mathbf{I}_{ref} and two sample ($\text{N}_2 + \text{NH}_3$ mixture) measurements $\mathbf{I}_{\text{gas+BB}}$ and \mathbf{I}_{gas} , one with and one without signal from the BB (at 1800 K) [13, 32]:

$$\tau_{\text{exp}}(\nu, T) = \frac{\mathbf{I}_{\text{gas+BB}} - \mathbf{I}_{\text{gas}}}{\mathbf{I}_{\text{ref+BB}} - \mathbf{I}_{\text{ref}}} \quad (1)$$

Spectra without signal from the BB are measured from a cold (room temperature) beam stopper placed at 90 degrees from the optical axis of the setup using a movable mirror in the BB adapter. The absorption spectra are then calculated from the reference, a_0 ($= \mathbf{I}_{\text{ref+BB}} - \mathbf{I}_{\text{ref}}$), and sample, a_1 ($= \mathbf{I}_{\text{gas+BB}} - \mathbf{I}_{\text{gas}}$), measurements:

$$A_{\text{exp}}(\nu, T) = \log_{10} \left[\frac{a_0}{a_1} \right] \quad (2)$$

3.2. Calculating Theoretical Absorption Spectra

The method for calculating theoretical absorption spectra follows Ref. [13]. First the 'true' transmission spectrum was computed as:

$$\tau_{\text{calc}}^{\text{true}}(\nu, T) = \exp(-\sigma(\nu, T)lc) \quad (3)$$

where l is the absorption path length in cm, c is the NH_3 concentration in cm^{-3} and $\sigma(\nu, T)$ is the pressure-broadened NH_3 absorption cross-section calculated using BYTe and the procedure laid out in [36], but replacing the Gaussian line shape with a Voigt line shape. Lorentz half-widths were estimated from the experimental spectra and with reference to measured widths compiled in the HITRAN database. The measured (effective) transmittance spectrum is derived by convolving $\tau_{\text{calc}}^{\text{true}}(\nu, T)$ with the instrument line shape (ILS) function $\Gamma(\nu - \nu_0)$:

$$\tau_{\text{calc}}^{\text{eff}}(\nu, T) = \int_0^\infty \tau_{\text{calc}}^{\text{true}}(\nu_0, T) \Gamma(\nu - \nu_0) d\nu_0 \quad (4)$$

For boxcar apodization, the ILS is a sinc function:

$$\Gamma(\nu) = \Lambda \text{sinc}(\Lambda\pi\nu) = \Lambda \frac{\sin(\Lambda\pi\nu)}{(\Lambda\pi\nu)} \quad (5)$$

For triangular apodization, the ILS is a sinc^2 function:

$$\Gamma(\nu) = \Lambda \text{sinc}^2(\Lambda\pi\nu) = 2\Lambda \frac{\sin^2(\Lambda\pi\nu)}{(\Lambda\pi\nu)^2} \quad (6)$$

where Λ is commonly termed the FTIR retardation and is generally defined as the inverse of the nominal resolution of the spectrometer [37].

The theoretical absorption spectrum is then computed as:

$$A_{\text{calc}}(\nu, T) = \log_{10} \left[\frac{1}{\tau_{\text{calc}}^{\text{eff}}(\nu, T)} \right] \quad (7)$$

3.3. The Assignment Procedure

First a list of observable BYTe lines for the experimental conditions was compiled. For this purpose the absorbance of each line, j , was approximated

as:

$$A_{\text{calc}}^{\text{approx}} = \frac{S_j^a l c}{\Delta L \ln(10)} \quad (8)$$

where ΔL is an effective line width which is assumed to be a constant for all lines in the spectrum and the quantity $\frac{S_j^a}{\Delta L}$ represents an effective cross section assuming rectangular line shapes with ΔL widths.

If both the upper and lower energies involved in a observable transition were known experimentally, the BYTe line position was replaced by the MARVEL line position generated by subtracting upper and lower state energies. This hybrid line list shall henceforth be referred to as "BARVEL" (Mr. Philip Coles).

Taking the resolution of the measurements and the accuracy of BYTe intensities into account (see Section 4.1), experimental peaks and BARVEL line positions were coupled using python scripts to produce a 'trivial' assignment list. In cases where multiple BARVEL lines corresponded to a single peak, the peak was assigned to the strongest line.

Trivial assignments for the same vibrational band provide an expected observed minus calculated (obs. - calc.) difference for all lines in that band. Lines present in the list of observable BYTe lines, but not in BARVEL, were shifted by this residual to make future assignments by the method of branches [38].

A list of all trivial and branch assignments, the final assignment list, was then compared to previous studies, namely those catalogued in the HITRAN database [39].

4. Results and Discussion

The absorption measurements were performed at a temperature of 1027 °C for the NH₃ volume concentration of 10%.

The measurements were used to test the accuracy of BYTe then analysed using BYTe to generate an assignment list for the data. Central wavenumbers for assigned peaks are compared to line positions measured by Hargreaves et al. [11] where possible.

The absorption spectra, a peak list (partially assigned) including line positions from Hargreaves et al. [11] for assigned lines where available, and new energy level information derived from the assignments are presented in the supplementary data.

4.1. Direct Comparison with BYTe

A comparison between the experimental and theoretical absorption spectra at 1027 °C for the whole region (2100 - 5500 cm⁻¹) is shown in Figure 3. Overall, taking into account the experimental noise, there is good agreement. However there are shifts in line position of the order ± 0.2 cm⁻¹ across the entire spectral range and shifts up to $\pm 1 - 2$ cm⁻¹ in a few regions, particularly at higher wavenumbers. Hence it was decided that assignments should only be made using MARVEL line positions or BYTe line positions corrected for the expected obs. - calc. difference derived from trivial assignments, and not by simple line list comparison. BARVEL line positions should have an obs. - calc. difference smaller than the nominal resolution of the measurements, 0.09 cm⁻¹, whilst the wavenumber threshold for the BYTe line positions was taken to within 0.1 cm⁻¹ of the expected obs. - calc. difference. On the

whole experimental line intensities are reproduced within 30 %. This is illustrated in Figure 4 for the region 4860 - 4900 cm^{-1} . As such experimental lines were coupled to BARVEL or BYTe lines using an intensity threshold of 30 %.




Figure 3: Comparison between experimental (upper) and calculated (BYTe, lower) absorption spectra at 500 °C for the range 2100 - 5500 cm^{-1} .

4.2. Assignments

Out of 3701 measured experimental peaks 2308 lines have been assigned. The remaining peaks either did not correspond to a BARVEL or BYTe line




Figure 4: Comparison between experimental (upper) and calculated (BYTe, lower) absorption spectra at 1027 °C for the range 4860 - 4900 cm⁻¹.

within the set wavenumber and intensity thresholds or corresponded to multiple lines with roughly equal contribution to the total intensity such that it could not be confidently assigned. 553 lines were previously assigned by studies included in the HITRAN database (see Table 1). The full 1027 °C (1300 K) peak list with assignments is available as supplementary material to this article.

Hargreaves et al. [11] presented high temperature line lists for the region 2100 - 4000 cm^{-1} constructed from emission spectra recorded at a resolution of 0.01 cm^{-1} . These line lists are currently being updated and extended (Rob Hargreaves and Peter Bernath, private communication) and hence were not the focus of the current work. Of the 1755 newly assigned lines in this work, 990 are also present in the line lists of Ref. [11]. In these cases line positions from Ref. [11] are included with the current central peak wavenumbers in the supplementary data and employed in the computation of upper state energies described below, as these were measured at a higher resolution.

For branch assignments with an experimentally known lower energy state, energies for the upper state were computed using MARVEL energies and the line position of the strongest assigned transition to that state. The calculated energies are available as supplementary material to this article.

As in our previous study [13], lines were assigned to a large number of different bands. Table 2 gives a summary of the observed bands including the number of lines assigned to each and whether the band was observed for the first time in this work. Bands are listed in order of theoretical vibrational band centre (VBC). $\text{VBC} = \text{VBO}' - \text{VBO}''$ where VBO is the vibrational band origin from BYTe, in wavenumbers. For simplicity abbreviated vibrational

Table 1: Summary of NH₃ lines assigned in the region 2100 - 5500 cm⁻¹.

	Lines
Experimental	3701
HITRAN	553
New trivial	272
New branch	1483
Total Assigned	2308

labels $(v_1v_2v_3^{L_3}v_4^{L_4}i)$ [40] are used to identify bands in this table and only the highest value of the rotational quantum number J , assigned in this work for each band, is indicated. If the observed J_{\max} in this work is bigger that quoted in the literature, the previous J_{\max} is also given. The full 26 quantum labels for each transition, 13 per vibration-rotation state as recommended by Down *et al.* [40], will be given in the partially assigned peak list and energies files.

Table 2: Summary of observed bands in the region 2100 - 5500 cm^{-1} in order of theoretical (BYTe) vibrational band centre (VBC = $\text{VBO}' - \text{VBO}''$ where VBO = vibrational band origin, in cm^{-1}) with maximum upper and lower J rotational quantum number (J'_{max} and J''_{max} respectively). N is the number of lines assigned to the band. If J_{max} in this work is higher than given in the literature, the previously known J_{max} is given in parentheses. VBO of 0^+ is set to 0.000000 cm^{-1} in line with the MARVEL study [17].

Band	VBC	N	J'_{max}	J''_{max}	Note
$\nu_3^{1,-} - \nu_2^-$	2475.50	59	17 (12)	16	
$\nu_3^{1,+} - \nu_2^+$	2511.55	52	19 (12)	18	
$(\nu_2 + \nu_3^1)^- - 2\nu_2^-$	2553.27	14	16 (11)	15	New Band
$(\nu_2 + \nu_3^1)^+ - 2\nu_2^+$	2553.27	6	9	9	New Band
$3\nu_2^- - 0^+$	2895.53	6	9	9	
$(\nu_1 + 2\nu_2)^+ - 2\nu_2^-$	3120.69	8	16 (11)	16	
$(\nu_2 + 2\nu_4^0)^+ - \nu_2^+$	3147.49	1	8	8	New Band
$(\nu_2 + 2\nu_4^2)^+ - \nu_2^+$	3167.81	19	18 (7)	19	New Band
$(2\nu_2 + \nu_4^1)^+ - 0^+$	3189.04	17	15 (11)	14	
$2\nu_4^{0,+} - 0^-$	3215.21	95	21 (12)	20	
$2\nu_4^{0,+} - 0^+$	3216.00	2	6	7	
$2\nu_4^{0,-} - 0^-$	3216.75	13	17 (12)	18	

Continued on next page

Table 2 – continued from previous page

Band	VBC	N	J'_{\max}	J''_{\max}	Note
$2\nu_4^{0,-} - 0^+$	3217.55	72	18 (12)	17	
$2\nu_4^{2,+} - 0^-$	3239.39	5	9	8	
$2\nu_4^{2,+} - 0^+$	3240.18	82	24 (13)	25	
$2\nu_4^{2,-} - 0^-$	3240.78	53	21 (13)	22	
$(\nu_2 + 2\nu_4^0)^- - \nu_2^-$	3240.81	1	7	8	New Band
$2\nu_4^{2,-} - 0^+$	3241.58	23	17 (13)	18	
$(\nu_2 + 2\nu_4^2)^- - \nu_2^-$	3260.70	8	16 (8)	17	New Band
$2\nu_1^+ - 2\nu_4^{0,-}$	3296.58	1	4	4	New Band
$(\nu_1 + \nu_2)^+ - \nu_2^-$	3326.39	62	22 (11)	22	
$(\nu_1 + \nu_4^1)^+ - \nu_4^{1,-}$	3328.35	19	14 (13)	15	New Band
$(\nu_1 + \nu_4^1)^- - \nu_4^{1,-}$	3329.51	11	15 (12)	16	New Band
$(\nu_1 + \nu_4^1)^- - \nu_4^{1,+}$	3330.61	10	13 (12)	14	New Band
$\nu_1^+ - 0^-$	3335.28	158	21 (12)	21	
$\nu_1^- - 0^+$	3337.07	155	21 (12)	21	
$(\nu_1 + \nu_2)^- - \nu_2^+$	3387.59	85	19 (12)	19	
$\nu_3^{1,-} - 0^-$	3443.20	142	22 (12)	22	
$(\nu_3^1 + \nu_4^1)^+ - \nu_4^{1,+}$	3443.60	2	6	7	New Band
$\nu_3^{1,+} - 0^+$	3443.62	160	23 (12)	24	
$(\nu_2 + \nu_3^1)^+ - \nu_2^+$	3448.80	110	19 (12)	20	
$(\nu_2 + \nu_3^1)^- - \nu_2^+$	3467.32	1	7	7	
$(\nu_2 + \nu_3^1)^- - \nu_2^-$	3503.01	73	19 (11)	20	
$(2\nu_2 + \nu_3^1)^- - 2\nu_2^-$	3470.63	18	14 (7)	15	New Band

Continued on next page

Table 2 – continued from previous page

Band	VBC	N	J'_{\max}	J''_{\max}	Note
$(2\nu_2 + \nu_3^1)^+ - 2\nu_2^+$	3548.80	74	21 (11)	22	New Band
$(2\nu_2 + \nu_3^1)^+ - \nu_2^+$	4178.16	89	18 (11)	17	New Band
$(\nu_1 + \nu_2)^+ - 0^-$	4293.73	11	12 (11)	12	
$(\nu_1 + \nu_2)^- - 0^+$	4320.03	12	13 (12)	12	
$(\nu_2 + \nu_3^1)^+ - 0^+$	4416.93	123	19 (12)	18	
$(2\nu_2 + \nu_3^1)^- - \nu_2^-$	4420.38	93	16 (7)	17	New Band
$(\nu_2 + \nu_3^1)^+ - 0^-$	4434.66	129	19 (11)	18	
$(\nu_1 + \nu_4^1)^+ - 0^+$	4955.73	100	18 (13)	18	
$(\nu_1 + \nu_4^1)^- - 0^-$	4956.10	130	19 (12)	20	
$(\nu_3^1 + \nu_4^1)^- - 0^-$	5069.59	3	9 (8)	9	
$(\nu_3^1 + \nu_4^1)^+ - 0^+$	5069.88	1	8	9	

15 bands have been observed for the first time in this work, although some of the energy levels involved are known from observations of other bands.

All trivial assignments are secure, as the MARVEL energies (and hence BARVEL line positions) are known to very high accuracy (of the order 10^{-4} cm^{-1} for the energies). The accuracy of branch assignments depends on the determination of the obs. - calc. difference for a given vibrational band.

For bands with many (> 10) assignments the obs. - calc. difference can be tracked through the band. As this remains relatively stable we have confidence in our assignments.

Bands for which only a few lines could be assigned are more tentative,

although every observed band in this work has at least one associated trivial assignment.

It is worth noting that the single lines assigned to $(\nu_2 + 2\nu_4^0)^+ - \nu_2^+$, $(\nu_2 + 2\nu_4^0)^- - \nu_2^-$, $2\nu_1^+ - 2\nu_4^{0,-}$, $(\nu_2 + \nu_3^1)^- - \nu_2^+$ and $(\nu_3^1 + \nu_4^1)^+ - 0^+$ are all trivial.

5. Summary

High-resolution absorption measurements of NH_3 in the region 2100 - 5500 cm^{-1} at atmospheric pressure and a temperature of 1027 °C have been reported and analysed.

A comparison between the measurements and BYTe shows in general good agreement through there are some shifts in line position (up to 2 cm^{-1}) and overall BYTe reproduces experimental intensities only within 30 %. Work towards a new NH_3 line list is currently being carried out as part of the ExoMol project [26].

The use of BYTe and MARVEL has allowed the assignment of 2308 lines. 553 lines were previously assigned by studies included in the HITRAN database. 1755 lines have been assigned for the first time in this work. The 272 lines assigned using MARVEL line positions, also known as trivial assignments, are secure as the accuracy of MARVEL energies is of the order 10^{-4} cm^{-1} . Of the 1483 branch assignments, those associated with bands which have numerous assignments in this work should be reliable because the observed-calculated differences remain relatively stable within a given band. The remaining assignments should also be valid, as all observed bands have at least one verified assignment in this work which provides an expected

observed-calculated difference for the band, however these are more tentative.

Acknowledgements

This work was supported by a grant from Energinet.dk project N. 2013-1-1027, by UCL through the Impact Studentship Program and the European Research Council under Advanced Investigator Project 267219.

References

- [1] D. C. Trimble, AIR QUALITY: Information on Tall Smokestacks and Their Contribution to Interstate Transport of Air Pollution, Tech. rep., GAO U.S. Government Accountability Office, published: May 11. Publicly Released: Jun 10 (2011).
- [2] A. Fateev, S. Clausen, High-resolution spectroscopy of gases at elevated temperatures for industrial applications, in: 22nd UCL Astrophysics Colloquium: Opacities in Cool Stars and Exoplanets, 2012.
- [3] M. R. Schmidt, J. H. He, R. Szczerba, V. Bujarrabal, J. Alcolea, J. Cernicharo, L. Decin, K. Justtanont, D. Teyssier, K. M. Menten, D. A. Neufeld, H. Olofsson, P. Planesas, A. P. Marston, A. M. Sobolev, A. de Koter, F. M. Schöier, Herschel/HIFI observations of the circumstellar ammonia lines in IRC+10216, *Astron. Astrophys.* Accepted. doi:10.1051/0004-6361/201527290.
- [4] D. A. Ladeyschikov, M. S. Kirsanova, A. P. Tsvilev, A. M. Sobolev, Molecular emission in dense massive clumps from the star-forming regions S231–S235, *Astrophysical Bulletin* 71 (2016) 208–224.

- [5] J. Harju, F. Daniel, O. Sipilä, P. Caselli, J. E. Pineda, R. K. Friesen, A. P. R. Güsten, L. Wiesenfeld, P. C. Myers, A. Faure, P. Hily-Blant, C. Rist, E. Rosolowsky, S. Schlemmer, Y. L. Shirley, Deuteration of ammonia in the starless core Ophiuchus/H-MM1, *Astron. Astrophys.* Submitted. (arXiv:1604.05602).
- [6] J. I. Canty, P. W. Lucas, J. Tennyson, S. N. Yurchenko, S. K. Leggett, C. G. Tinney, H. R. A. Jones, B. Burningham, D. J. Pinfield, R. L. Smart, Methane and Ammonia in the near-infrared spectra of late T dwarfs, *Mon. Not. R. Astron. Soc.* 450 (2015) 454–480. doi:10.1093/mnras/stv586.
- [7] S. K. Leggett, C. V. Morley, M. S. Marley, D. Saumon, NEAR-INFRARED PHOTOMETRY OF Y DWARFS: LOW AMMONIA ABUNDANCE AND THE ONSET OF WATER CLOUDS, *Astrophys. J.* 799 (2015) 37. doi:{10.1088/0004-637X/799/1/37}.
- [8] J. H. Woodman, L. Trafton, T. Owen, The abundances of ammonia in the atmospheres of Jupiter, Saturn, AND Titan, *Icarus* 32 (1977) 314–320.
- [9] V. N. Salinas, M. R. Hogerheijde, E. A. Bergin, L. Ilse, C. Brinch, G. A. Blake, D. C. Lis, G. J. Melnick, O. Panic, J. C. Pearson, L. Kristensen, U. A. Yildiz, E. F. van Dishoeck, First detection of gas-phase ammonia in a planet-forming disk, *Astron. Astrophys.* 591 (2016) A122.

- [10] R. J. Hargreaves, G. Li, P. F. Bernath, Ammonia line lists from 1650 to 4000 cm^{-1} , *J. Quant. Spectrosc. Radiat. Transf.* 113 (2012) 670–679.
- [11] R. J. Hargreaves, G. Li, P. F. Bernath, Hot NH_3 spectra for astrophysical applications, *Astrophys. J.* 735 (2012) 111.
- [12] N. F. Zobov, S. V. Shirin, R. I. Ovsyannikov, O. L. Polyansky, S. N. Yurchenko, R. J. Barber, J. Tennyson, R. Hargreaves, P. Bernath, Analysis of high temperature Ammonia spectra from 780 to 2100 cm^{-1} , *J. Mol. Spectrosc.* 269 (2011) 104–108.
- [13] E. J. Barton, S. N. Yurchenko, J. Tennyson, S. Clausen, A. Fateev, High-resolution absorption measurements of NH_3 at high temperatures: 500 - 2100 cm^{-1} , *J. Quant. Spectrosc. Radiat. Transf.* 167 (2015) 126–134. doi:10.1016/j.jqsrt.2015.07.020.
- [14] E. J. Barton, S. N. Yurchenko, J. Tennyson, S. Béguier, A. Campargue, A near infrared line list for NH_3 : Analysis of a Kitt Peak spectrum after 35 years, *J. Mol. Spectrosc.* 325 (2016) 7–12. doi:10.1016/j.jms.2016.05.001.
- [15] P. Cermák, J. Hovorka, P. Veis, P. Cacciani, J. Cosléou, J. E. Romh, M. Khelkhal, Spectroscopy of $^{14}\text{nh}_3$ and $^{15}\text{nh}_3$ in the 2.3 μm spectral range with a new vecsel laser source, *J. Quant. Spectrosc. Radiat. Transfer* 137 (2014) 13–22.
- [16] P. Cacciani, P. Cermák, J. Cosléou, J. El Rohm, J. Hovorka, M. Khelkhal, Spectroscopy of ammonia in the range 6626–6805 cm^{-1} :

- using temperature dependence towards a complete list of lower state energy transitions, *Mol. Phys.* 112 (2014) 2476–2485.
- [17] A. R. Al Derzi, T. Furtenbacher, S. N. Yurchenko, J. Tennyson, A. G. Császár, MARVEL analysis of the measured high-resolution spectra of $^{14}\text{NH}_3$, *J. Quant. Spectrosc. Radiat. Transf.* 161 (2015) 117–130. doi: 10.1016/j.jqsrt.2015.03.034.
- [18] T. Furtenbacher, A. G. Császár, J. Tennyson, MARVEL: measured active rotational-vibrational energy levels, *J. Mol. Spectrosc.* 245 (2007) 115–125.
- [19] T. Furtenbacher, A. G. Császár, MARVEL: measured active rotational-vibrational energy levels. II. Algorithmic improvements, *J. Quant. Spectrosc. Radiat. Transf.* 113 (2012) 929–935.
- [20] K. Sung, S. Yu, J. Pearson, O. Pirali, F. K. Tchana, L. Manceron, Far-infrared $^{14}\text{NH}_3$ line positions and intensities measured with a ft-ir and {AILES} beamline, synchrotron {SOLEIL}, *J. Mol. Spectrosc.* (2016) –<http://dx.doi.org/http://dx.doi.org/10.1016/j.jms.2016.06.011> doi:<http://dx.doi.org/10.1016/j.jms.2016.06.011>.
- [21] S. N. Yurchenko, R. J. Barber, A. Yachmenev, W. Thiel, P. Jensen, J. Tennyson, A variationally computed $T=300$ K line list for NH_3 , *J. Phys. Chem. A* 113 (2009) 11845–11855.
- [22] S. N. Yurchenko, R. J. Barber, J. Tennyson, A variationally computed hot line list for NH_3 , *Mon. Not. R. Astron. Soc.* 413 (2011) 1828–1834.

- [23] X. Huang, D. W. Schwenke, T. J. Lee, Rovibrational spectra of ammonia. II. Detailed analysis, comparison, and prediction of spectroscopic assignments for $^{14}\text{NH}_3$, $^{15}\text{NH}_3$, and $^{14}\text{ND}_3$, *J. Chem. Phys.* 134 (2011) 044321. doi:10.1063/1.3541352.
- [24] S. N. Yurchenko, R. J. Barber, J. Tennyson, W. Thiel, P. Jensen, Towards efficient refinement of molecular potential energy surfaces: Ammonia as a case study, *J. Mol. Spectrosc.* 268 (2011) 123–129.
- [25] S. N. Yurchenko, W. Thiel, P. Jensen, Theoretical ROVibrational Energies (TROVE): A robust numerical approach to the calculation of rovibrational energies for polyatomic molecules, *J. Mol. Spectrosc.* 245 (2007) 126–140. doi:10.1016/j.jms.2007.07.009.
- [26] J. Tennyson, S. N. Yurchenko, ExoMol: molecular line lists for exoplanet and other atmospheres, *Mon. Not. R. Astron. Soc.* 425 (2012) 21–33.
- [27] J. Tennyson, S. N. Yurchenko, A. F. Al-Refaie, E. J. Barton, K. L. Chubb, P. A. Coles, S. Diamantopoulou, M. N. Gorman, C. Hill, A. Z. Lam, L. Lodi, L. K. McKemmish, Y. Na, A. Owens, O. L. Polyansky, T. Rivlin, C. Sousa-Silva, D. S. Underwood, A. Yachmenev, E. Zak, The ExoMol database: molecular line lists for exoplanet and other hot atmospheres, *J. Mol. Spectrosc.* 372 (2016) 73–94. doi:10.1016/j.jms.2016.05.002.
- [28] X. Huang, D. W. Schwenke, T. J. Lee, Rovibrational spectra of ammonia. i. unprecedented accuracy of a potential energy surface used with nonadiabatic corrections, *J. Chem. Phys.* 134 (2011) 044320.

- [29] X. Huang, D. W. Schwenke, T. J. Lee, Rovibrational spectra of ammonia. II. Detailed analysis, comparison, and prediction of spectroscopic assignments for $^{14}\text{NH}_3$, $^{15}\text{NH}_3$, and $^{14}\text{ND}_3$, *J. Chem. Phys.* 134 (2011) 044321.
- [30] K. Sung, L. R. Brown, X. Huang, D. W. Schwenke, T. J. Lee, S. L. Coy, K. K. Lehmann, Extended line positions, intensities, empirical lower state energies and quantum assignments of nh_3 from 6300 to 7000 cm^{-1} , *J. Quant. Spectrosc. Radiat. Transfer* 113 (2012) 1066–1083. doi:10.1016/j.jqsrt.2012.02.037.
- [31] S. Clausen, K. A. Nielsen, A. Fateev, Ceramic gas cell operating up to 1873 K, *Measurement Science and Technology*In preparation.
- [32] V. Evseev, A. Fateev, S. Clausen, High-resolution transmission measurements of CO_2 at high temperatures for industrial applications, *J. Quant. Spectrosc. Radiat. Transf.* 113 (2012) 2222–2233.
- [33] V. Bercher, S. Clausen, A. Fateev, H. Spliethoff, *Oxyfuel combustion, International Greenhouse Gas control* 5 (2011) S76–99.
- [34] M. Alberti, R. Weber, M. Mancini, A. Fateev, S. Clausen, Validation of HITEMP-2010 for carbon dioxide and water vapour at high temperatures and atmospheric pressure in 450-7600 cm^{-1} spectral range, *J. Quant. Spectrosc. Radiat. Transf.* 157 (2015) 14–33.
- [35] A. Fateev, S. Clausen, Online non-contact gas analysis, Tech. rep., Technical University of Denmark, contract no.: Energinet.dk no. 2006 1 6382. (2008).

- [36] C. Hill, S. N. Yurchenko, J. Tennyson, Temperature-dependent molecular absorption cross sections for exoplanets and other atmospheres, *Icarus* 226 (2013) 1673–1677.
- [37] P. R. Griffiths, J. A. de Haseth, Fourier transform infrared spectrometry: chemical analysis, John Wiley and Sons, New York, 1986.
- [38] O. L. Polyansky, N. F. Zobov, S. Viti, J. Tennyson, P. F. Bernath, L. Wallace, K band spectrum of water in sunspots, *Astrophys. J.* 489 (1997) L205–L208.
- [39] L. S. Rothman, I. E. Gordon, Y. Babikov, A. Barbe, D. C. Benner, P. F. Bernath, M. Birk, L. Bizzocchi, V. Boudon, L. R. Brown, A. Campargue, K. Chance, E. A. Cohen, L. H. Coudert, V. M. Devi, B. J. Drouin, A. Fayt, J.-M. Flaud, R. R. Gamache, J. J. Harrison, J.-M. Hartmann, C. Hill, J. T. Hodges, D. Jacquemart, A. Jolly, J. Lamouroux, R. J. Le Roy, G. Li, D. A. Long, O. M. Lyulin, C. J. Mackie, S. T. Massie, S. Mikhailenko, H. S. P. Müller, O. V. Naumenko, A. V. Nikitin, J. Orphal, V. Perevalov, A. Perrin, E. R. Polovtseva, C. Richard, M. A. H. Smith, E. Starikova, K. Sung, S. Tashkun, J. Tennyson, G. C. Toon, V. G. Tyuterev, G. Wagner, The *HITRAN* 2012 molecular spectroscopic database, *J. Quant. Spectrosc. Radiat. Transf.* 130 (2013) 4 – 50. doi:10.1016/j.jqsrt.2013.07.002.
- [40] M. J. Down, C. Hill, S. N. Yurchenko, J. Tennyson, L. R. Brown, I. Kleiner, Re-analysis of ammonia spectra: Updating the HITRAN $^{14}\text{NH}_3$ database, *J. Quant. Spectrosc. Radiat. Transf.* 130 (2013) 260–272.