



Review

Learning from epidemiological, clinical, and immunological studies on *Mycobacterium africanum* for improving current understanding of host–pathogen interactions, and for the development and evaluation of diagnostics, host-directed therapies, and vaccines for tuberculosis



Alimuddin Zumla^{a,b,c}, Isaac Darko Otchere^d, Gloria Ivy Mensah^d, Adwoa Asante-Poku^d, Florian Gehre^{e,f}, Markus Maeurer^g, Matthew Bates^{a,c}, Peter Mwaba^c, Francine Ntoumi^{h,i}, Dorothy Yeboah-Manu^{d,*}

^a Center for Clinical Microbiology, Division of Infection and Immunity, University College London, London, UK

^b National Institute of Health Research, Biomedical Research Centre at UCL Hospitals, London, UK

^c UNZA-UCLMS Research and Training Program, University Teaching Hospital, Lusaka, Zambia

^d Department of Bacteriology, Noguchi Memorial Institute for Medical Research, Accra, Ghana

^e Mycobacterial Unit, Biomedical Sciences, Institute of Tropical Medicine, Antwerp, Belgium

^f Vaccines and Immunity Theme, Medical Research Council (MRC) Unit, Fajara, The Gambia

^g Therapeutic Immunology (TIM) Division, Department of Laboratory Medicine, Karolinska University Hospital Huddinge, and Centre for Allogeneic Stem Cell Transplantation, Karolinska University Hospital Huddinge, Stockholm, Sweden

^h Fondation Congolaise pour la Recherche Médicale, Faculté des Sciences de la Santé, Marien Ngouabi University, Brazzaville, Congo

ⁱ Institute for Tropical Medicine, University of Tübingen, Tübingen, Germany

ARTICLE INFO

Article history:

Received 9 November 2016

Received in revised form 1 December 2016

Accepted 5 December 2016

Corresponding Editor: Eskild Petersen, Aarhus, Denmark

Keywords:

TB

Mycobacterium africanum

Mycobacterium tuberculosis complex

Host

Diagnostics

Vaccines

Africa

EDCTP

SUMMARY

Mycobacterium africanum comprises two phylogenetic lineages within the *Mycobacterium tuberculosis* complex (MTBC). *M. africanum* was first described and isolated in 1968 from the sputum of a Senegalese patient with pulmonary tuberculosis (TB) and it has been identified increasingly as an important cause of human TB, particularly prevalent in West Africa. The restricted geographical distribution of *M. africanum*, in contrast to the widespread global distribution of other species of MTBC, requires explanation. Available data indicate that *M. africanum* may also have important differences in transmission, pathogenesis, and host–pathogen interactions, which could affect the evaluation of new TB intervention tools (diagnostics and vaccines)—those currently in use and those under development. The unequal geographical distribution and spread of MTBC species means that individual research findings from one country or region cannot be generalized across the continent. Thus, generalizing data from previous and ongoing research studies on MTBC may be inaccurate and inappropriate. A major rethink is required regarding the design and structure of future clinical trials of new interventions. The West, Central, East, and Southern African EDCTP Networks of Excellence provide opportunities to take forward these pan-Africa studies. More investments into molecular, epidemiological, clinical, diagnostic, and immunological studies across the African continent are required to enable further understanding of host–*M. africanum* interactions, leading to the development of more specific diagnostics, biomarkers, host-directed therapies, and vaccines for TB.

© 2016 The Author(s). Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Tuberculosis (TB) in humans is caused by one of several genetically related groups of organisms comprising the

Mycobacterium tuberculosis complex (MTBC). Genomic studies indicate that all MTBC species have between 95% and 100% DNA relatedness, with the same 16S rRNA.^{1,2} *Mycobacterium tuberculosis* sensu stricto is responsible for a large proportion of human TB cases worldwide.³ Other MTBC species that cause human disease include *Mycobacterium bovis*, *Mycobacterium africanum*, *Mycobacterium microti*, *Mycobacterium canettii*, and *Mycobacterium pinnipedii*.^{4–6} Globally, TB continues to be a major public health

* Corresponding author.

E-mail address: dyeboah-manu@noguchi.ug.edu.gh (D. Yeboah-Manu).

problem and is now the most common infectious disease cause of death. The World Health Organization annual TB report for 2016 states that there were 10 million cases of active TB in 2015, with an estimated 1.5 million associated deaths.⁷

Over the past six decades, the common dogma prevailing has been that all members of the MTBC are genetically similar, with a clonal population. Thus, it has generally been assumed that the small genotypic or phenotypic differences do not translate into differences in transmission, pathogenesis, host–pathogen interaction, disease outcome, and efficacy of diagnostics and vaccines. A huge amount of resources and effort have been invested in developing new TB diagnostics, TB drugs, treatment regimens, and vaccines. Currently available diagnostics and vaccines, and those under development, have been based on this assumption. However, molecular and clinical studies combined with genomic analyses have shown important strain diversity among the different members of the MTBC.^{4,8–12} Differences between lineages could affect the performance of diagnostics, treatments, and vaccines.

Available data obtained from epidemiological, clinical, and immunological studies on *M. africanum* in West Africa are highlighted in this viewpoint;^{10,13} these indicate important differences in transmission, pathogenesis, and host–pathogen interactions. The need for a radical rethink to improve current understanding of host–pathogen interactions and for the development and evaluation of new diagnostics, biomarkers, host-directed therapies, and specific vaccines for TB is also discussed.

2. Geographical distribution of *M. africanum*

M. africanum was first described and isolated in 1968 from the sputum of a Senegalese patient with pulmonary TB.⁵ *M. africanum* is made up of two lineages within the MTBC: MTBC lineage 5 (also called *M. africanum* West African type I) and MTBC lineage 6 (*M. africanum* West African type II).^{13,14} With advances in diagnostic technologies, *M. africanum* has been identified increasingly as an important cause of human TB, causing nearly 50% of all TB cases reported in West Africa. The following country prevalence of *M. africanum* has been reported: Benin 39%, Burkina Faso 18%, Cameroon 56%, The Gambia 39%, Ghana 20%, Guinea Bissau 47%, Ivory Coast 55%, Nigeria 8%, Senegal 20%, Sierra Leone 24%.^{13,15–20} *M. africanum* has also been reported infrequently among TB patients from across Europe, South America, and the USA.^{21–26} It is assumed that TB caused by *M. africanum* in non-African countries is due to human migration from disease-endemic West African regions.

Recent advances in molecular technology have allowed studies on the evolution of the nature of *M. africanum* genotypes, its position within the MTBC, and its genotypic and phenotypic expression, specific clade identification, transmission patterns, infectivity, and disease expression in humans.^{10–33} Whether there are significant differences between *M. tuberculosis* and *M. africanum* in their clinical presentation, epidemiology, association with migration, transmission patterns, pathogenesis, virulence, rate of progression of disease, and treatment outcomes needs to be defined.

3. Clinical manifestations, transmission, and genotype studies of TB due to *M. africanum*

Several small clinical cohort studies from West Africa have evaluated differences in clinical presentation between TB caused by *M. africanum* and TB caused by *M. tuberculosis* sensu stricto.^{10–12,27–33} Aerosol spread and symptoms at presentation seem similar in both. No significant associations of *M. africanum* with HIV status and chest X-ray changes have been noted. Contacts of patients with

M. africanum infection appear to have a lower rate of progression to active TB compared with contacts of *M. tuberculosis*-infected patients.³⁴ A lower rate of genotype clustering for *M. africanum* compared to *M. tuberculosis* has been reported.³⁵ Other studies have reported that *M. africanum* West African type II has a similar rate of transmission as *M. tuberculosis* sensu stricto, although it may be associated with a lower rate of progression to disease in contacts.^{36,37}

4. *M. africanum* genotype, immunological responses, and diagnostic tests

Immunological studies reported from The Gambia have shown marked differences in clinical features, immune response, gene expression, and proteins associated with inflammation between *M. tuberculosis* sensu stricto-infected patients and *M. africanum*-infected patients.^{35,38,39} A number of genomic polymorphisms unique to *M. africanum* in West Africa and a study using whole genome sequencing of four *M. africanum* isolates against H37Rv,¹¹ have shown a number of frame-shifts and non-synonymous mutations within some functional operons, including genes associated with immunogenicity and host invasion. It was highlighted that potential differential gene expression could have implications for diagnostic tests and management algorithms in West Africa, especially when these genes are being targeted for diagnostic test purposes.

The gene product of *mpt64* is the target of three widely used rapid speciation lateral flow assays for the identification of the MTBC in culture: BD MGIT TBc Identification Test (BD TBc ID; Becton Dickinson Diagnostics, Becton, Dickinson and Company, Sparks, MD, USA); SD Bioline Ag MPT64 Rapid (SD Bioline; Standard Diagnostics, Inc., Yongin-si, Gyeonggi-do, Republic of Korea); and Capilia TB-Neo (TAUNS Laboratories, Inc., Numazu, Shizuoka, Japan). Despite the advantage of lateral flow assays, there have been reports of the failure of MPT64 tests to detect MTBC isolates, resulting in erroneous reporting of a non-tuberculous mycobacteria (NTM) diagnosis. A systematic evaluation of two commonly used rapid TB tests that detect the product of the mycobacterial *mpt64* gene compared the abundance of the *mpt64* gene product in sputum samples of patients with untreated pulmonary TB caused by *M. africanum* type II or *M. tuberculosis* sensu stricto. Ofori-Anyinam et al. compared the expression of the *mpt64* gene in sputa from *M. africanum* and *M. tuberculosis* sensu stricto-infected Gambian patients and found that the *mpt64* gene was 2.5-fold less expressed in *M. africanum* than in *M. tuberculosis* sensu stricto.³³ Thus the sensitivity of rapid tests was 20% lower for *M. africanum* cultures than for *M. tuberculosis* sensu stricto cultures.³³ A non-synonymous single nucleotide polymorphism (SNP) in the *mpt64* gene of that lineage was identified. The *mpt64* (Rv1980c) mRNA transcript was significantly less abundant in the sputa of TB patients infected with *M. africanum* type II compared to patients infected with *M. tuberculosis* sensu stricto, suggesting that the MPT64 protein may be produced at a slower rate. There also appeared to be lineage-dependent and time-specific differences in conversion to MPT64 test positivity. Thus, in West Africa, tests to identify MTBC in culture miss a substantial fraction of TB cases, and this may have direct consequences for the patients and for TB control efforts. These data have important implications for the performance of diagnostic tests and any new TB vaccine in West Africa.

5. Bugbear of current portfolio of geographically restricted TB research

Recent debates have focused on whether the currently available diagnostic tests are specific and reliable for all types of MTBC

lineage, or whether treatment or vaccine success or failure will be determined by the specific nature of MTBC lineages. Given the limited efficacy of the current TB vaccine and the recent clinical failure of the most advanced new TB vaccine candidate, novel concepts for vaccine design should be explored. Over the past 5 years, advances in molecular technologies have shown that MTBC lineages differ in their phenotypes and that they may also differ in their pathogenicity. Thus any generalizing of data from previous and ongoing research studies on TB may be inappropriate and inaccurate. The unequal geographical distribution and spread of MTBC species across Africa means that individual research findings from one country or region cannot be generalized across the continent.¹³ Thus, a major rethink is required for the design and structure of future clinical trials of new diagnostic and preventive interventions. Phylogeographic studies across the African continent are now required to improve the design and interpretation of epidemiological research, translational clinical research, and evaluation trials of new interventions (rapid diagnostics, new drugs and treatment regimens, and vaccines).

6. Conclusions and the way forward

Recent phylogenomic analysis of global strains has indicated that while some lineages and sub-lineages casually referred to as generalist can infect different populations, the majority of tuberculous lineages and sub-lineages are 'specialized' to cause TB disease in distinct human populations. *M. africanum* classified into L5 and L6 seems to have specialized and remains an important cause of TB in West Africa. With the wide spectrum of MTBC lineages present across Africa, these patient cohorts provide ideal trial sites for evaluation studies. Increased investment in the study of *M. africanum* is required to further our understanding of MTBC epidemiology, evolution, and pathogenesis. The diversity within *M. africanum* must be factored into priority agendas of funders and current investments into new TB biomarkers and vaccine development. With the limited efficacy of bacille Calmette–Guérin (BCG) and the failure of the Oxford TB vaccine, novel concepts for vaccine design should be explored, such as the highly variable T-cell epitopes. Moreover, there is increasing evidence that the human-adapted MTBC has been co-evolving with the human host for a long time. Hence, studying the interaction between bacterial and human genetic diversity might help identify additional targets that could be exploited for TB vaccine development. It is important that systematic collaborative research studies are also conducted across Africa to determine the contribution of host genetics, ethnicity, or other factors in the persistent circulation of *M. africanum* in West Africa. Moreover, such studies also have the potential to help in developing host-specific adjunct host-directed therapies against both *M. africanum* and *M. tuberculosis* sensu stricto.⁴⁰ Whilst the infrastructure and resources for the performance of these pan-African studies have been lacking, the second European Developing Countries Clinical Trials Partnership program (EDCTP2)⁴¹ and the EDCTP regional networks of excellence^{42,43} now provide new opportunities to take forward these important studies.

Acknowledgements

DYM acknowledges support from the Wellcome Trust, UK. Authors AZ, MB, and FN acknowledge support from the European and Developing Countries Clinical Trials Partnership (EDCTP), The Hague, The Netherlands. AZ receives support from the NIHR Biomedical Research Centre at UCL Hospitals, London, UK.

Conflict of interest: The authors declare no conflicts of interest.

References

- Musser JM, Amin A, Ramaswamy S. Negligible genetic diversity of *Mycobacterium tuberculosis* host immune system protein targets: evidence of limited selective pressure. *Genetics* 2000;**155**:7–16.
- Sreevatsan S, Pan X, Stockbauer KE, Connell ND, Kreiswirth BN, Whittam TS, Musser JM. Restricted structural gene polymorphism in the *Mycobacterium tuberculosis* complex indicates evolutionarily recent global dissemination. *Proc Natl Acad Sci U S A* 1997;**94**:9869–74.
- Gagneux S, DeRiemer K, Van T, Kato-Maeda M, de Jong BC, Narayanan S, et al. Variable host–pathogen compatibility in *Mycobacterium tuberculosis*. *Proc Natl Acad Sci U S A* 2006;**103**:2869–73. <http://dx.doi.org/10.1073/pnas.0511240103>
- Frota CC, Hunt DM, Buxton RS, Rickman L, Hinds J, Kremer K, et al. Genome structure in the vole bacillus, *Mycobacterium microti*, a member of the *Mycobacterium tuberculosis* complex with a low virulence for humans. *Microbiology* 2004;**150**:1519–27. <http://dx.doi.org/10.1099/mic.0.26660-0>
- Castets M, Boisvert H, Grumbach F, Brunel M, Rist N. Tuberculosis bacilli of the African type: preliminary note. *Rev Tuberc Pneumol (Paris)* 1968;**32**:179–84.
- Brosch R, Gordon SV, Marmiesse M, Brodin P, Buchrieser C, Eiglmeier K, et al. A new evolutionary scenario for the *Mycobacterium tuberculosis* complex. *Proc Natl Acad Sci U S A* 2002;**99**:3684–9. <http://dx.doi.org/10.1073/pnas.052548299>
- World Health Organization. Global tuberculosis report 2016. Geneva: WHO; 2016.
- Cole ST, Brosch R, Parkhill J, Garnier T, Churcher C, Harris D, et al. Deciphering the biology of *Mycobacterium tuberculosis* from the complete genome sequence. *Nature* 1998;**393**:537–44. <http://dx.doi.org/10.1038/31159>
- Comas I, Homolka S, Niemann S, Gagneux S. Genotyping of genetically monomorphic bacteria: DNA sequencing in *Mycobacterium tuberculosis* highlights the limitations of current methodologies. *PLoS One* 2009;**4**:e7815. <http://dx.doi.org/10.1371/journal.pone.0007815>
- Comas I, Coscolla M, Luo T, Borrell S, Holt KE, Kato-Maeda M, et al. Out-of-Africa migration and Neolithic coexpansion of *Mycobacterium tuberculosis* with modern humans. *Nat Genet* 2013;**45**:1176–82. <http://dx.doi.org/10.1038/ng.2744>
- Gehre F, Otu J, DeRiemer K, de Sessions PF, Hibberd ML, Mulders W, et al. Deciphering the growth behaviour of *Mycobacterium africanum*. *PLoS Negl Trop Dis* 2013;**7**:e2220. <http://dx.doi.org/10.1371/journal.pntd.0002220>
- Coscolla M, Gagneux S. Consequences of genomic diversity in *Mycobacterium tuberculosis*. *Semin Immunol* 2014;**26**:431–44. <http://dx.doi.org/10.1016/j.smim.2014.09.012>
- De Jong BC, Antonio M, Gagneux S. *Mycobacterium africanum*—review of an important cause of human tuberculosis in West Africa. *PLoS Neglected Tropical Diseases* 2010;**4**:e744. <http://dx.doi.org/10.1371/journal.pntd.0000744>
- Niemann S, Kubica T, Bange FC, Adjei O, Browne EN, Chinbuah MA, et al. The species *Mycobacterium africanum* in the light of new molecular markers. *J Clin Microbiol* 2004;**42**:3958–62. <http://dx.doi.org/10.1128/JCM.42.9.3958-3962.2004>
- Affolabi D, Faihun F, Sanoussi N, Anyo G, Shamputa IC, Rigouts L, et al. Possible outbreak of streptomycin-resistant *Mycobacterium tuberculosis* Beijing in Benin. *Emerg Infect Dis* 2009;**15**:1123–5. <http://dx.doi.org/10.3201/eid1507.080697>
- Groenheit R, Ghebremichael S, Svensson J, Rabna P, Colombatti R, Riccardi F, et al. The Guinea-Bissau family of *Mycobacterium tuberculosis* complex revisited. *PLoS One* 2011;**6**:e18601. <http://dx.doi.org/10.1371/journal.pone.0018601>
- Källénus G, Koivula T, Ghebremichael S, Hoffner SE, Norberg R, Svensson E, et al. Evolution and clonal traits of *Mycobacterium tuberculosis* complex in Guinea-Bissau. *J Clin Microbiol* 1999;**37**:3872–8.
- Frothingham R, Strickland PL, Bretzel G, Ramaswamy S, Musser JM, Diana L, Williams DL. Phenotypic and genotypic characterization of *Mycobacterium africanum* isolates from West Africa. *Journal of Clinical Microbiology* 1999;**37**:1921–6.
- Viana-Niero C, Gutierrez C, Sola C, Filioli I, Boulahbal F, Vincent V, Rastogi N. Genetic diversity of *Mycobacterium africanum* clinical isolates based on IS6110-restriction fragment length polymorphism analysis, spoligotyping, and variable number of tandem DNA repeats. *J Clin Microbiol* 2001;**39**:57–65. <http://dx.doi.org/10.1128/JCM.39.1.57-65.2001>
- Yeboah-Manu D, Asare P, Asante-Poku A, Otchere ID, Osei-Wusu S, Danso E, et al. Spatio-temporal distribution of *Mycobacterium tuberculosis* complex strains in Ghana. *PLoS One* 2016;**11**:e0161892. <http://dx.doi.org/10.1371/journal.pone.0007815>
- Jungbluth H, Fink H, Reusch F. Tuberculous infection caused by *Myco. africanum* in black Africans resident in the German Federal Republic (author's transl). *Prax Klin Pneumol* 1978;**32**:306–9.
- Schröder KH. Isolation of *M. africanum* from German patients. *Bull Int Union Tuberc* 1982;**57**:249–53.
- Brown T, Nikolayevskiy V, Velji P, Drobniowski F. Associations between *Mycobacterium tuberculosis* strains and phenotypes. *Emerging Infectious Diseases* 2010;**16**. <http://dx.doi.org/10.3201/eid1602.091032>
- Frottier J, Eliasiewicz M, Arlet V, Gaudillat C. Infections caused by *Mycobacterium africanum*. *Bull Acad Natl Med* 1990;**174**:29–33. discussion 34–5.
- Pérez-de Pedro I, Bermúdez P, Artero I, Jiménez MS. Orchiepididymitis due to *Mycobacterium africanum*. *Enferm Infecc Microbiol Clin* 2008;**26**:600–2.
- Desmond E, Ahmed AT, Probert WS, Ely J, Jang Y, Sanders CA, et al. *Mycobacterium africanum* cases, California. *Emerg Infect Dis* 2004;**10**:921–3. <http://dx.doi.org/10.3201/eid1005.030016>
- Gehre F, Antonio M, Otu JK, Sallah N, Secka O, Faal T, et al. Immunogenic *Mycobacterium africanum* strains associated with ongoing transmission in The Gambia. *Emerg Infect Dis* 2013;**19**:1598–604. <http://dx.doi.org/10.3201/eid1910.121023>

28. Leone S, Odoun M, Uwizye C, De Rijk P, Gehre F, Antonio M, et al. The first phylogeographic population structure and analysis of transmission dynamics of *M. africanum* West African 1—combining molecular data from Benin, Nigeria and Sierra Leone. *PLoS One* 2013;**8**:e77000. <http://dx.doi.org/10.1371/journal.pone.0077000>
29. Winglee K, Manson McGuire A, Maiga M, Abeel T, Shea T, Desjardins CA, et al. Whole genome sequencing of *Mycobacterium africanum* strains from Mali provides insights into the mechanisms of geographic restriction. *PLoS Negl Trop Dis* 2016;**10**:e0004332. <http://dx.doi.org/10.1371/journal.pntd.0004332>
30. Asante-Poku A, Yeboah-Manu D, Otchere ID, Aboagye SY, Stucki D, Hattendorf J, et al. *Mycobacterium africanum* is associated with patient ethnicity in Ghana. *PLoS Negl Trop Dis* 2015;**9**:e3370. <http://dx.doi.org/10.1371/journal.pntd.0003370>
31. Bryant JM, Harris SR, Parkhill J, Dawson R, Diacon AH, van Helden P, et al. Whole-genome sequencing to establish relapse or re-infection with *Mycobacterium tuberculosis*: a retrospective observational study. *Lancet Respir Med* 2013;**1**:786–92. [http://dx.doi.org/10.1016/S2213-2600\(13\)70231-5](http://dx.doi.org/10.1016/S2213-2600(13)70231-5)
32. Bentley SD, Comas I, Bryant JM, Walker D, Smith NH, Harris SR, et al. The genome of *Mycobacterium africanum* West African 2 reveals a lineage-specific locus and genome erosion common to the *M. tuberculosis* complex. *PLoS Negl Trop Dis* 2012;**6**:e1552. <http://dx.doi.org/10.1371/journal.pntd.0001552>
33. Ofori-Anyinam B, Kanuteh F, Agbla SC, Adetifa I, Okoi C, Dolganov G, et al. Impact of the *Mycobacterium africanum* West Africa 2 lineage on TB diagnostics in West Africa: decreased sensitivity of rapid identification tests in The Gambia. *PLoS Negl Trop Dis* 2016;**10**:e0004801. <http://dx.doi.org/10.1371/journal.pntd.0004801>
34. de Jong BC, Hill PC, Brookes RH, Gagneux S, Jeffries DJ, Otu JK, et al. *Mycobacterium africanum* elicits an attenuated T cell response to early secreted antigenic target, 6 kDa, in patients with tuberculosis and their household contacts. *J Infect Dis* 2006;**193**:1279–86. <http://dx.doi.org/10.1086/502977>
35. Gehre F, Kumar S, Kendall L, Ejo M, Secka O, Ofori-Anyinam B, et al. A mycobacterial perspective on tuberculosis in West Africa: significant geographical variation of *M. africanum* and other *M. tuberculosis* complex lineages. *PLoS Negl Trop Dis* 2016;**10**:1–11. <http://dx.doi.org/10.1371/journal.pntd.0004408>
36. Pai M, Riley LW, Colford JM. Interferon-gamma assays in the immunodiagnosis of tuberculosis: a systematic review. *Lancet Infect Dis* 2004;**4**:761–76. [http://dx.doi.org/10.1016/S1473-3099\(04\)01206-X](http://dx.doi.org/10.1016/S1473-3099(04)01206-X)
37. de Jong BC, Hill PC, Aiken A, Awine T, Antonio M, Adetifa IM, et al. Progression to active tuberculosis, but not transmission, varies by *Mycobacterium tuberculosis* lineage in The Gambia. *J Infect Dis* 2008;**198**:1037–43. <http://dx.doi.org/10.1086/591504>
38. Tientcheu LD, Haks MC, Agbla SC, Sutherland JS, Adetifa IM, Donkor S, et al. Host immune responses differ between *M. africanum*- and *M. tuberculosis*-infected patients following standard anti-tuberculosis treatment. *PLOS Neglected Tropical Diseases* 2016;**10**:e0004701. <http://dx.doi.org/10.1371/journal.pntd.0004701>
39. Tientcheu LD, Maertzdorf J, Weiner J, Adetifa IM, Mollenkopf HJ, Sutherland JS, et al. Differential transcriptomic and metabolic profiles of *M. africanum*- and *M. tuberculosis*-infected patients after, but not before, drug treatment. *Genes Immun* 2015;**16**:347–55. <http://dx.doi.org/10.1038/gene.2015.21>
40. Zumla A, Maeurer M, Host-Directed Therapies Network. Towards host-directed therapies for tuberculosis. *Nat Rev Drug Discov* 2015;**14**:511–2. <http://dx.doi.org/10.1038/nrd4696>
41. Zumla A, Makanga M, Nyirenda T, Beattie P, Olesen OF, Breugelmans JG, et al. Genesis of EDCTP2. *Lancet Infect Dis* 2015;**15**:11–3. [http://dx.doi.org/10.1016/S1473-3099\(14\)71034-5](http://dx.doi.org/10.1016/S1473-3099(14)71034-5)
42. Ntoumi F, Kaleebu P, Macete E, Mfinanga S, Chakaya J, Yeboah-Manu D, et al. Taking forward the World TB Day 2016 theme 'Unite to End Tuberculosis' for the WHO Africa Region. *Int J Infect Dis* 2016;**46**:34–7. <http://dx.doi.org/10.1016/j.ijid.2016.03.003>
43. Zumla A, Petersen E, Nyirenda T, Chakaya J. Tackling the tuberculosis epidemic in Sub-Saharan Africa—unique opportunities arising from the second European Developing Countries Clinical Trials Partnership (EDCTP) programme 2015–2024. *Int J Infect Dis* 2015;**32**:46–9.