

Discovery of Multi-Target Agents Active as Broad-Spectrum Antivirals and Correctors of Cystic Fibrosis Transmembrane Conductance Regulator (CFTR) for Associated Pulmonary Diseases

Sabrina Tassini, Liang Sun, Kristina Lanko, Emmanuele Crespan, Emily Langron, Federico Falchi, Miroslava Kissova, Jorge I. Armijos-Rivera, Leen Delang, Carmen Mirabelli, Johan Neyts, Marco Pieroni, Andrea Cavalli, Gabriele Costantino, Giovanni Maga, Paola Vergani, Pieter LEYSEN, and Marco Radi

J. Med. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.jmedchem.6b01521 • Publication Date (Web): 25 Jan 2017

Downloaded from <http://pubs.acs.org> on January 26, 2017

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discovery of Multi-Target Agents Active as Broad-Spectrum Antivirals and Correctors of Cystic Fibrosis Transmembrane Conductance Regulator (CFTR) for Associated Pulmonary Diseases

Sabrina Tassini,^a Liang Sun,^b Kristina Lanko,^b Emmanuele Crespan,^c Emily Langron,^d
Federico Falchi,^{e,f} Miroslava Kissova,^c Jorge I. Armijos-Rivera,^c Leen Delang,^b Carmen
Mirabelli,^b Johan Neyts,^{b,*} Marco Pieroni,^a Andrea Cavalli,^{e,f} Gabriele Costantino,^a Giovanni
Maga,^c Paola Vergani,^d Pieter Leyssen,^b and Marco Radi^{a,*}

^aP4T Group, Dipartimento di Scienze degli Alimenti e del Farmaco, Università degli Studi di
Parma, Viale delle Scienze, 27/A, 43124 Parma, Italy;

^bLaboratory of Virology and Experimental Chemotherapy, Rega Institute for Medical
Research, KU Leuven, Minderbroedersstraat 10, 3000, Leuven, Belgium;

^cIstituto di Genetica Molecolare, IGM-CNR, Via Abbiategrasso 207, 27100 Pavia, Italy;

^dDepartment of Neuroscience, Physiology and Pharmacology, University College London,
Gower Street, WC1E 6BT London, UK;

^eCompuNet, Istituto Italiano di Tecnologia, Via Morego 30, I-16163 Genova, Italy;

1
2
3 †Department of Pharmacy and Biotechnology, University of Bologna, Via Belmeloro 6, I-
4
5 40126 Bologna, Italy.
6

7
8 KEYWORDS: PI4KIII β , F508del-CFTR, broad-spectrum antivirals, enterovirus, cystic
9
10 fibrosis, multi-target.

11 12 13 **ABSTRACT**

14
15
16
17 Enteroviruses (EVs) are among the most frequent infectious agents in humans worldwide
18 and represent the leading cause of upper respiratory tract infections. No drugs for the
19 treatment of EV infections are currently available. Recent studies have also linked
20 treatment of EV infections are currently available. Recent studies have also linked
21 enterovirus infection with pulmonary exacerbations, especially in cystic fibrosis (CF)
22 patients, and the importance of this link is probably underestimated. The aim of this work
23 was to develop a new class of multi-target agents active both as broad-spectrum antivirals
24 and as correctors of the F508del-CFTR folding defect responsible for >90% of CF cases. We
25 report herein the discovery of the first small-molecules able to simultaneously act as
26 correctors of the F508del-CFTR folding defect and as broad-spectrum antivirals against a
27 panel of enteroviruses representative of all major species.
28
29
30
31
32
33
34
35
36
37
38
39
40

41 42 **INTRODUCTION**

43
44 Enteroviruses (EVs) are positive-sense single stranded RNA viruses, classified into 12
45 species, including four human enterovirus species (EV-A to EV-D), three species of human
46 rhinoviruses (RV-A to RV-C) and five enterovirus species that only infect animals.¹ EVs are
47 responsible for a great variety of clinical manifestations, especially in young children, which
48 may result in life-threatening neurological complications (e.g. encephalitis, meningitis and
49 poliomyelitis-like paralysis).²⁻⁴ Furthermore, RV infections are now considered one of the
50 major causes of acute exacerbations in chronic pulmonary diseases like asthma, chronic
51
52
53
54
55
56
57
58
59
60

1
2
3 obstructive pulmonary disorder (COPD) and cystic fibrosis (CF) in children and adults.⁵
4
5 Physicians pay particular attention to patients that already suffer from respiratory diseases,
6
7 such as CF or asthma, as they could be particularly affected by an additional enterovirus
8
9 infection.⁶ An increasing number of studies also suggest that respiratory viruses, in particular
10
11 enterovirus and rhinovirus, contribute significantly to CF pulmonary exacerbations,
12
13 hospitalization, decreased lung function and predisposition to bacterial colonization.⁷ The
14
15 mechanistic link between viral infections and deterioration of CF lung function is not fully
16
17 understood and their impact is probably underestimated, especially in young children.⁸
18
19 Despite their high clinical and socioeconomic impact, to date there is no approved antiviral
20
21 therapy for the prophylaxis and/or the treatment of enterovirus infections, and the
22
23 management of patients is currently limited to symptomatic treatment and supportive care.
24
25 Therefore, there is an unmet need for broad-spectrum antiviral drugs as a rapid defense
26
27 strategy against enterovirus infections and virus-related exacerbations.
28
29

30
31
32 Nowadays, host factors are considered as very attractive targets for the development of
33
34 antiviral drugs because they are unlikely to mutate and develop resistance in response to
35
36 therapy.⁹ Moreover, since viruses belonging to the same genus or family usually share the
37
38 same cellular pathways for replication, targeting a host factor may allow the development of
39
40 effective broad-spectrum antiviral compounds.¹⁰ Although some toxicity risks may be
41
42 expected from inhibiting a host factor, it should be kept in mind that most drugs currently
43
44 used in therapy target host proteins with excellent therapeutic outcomes and acceptable safety
45
46 profiles. In particular, it has been well documented that the host lipid kinase
47
48 phosphatidylinositol 4-kinase III β (PI4KIII β) is critical for RNA replication of several
49
50 enteroviruses.¹¹⁻¹⁴ PI4KIII β belongs to the phosphatidylinositol 4-kinases (PI4Ks) that
51
52 synthesize phosphatidylinositol 4-phosphate (PI4P) from phosphatidylinositol (PI). PI4P is
53
54 involved in signaling and cellular trafficking mainly at the Golgi and trans-Golgi network
55
56
57
58
59
60

(TGN), it contributes to defining the characteristics of plasma membranes and it activates a variety of ion channels, including CFTR.¹⁵⁻¹⁸ Four PI4K isoforms have been identified in mammals, classified as type II (PI4KII α and PI4KII β) or type III (PI4KIII α and PI4KIII β) based on their primary sequences and catalytic properties.¹⁹ Type III PI4Ks are hijacked by several ss(+)RNA viruses (especially from *Flaviviridae*, *Picornaviridae* and *Coronaviridae* families) to remodel cellular membranes and generate PI4P lipid-enriched organelles specialized for viral replication.²⁰

A few PI4KIII β inhibitors with antiviral activity against a panel of picornaviruses have been reported recently (Figure 1).²¹⁻²³ Generally, chemical inhibition of PI4KIII β does not influence cell viability.²⁴ One possible explanation might be that while the small amounts of PI4P produced by other PI4K isoforms could be enough to support cell trafficking and signaling, it would not be sufficient to sustain viral RNA synthesis.²⁰ A major aim in the development of PI4KIII β inhibitors is to achieve selective inhibition of the α or β isoforms. Among known PI4KIII β inhibitors, compound **1** (PIK93) is about 100-fold more potent against the PI4KIII β isoform, although it also has detectable activity towards PI3-kinases.^{25,26}




Figure 1. Representative PI4KIII β inhibitors.

1
2
3 Considering the growing need for novel broad-spectrum antivirals to fight emerging
4 epidemics and the link between respiratory viruses and pulmonary exacerbation in cystic
5 fibrosis patients, our aim was the development of a new class of multi-target agents active
6 both as broad-spectrum antivirals (by targeting PI4KIII β) and as correctors of the F508del-
7 CFTR folding defect responsible for >90% of CF cases. We here report the discovery of the
8 first small-molecule compounds able to simultaneously act as moderately efficacious
9 correctors of the F508del-CFTR folding defect and broad-spectrum antivirals against a panel
10 of enteroviruses (linked to CF pulmonary exacerbations).
11
12
13
14
15
16
17
18
19

20 21 22 23 **RESULTS AND DISCUSSION**

24
25 Drug repurposing and polypharmacology are two very attractive approaches in modern
26 drug discovery. The first offers the possibility of recycling known drugs or advanced drug
27 candidates developed for a different disease. The second results in simultaneous action on
28 different targets/diseases with a single, rationally designed drug.^{27,28} In particular,
29 polypharmacology aims at producing multi-target agents whose interference with multiple
30 biochemical pathways offers an advantage - in terms of drug load, efficacy and safety - over
31 combination therapy. This approach is well suited to complex diseases that generally require
32 the simultaneous administration of many different drugs. Considering the increasing number
33 of reports on the connection between enterovirus infections and pulmonary exacerbations in
34 CF patients, we reasoned that an ideal drug candidate for such closely related diseases might
35 be a multi-target agent able to act, at the same time, on proteins/pathways implicated in
36 enterovirus replication (PI4KIII β) and on F508del-CFTR biogenesis. At the beginning of this
37 work, the X-ray structures of the above targets were not available for a structure-based study.
38 We therefore developed a PI4KIII β homology model to be used for the design of PI4KIII β
39 inhibitors, selecting those whose chemical scaffolds resemble known CFTR
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 correctors/potentiators. The structure of the complex of PI3K γ with compound **1** (PDB ID:
4 2CHZ)²⁶ has been used to build the homology model of PI4KIII β by using Prime software
5
6 (see methods): this structure shows an identity of 30%, a positive of 52% and a score of 322.
7
8 The presence of **1** in the structure of PI3K γ allowed us to identify its likely binding site in
9
10 PI4KIII β and hypothesize its binding mode. A 10 ns molecular dynamics simulation on the
11
12 modeled PI4KIII β protein containing compound **1** was performed using the software
13
14 Desmond.²⁹ In the latter (equilibrated) part of the trajectory (last 2 ns) 100 frames were
15
16 extracted and clusterized on the basis of RMSD. Five clusters were obtained. All PI4K β
17
18 inhibitors available in Pubchem³⁰ were docked in the compound **1** binding site of each cluster
19
20 and the frame with the best correlation between docking score and enzymatic activity was
21
22 selected for virtual screening. A high-throughput docking (HTD) approach was then applied
23
24 to the compound **1** binding site in our PI4KIII β model to identify high affinity hits within the
25
26 Asinex database collection.³¹ Compound selection was based on the ranking score and visual
27
28 inspection of the PI4KIII β catalytic site, but also took into account the 2D similarity to
29
30 known CFTR correctors/potentiators. Thirteen commercially available compounds (**5-17**,
31
32 Figure 2), four of which (**6, 11, 16, 17**) resemble known CFTR correctors,^{32,33} were selected
33
34 for biological investigation. These computational results were confirmed on the recently
35
36 released crystal structure of PI4KIII β co-crystallized with compound **1**, and this structure
37
38 (PDB ID: 4D0L) was used for all the following simulations.³⁴
39
40
41
42
43
44

45 These commercially available compounds were then tested both against the PI4KIII β
46
47 enzyme and in a virus-cell-based replication assay. In particular these compounds were
48
49 evaluated for antiviral activity against a panel of enteroviruses that are representative of all
50
51 major species: enterovirus group A (EV71), group B (coxsackievirus B3, CVB3, and
52
53 echovirus 11, ECHO11), group C (poliovirus 1, PV1), group D (enterovirus 68, EV68),
54
55 rhinovirus group A (RV02) and rhinovirus group B (RV14). Among the selected compounds,
56
57
58
59
60

only the bithiazole **17** showed activity in cell-free and cell-based assays and possesses a chemical scaffold (the bithiazole) of a known family of CFTR correctors (Figure 2).^{33,35}

Compound **17** was therefore selected as a promising starting point for further structure-based optimization.





Figure 2. Chemical structure of compounds selected by virtual screening and activity profile of the hit compound **17**.

Chemistry

Compound **17** was initially resynthesized to validate the biological activity of the commercial sample and to set up a synthetic protocol for its chemical diversification starting from cheap and commercially available building blocks (Scheme 1).




Scheme 1. Reagents and conditions: (a) EtOH, reflux, 12 h, 95%; (b) 48% aqueous HBr, Br₂, 1,4-dioxane, 60 °C, 3 h, 90%; (c) (i) benzoyl isothiocyanate, DCM, rt, 12 h, (ii) NaOH 1N, THF, reflux, 3 h, 72%; (d) EtOH, reflux, 1 h, 84%; (e) acetyl chloride, Et₃N, DCM, rt, 15 h, 77%.

Thiourea **18** was condensed with 3-chloro-2,4-pentadione **19** in refluxing ethanol to afford 1-(2-amino-4-methylthiazol-5-yl)ethanone **20** in nearly quantitative yield,³⁶ followed by bromination α to the carbonyl to give compound **21**. The subsequent condensation of intermediate **21** with 1-(3-acetylphenyl)thiourea **23** gave bithiazole **24** that was finally N-acetylated to obtain the desired compound **17**. Thiourea **23** was synthesized by reaction of 3'-aminoacetophenone **22** with benzoyl isothiocyanate, followed by a basic hydrolysis to remove the benzoyl group.³⁷

Docking studies on compound **17** (see Molecular modeling and SAR section) showed a pattern of interactions within the ATP-binding pocket of PI4KIII β very similar to that of the reference compound **1**. The proposed binding mode of **17** suggested that two main portions of this molecule could be functionalized to explore the biologically relevant chemical space: *i*)


the 2-amino group on the 4-methylthiazole ring (left part) and, *ii*) the phenyl ring (right part). We first explored the chemical space around the left part of compound **17**, introducing bulkier groups and urea/thiourea functions in place of the acetamide moiety. The intermediate **24** represents in fact an advanced intermediate that could be easily functionalized on the 2-amino group to give a series of functionalized derivatives (**25a-g** and **26a-d**) (Scheme 2).



Scheme 2. Reagents and conditions: (a) method A (for **25a-c**), R¹COCl (for **25a,b**) or (R¹CO)₂O (for **25c**), Et₃N, DCM, rt, 12-15 h, 65-80%; method B (for **25d-f**) R¹COCl, Et₃N, DCM, reflux, 15 h, 65-75%; method C (for **25g**), (R¹CO)₂O, Et₃N, DMF, 50 °C, 12 h, 63%; (b) R²NCX, pyridine, reflux, 12-18 h, 52-69%.


Compound **24** was first reacted with different acyl chlorides or anhydrides to obtain compounds **25a-g**, while the urea/thiourea derivatives **26a-d** were synthesized by reacting **24** with the appropriate isocyanates/isothiocyanates. We also decided to replace the acetamide moiety of compound **17** with chain-extended ureidic groups for the SAR development. Unfortunately, the synthesis of compounds **30a,b** following the approach described above would have required very expensive isocyanates. Thus, an alternative synthetic approach was used for the synthesis of **30a,b** (Scheme 3): starting from 1-(2-amino-4-methylthiazol-5-yl)ethanone **20**, reaction with diphenyl carbonate gave good yields of the desired phenyl

carbamate **27** that reacted readily with the appropriate amines to give the urea intermediates **28a,b**.³⁸ Similar to the synthesis of compound **17**, the bromination α to the carbonyl and the subsequent condensation of intermediates **29a,b** with the 1-(3-acetylphenyl)thiourea **23**, gave the desired compounds **30a,b**.



Scheme 3. Reagents and conditions: (a) diphenyl carbonate, NaH, DMF, rt, 30 min, 67%; (b) 4-chlorobenzylamine (for **28a**) or 1-adamantylamine (for **28b**), THF, 50 °C, 5-6 h, 61-93%; (c) 48% aqueous HBr, Br₂, 1,4-dioxane, 60 °C, 3 h, 88-92%; (d) EtOH, reflux, 1 h, 78-83%.


We next explored the right part of compound **17**, keeping the 2-acetamido group on the left part of the molecule unchanged and modifying the substitution pattern of the phenyl ring on the right part. Since the acetamide moiety on the left part of the molecule was conserved, intermediate **20** was conveniently acetylated before the bromination α to the carbonyl (Scheme 4). Final compounds **35a-d** were quickly obtained in good yields by reacting **32** with substituted thioureas **34a-d**, previously synthesized from the corresponding amines **33a-d**.




Scheme 4. Reagents and conditions: (a) acetyl chloride, pyridine, THF/DCM, 0 °C, 3 h, 87%; (b) Br₂, 1,4-dioxane, 50 °C, 22 h, 84%; (c) (i) benzoyl isothiocyanate, DCM, rt, 12 h, (ii) NaOH 1N, THF, reflux, 2 h, 72-78%; (d) EtOH, reflux, 1h, 71-85%.

Then we decided to modify the central bithiazole scaffold of the hit compound **17**, to get additional SAR information. As described in Scheme 5, we first introduced an imidazole ring by reacting intermediate **21** with the 1-(3-acetylphenyl)guanidine **36**, obtained by treating 3'-aminoacetophenone **22** with cyanamide.³⁹ Compound **38** was thus synthesized by acylation of the intermediate **37** with acetyl chloride.

A scaffold hopping approach (FAF-drugs2 server)⁴⁰ was also employed to identify alternatives to the bithiazole scaffold: among the molecules proposed by the software, the asymmetrical N,N'-diarylthiourea scaffold was considered the most promising on the basis of its synthetic accessibility and the antiviral activity of some closely related analogues reported in the literature.⁴¹ Intermediate **41** was easily obtained by addition of 3'-aminoacetophenone **22** to 3-nitrophenyl isothiocyanate **39**, followed by reduction of the nitro group with iron powder in acidic ethanol (Scheme 6). The subsequent acylation of the amino group led to final compounds **42a,b**.




Scheme 5. Reagents and conditions: (a) cyanamide, HNO₃, EtOH/H₂O, reflux, 24 h, 73%; (b) Et₃N, EtOH, reflux, 12 h, 82%; (c) acetyl chloride, Et₃N, DCM, rt, 8 h, 47%.




Scheme 6. Reagents and conditions: (a) DCM, rt, 18 h, 88%; (b) Fe, HCl, EtOH, reflux, 2 h, 75%; (c) acetyl chloride (for 42a) or trimethylacetyl chloride (for 42b), pyridine, THF, rt, 2 h, 67-69%.

Moreover we noted high chemical similarity between compound **17** and compound **46**, a known inhibitor of DC-SIGN (dendritic cell (DC)-specific intercellular adhesion molecule-3 grabbing nonintegrin).⁴² The role of DC-SIGN in the binding and transmission of different pathogens, including enteroviruses, has been well investigated.⁴³ So we decided to evaluate the antiviral effect of compound **46** in a virus-cell-based assay and its activity on PI4KIII β . As described in Scheme 7, compound **46** was synthesized following the procedure previously reported for compound **17**.



Scheme 7. Reagents and conditions: (a) (i) benzoyl isothiocyanate, DCM, rt, 12 h, (ii) NaOH 1N, THF, reflux, 3 h, 80%; (b) EtOH, reflux, 30 min, 77%; (c) propionyl chloride, Et₃N, DCM, rt, 8 h, 68%.

Finally, we decided to prepare two compounds (related to the hit **17**) known in the CFTR field to evaluate the potential role of the CFTR channel in viral replication: *i*) compound **47a**, which is active in correcting the F508del-CFTR defect and, *ii*) compound **47b**, which is inactive in correcting the F508del-CFTR defect (Figure 3). Compound **47a,b** were synthesized following the procedure reported in literature.⁴⁴



47a: X = O

47b: X = S

Figure 3. Chemical structures of compounds **47a,b**.

Biology

All the synthesized compounds were initially evaluated for their inhibitory potency against PI4KIII β kinase *in vitro* and for their cell-based antiviral activity: EV71 was used as the primary target for SAR exploration since compound **17** revealed the best and most reproducible antiviral activity against this virus. In particular, the antiviral activity against EV71 was evaluated in EV71-induced CPE-reduction assay in rhabdosarcoma (RD) cells. Both the EC₅₀ values and the CC₅₀ values were measured. Uninfected, treated cells were also inspected under the microscope to evaluate whether the compounds altered normal cell morphology. The EC₅₀ and CC₅₀ values allowed us to calculate the selectivity index (SI), defined as CC₅₀/EC₅₀. Compound **3** was used as a positive control. Results are summarized in Table 1.

A close correlation between the antiviral activity measured in the cell-based assay and the inhibitory potency of the PI4KIII β kinase was observed, with only a few exceptions. The best results were obtained via modifying the left part of the molecule. In particular compounds **25a,b**, bearing respectively a propanamide and a pivalamide moiety instead of the acetamide function of compound **17**, showed a very promising antiviral activity.

Table 1. Activity of synthesized derivatives in PI4KIII β inhibition assay and in virus-cell-based EV71 assay.

Compd	PI4K III β IC ₅₀ (μ M) ^a	EV71 EC ₅₀ (μ M)	EV71 CC ₅₀ (μ M) ^b	EV71 CC ₅₀ (μ M) ^c	SI ^d	SI ^d
17	0.48	0.92 \pm 2.75	16.5 \pm 9.04	9.73 \pm 0.87	17.9	10.6
25a	0.27	0.38 \pm 0.10	10.1 \pm 4.82	6.37 \pm 2.46	26.6	23.6
25b	0.32	0.27 \pm 0.05	7.94 \pm 1.24	8.83 \pm 0.89	29.4	32.7
25c	21.89	2.0 \pm 0.95	25.8 \pm 5.06	30.1 \pm 9.93	12.9	15.0
25d	18.85	0.51 \pm 0.14	5.75 \pm 2.12	11.8 \pm 4.56	11.3	23.1
25e	>50	>44.6	ND ^e	ND	ND	ND
25f	>50	>55.7	ND	ND	ND	ND
25g	4.67	1.42 \pm 0.04	42.3 \pm 4.94	101.0 \pm 15.8	29.7	71.1
26a	7.69	NA ^f	ND	ND	ND	ND
26b	>50	NA	ND	ND	ND	ND
26c	1.82	2 \pm 0.04	8.87 \pm 3.85	ND	4.4	ND
26d	>50	NA	ND	ND	ND	ND
30a	3.95	4.77 \pm 0.28	20.9 \pm 6.97	ND	4.4	ND
30b	12.40	NA	ND	ND	ND	ND
35a	2.48	1.93 \pm 0.81	18.6 \pm 6.01	ND	9.64	ND
35b	2.63	NA	ND	ND	ND	ND
35c	1.55	1.2 \pm 0.16	9.17 \pm 0.85	ND	5.9	ND
35d	3.71	0.68 \pm 0.04	5.59 \pm 0.32	ND	8.2	ND
38	>50	>90.2	85.7	64.2	ND	ND
42a	NA	NA	ND	ND	ND	ND
42b	NA	NA	ND	ND	ND	ND
46	50.00	8.58 \pm 0.77	51.6 \pm 27.1	142.0 \pm 62.8	6	ND
47a	NA	NA	ND	ND	ND	ND
47b	NA	NA	ND	ND	ND	ND
3^g	0.06	0.73	>125	-	-	-

^aValues are the mean of at least three independent experiments. ^bCC₅₀ values were assessed by MTS method. ^cCC₅₀ values were determined by microscopically detectable alteration of cell morphology. ^dSelectivity index (SI = CC₅₀/EC₅₀). ^eND = not determined. ^fNA = not active. ^gReference 13

Compounds **25a,b** inhibited PI4KIII β and exhibited a significant antiviral effect at sub-micromolar concentrations, demonstrating a better activity than compound **17**. Compound **25g**, characterized by the Boc amino group, proved to be the most interesting compound of the entire series showing the highest selectivity index in the EV71 cell-based assay. Also changing the right portion of hit compound **17** gave interesting results (compounds **35a,c,d**). The central bithiazole scaffold proved to be essential for antiviral activity, as changing it gave inactive compounds (compounds **38, 42a,b**). Finally, the reported compounds **46, 47a,b** were devoid of antiviral activity and PI4KIII β inhibition activity.

Based on these activity data, and considering that the SI of a promising antiviral candidate should be at least greater than 10, compounds **17, 25a-d, 25g** were selected for further studies. The broad-spectrum activity of the six selected compounds was evaluated against a panel of enteroviruses representative of all major groups: enterovirus group B (coxsackievirus B3 and echovirus11, ECHO11), group C (poliovirus 1), group D (enterovirus 68), rhinovirus group A (RV02) and rhinovirus group B (RV14) (see the Experimental Section for details). Results are reported in Table 2. The selected compounds showed micromolar and sub-micromolar activity against different enteroviruses within the tested panel. In addition, the antiviral activity of the less toxic compound **25g** was confirmed against a representative panel of EV71 clinical isolates. As shown in Table 3, we could confirm the activity of compounds **25g** against the clinically relevant EV71 specimens. Only the (sub)genogroup B5 appeared to be less sensitive. Furthermore we evaluated the lipid kinase isoform selectivity of our best PI4KIII β inhibitors by testing them in an *in vitro* inhibition assay on the related enzyme PI4KIII α and PI3K- α /p85 α (Table 4).

Table 2. Evaluation of the broad-spectrum antiviral activity of the most potent derivatives against a representative panel of enteroviruses.

Compd	CVB3		ECHO11		PV1		EV68		RV14		RV02	
	EC ₅₀ (μ M)	CC ₅₀ (μ M)	EC ₅₀ (μ M)	CC ₅₀ (μ M)	EC ₅₀ (μ M)	CC ₅₀ (μ M)	EC ₅₀ (μ M)	CC ₅₀ (μ M)	EC ₅₀ (μ M)	CC ₅₀ (μ M)	EC ₅₀ (μ M)	CC ₅₀ (μ M)
17	2.17	101 \pm 33.6	1.57 \pm 0.23	>268	3.09	>269	NA ^a	ND ^b	>268	ND	NA ^b	ND
25a	2.16	89.1 \pm 16.2	0.97	12 \pm 4. 07	<1.52	5.49 \pm 1.42	NA	ND	NA	ND	>259	ND
25b	ND	20.9 \pm 3.83	0.72 \pm 0.05	5.07 \pm 1.24	<1.41	5.07 \pm 1.3	1.38	3.09	4.85 \pm 1.09	ND	2.01 \pm 0.05	ND
25c	3.87 \pm 0.23	58.8 \pm 7.14	3.51	29.3 \pm 1.78	2.86 \pm 0.36	30.3 \pm 3.66	ND	ND	10.6 \pm 0.7	ND	10.6 \pm 0.2	ND
25d	2.19 \pm 0.23	45.1 \pm 4.44	1.77 \pm 0.13	145 \pm 2 7.6	2.23 \pm 0.4	145 \pm 2 7.6	NA	ND	>230	ND	2.05 \pm 0.36	ND
25g	2.74 \pm 0.17	80.1 \pm 6.97	2.93	124 \pm 7 .94	13.3 \pm 1.9	124 \pm 7 .94	ND	ND	>232	ND	ND	ND

^aNA = not active. ^bND = not determined

Results showed a higher specificity of the tested bithiazole derivatives for the PI4KIII β isoform with poor inhibition of both PI4KIII α and PI3K- α /p85 α at 100 μ M concentration of each compound. The specificity of compound **25g** was also tested on a small panel of unrelated kinases: it shows only a low inhibitory effect on Src and CDK6. Despite the latter enzymes being involved in cell cycle regulation and representing common targets of antitumor compounds, **25g** did not show any toxicity or morphology alteration at antiviral concentration in the tested cell lines. In addition, recent studies indicated that Src inhibitors have no effect on EV71 replication⁴⁵ while CDK6 seems to be down-regulated in response to EV71 infection.⁴⁶ Finally, compounds reported in Table 4 were evaluated for their CFTR

Table 3. Evaluation of the antiviral activity of compound **25g** against EV71 clinical isolates.

Genogroup	Strain	Genbank	EC ₅₀ (μM) ^a
			Compd 25g
B2	11316	AB575927	<1.39
B5	TW/96016/08	GQ231942	21.00
	TW/70902/08	GQ231936	3.58
C2	H08300 461#812	-	0.97
C4	TW/1956/05	GQ231926	<1.39
	TW/2429/04	GQ231927	1.17

^aAll values are based on at least three independent dose-response curves.

Table 4. Inhibitory effect of selected compounds against members of PIK family and profiling of compound **25g** against a small panel of unrelated kinases.

Compd	PI4KIIIβ	PI4KIIIα	PI3K- α/p85α	Compd	kinase	% Residual activity at 100 μM ^a
	IC ₅₀ (μM) ^a	% Residual activity at 100 μM ^a				
17	0.48	52	37	25g	Src FL	32
25a	0.27	72	40		GSK3β	79
25b	0.32	71	93		Hck FL	100
25c	21.89	73	85		FAK	82
25d	18.85	81	54		DYRK1A	88
25g	4.67	69	64		ABL FL	53
35a	2.48	58	81		FLT3	59
35c	1.55	71	51		CDK2/cA2	62
35d	3.71	73	100		CDK9/cT1	57
					CDK9/cK	49
				CDK6/cD1	25	
				Pim1	70	

^aValues are the mean of two independent replicates

corrector/potentiator activity, to identify molecules that may be endowed with dual antiviral/CFTR modulator activity. As shown in Figure 4A, some of the compounds (**25a**, **25d**, **25g**) acted as CFTR correctors, increasing steady-state levels of F508del-CFTR at the plasma membrane after chronic (24 hour) incubation. Compound **48** (Lumacaftor), the leading corrector drug,⁴⁷ was used as a benchmark. This increased CFTR plasma membrane density was measured with a recently developed assay exploiting a CFTR fusion to a pH-sensitive protein.⁴⁸ The improvement in biogenesis also lead to increased anion permeability, estimated from fluorescence quenching of a CFTR-fused YFP probe following extracellular I⁻ addition (Figure 4B). None of the compounds acted as "potentiators" rapidly increasing anion permeability, when added only immediately prior to I⁻ addition (Figure 4C). The approved potentiator drug **49** (Ivacaftor) was used as a comparison.⁴⁹ Overall, the drug-induced changes in the iodide entry rate and in membrane density followed similar patterns, suggesting that the chemically corrected molecules of F508del-CFTR that reached the plasma membrane displayed an ion-channel function similar to those corrected by treatment with **48**. However, compound **25d** appears to increase CFTR membrane density more than expected from its effect on anion permeability (Figure 4D). Further studies will be required to understand the underlying mechanism. Overall, the collected biological data indicates that a fine chemical tuning of the bithiazole substituents is needed to generate compounds able to specifically inhibit the PI4KIII β kinase and block the replication of different enteroviruses while also correcting the F508del-CFTR folding defect. It is interesting to note that the most promising CFTR correctors (**25a**, **25d**, **25g**) are also the most active broad-spectrum antivirals and represent the first example of multi-target agents for tightly associated pulmonary diseases like enterovirus infection and cystic fibrosis.




Figure 4. Effects of selected compounds on CFTR biogenesis and function. All treatments were carried out alongside low temperature incubation, known to improve F508del-CFTR membrane-localization and used to increase the fluorescence signal. A) F508del-CFTR-

1
2
3 pHTomato present at the plasma membrane was quantified following 24 h incubation in 10
4
5 μM of each drug. Incubation with **48** (Lumacaftor) was assessed in parallel, as a positive
6
7 control. B) Anion permeability quantified using a YFP-F508del-CFTR probe following 24 h
8
9 treatment as in A). C) Compounds (10 μM) do not cause an immediate change in anion
10
11 permeability. Potentiator compound **49** (Ivacaftor) was used as a positive control. D) For
12
13 most compounds, there is a similar ratio describing increase in membrane density over anion
14
15 permeability as caused by **48** (Lumacaftor).
16
17
18
19
20
21
22

23 **Molecular modelling and SAR**

24 The hit compound **17** was docked with the Glide software⁵⁰ (SP) in the ATP binding site of
25
26 the PI4KIII β crystal structure (PDB ID: 4D0L)³⁴ centering the grid on compound **1**. The
27
28 predicted binding mode and interaction profile of compound **17** is very similar to that of
29
30 compound **1** (Figure 5). The NH-acetamide moiety of **17** is hydrogen bonded to VAL598,
31
32 which also interact with the thiazole nitrogen. The thiazole ring is involved in a Pi-Pi
33
34 stacking with TYR583, while the phenyl ring is involved in a Pi-cation interaction with
35
36 LYS549. The O carbonyl moiety is hydrogen bonded to LYS377. Moreover, the binding
37
38 mode is completed by a series of hydrophobic interactions involving LEU383, ALA602,
39
40 VAL599, VAL602, LEU663, ILE595, TYR583, ILE671, ILE673, PRO381 and LEU374.
41
42 The first series of derivatives of the hit compound **17** encompasses different substitution on
43
44 the left part of the molecule (right part as per Figure 5-7 representation) by replacing the
45
46 methyl group of the acetamide moiety with different groups. Changing the methyl of the
47
48 acetamide with an ethyl or *tert*-butyl group (**25a,b**) caused a 2-fold increase of potency, but
49
50 the substitution with more hydrophobic and therefore more sterically bulky groups first
51
52 reduced the potency (**25d**) and then led to a complete loss of activity (**25e,f**). The limit seems
53
54
55
56
57
58
59
60

to be a *tert*-butoxycarbonyl group (**25g**) which still maintains low micromolar activity (Figure 6).




Figure 5. a) Predicted binding mode of the hit **17** (magenta sticks) superimposed to compound **1** (green sticks) into the binding site of PI4KIIIβ. b) 2D ligand interaction diagram of **17**.

The substitution of the acetamide methyl with a CF₃ (**25c**) weakened the inhibition (45-fold decrease) not due to steric reasons, but probably because of the electron withdrawing properties of the CF₃ group. Changing the acetamide moiety with a ureidic moiety caused a slight decrease in potency (**26a,c** and **30a,b**). The ureidic portion seems to interact with a double hydrogen bond to VAL598 but at the same time, the hydrophobic NH substituent moves away from the protein resulting in a solvation penalty (Figure 6). Finally, modifying a ureidic group with a thioureidic moiety caused a complete loss of activity (**26b,d**).




Figure 6. a) Superimposed binding modes of compounds **17** (purple sticks), **25a** (orange sticks), **25b** (green sticks) and **25g** (pink sticks) into the binding site of PI4KIIIβ. b) Superimposed binding modes of compounds **26a** (orange sticks), **26c** (green sticks) and **30a** (pink sticks) into the binding site of PI4KIIIβ. c) 2D ligand interaction diagram of **25b**. d) 2D ligand interaction diagram of **30a**.

The second series of derivatives of the hit compound **17** comprises different substitutions on the right part of the molecule (left part as per Figure 5-7 representation) by changing the substitution pattern of the phenyl ring. A change in the position of the acetyl group reduces the potency by about 10-fold (**35d**), and the same happens by introducing hydroxy and

1
2
3 methoxy groups in the meta position (**35b,c**). The precise positioning of the hydrogen-bond
4
5 acceptor on the phenyl ring seems therefore important in improving the affinity for the
6
7 enzyme. Surprisingly, the deletion of the ketonic group reduces but does not abolish activity
8
9 (**35a**). The lack of the substituent on the phenyl ring deletes a hydrogen bond, but allows
10
11 optimization of the other interactions, in particular the Pi-cation with lysine 549.
12
13

14
15 The last series of derivatives of the hit compound **17** encompasses modifications of the
16
17 bithiazole scaffold and closely related analogues reported as DC-SIGN inhibitors and CFTR
18
19 correctors. Replacement of the thiazole group of compound **17** with an imidazole (**38**) or
20
21 conversion of the bithiazole scaffold into a N,N'-diarylthiourea (**42a,b**) changes the binding
22
23 mode and abolishes the activity. Finally, also the DC-SIGN inhibitor **46** and CFTR correctors
24
25 **47a,b** presented a suboptimal interaction profile with PI4KIII β and resulted in a complete
26
27 loss of activity. As reported by Warrem et al.⁵¹, docking programs and scoring functions
28
29 present a few limitations in correlating subtle structural differences of active ligands with
30
31 their enzymatic activity. This error is quite limited within homologous series of compounds
32
33 but it can be very important for structurally unrelated compounds.⁵² Our docking studies were
34
35 in fact able to distinguish between active and inactive compounds but it is no coincidence that
36
37 the reference compound **3**, whose scaffold is very different from those of our series, showed a
38
39 docking score that is not in line with its enzymatic potency (Table 5). The most active
40
41 compounds **17**, **25a** and **25b** also showed the best specificity for PI4KIII β over PI4KIII α ,
42
43 which seems to depend on the steric hindrance of the acetamide substitution. In fact,
44
45 considering compounds **17**, **25a**, **25b**, **25g** and **25d** that differ only for the bulkiness of the
46
47 amide substituents in position C2 of the thiazole (methyl, ethyl, *tert*-butyl, *tert*-butoxy and
48
49 phenyl, respectively), the affinity for PI4KIII α decreases going from **17** to **25d** (see Table 4).
50
51
52
53
54
55
56
57
58
59
60

Table 5. Correlation between IC₅₀ and docking score for the synthesized compounds.

Compd	PI4K IIIβ IC ₅₀ (μM) ^a	Docking score ^b	Compd	PI4K IIIβ IC ₅₀ (μM) ^a	Docking score ^b
35d	3.71	-8,297	25e	>50	-7,431
30b	12.40	-8,222	26d	>50	-7,4
26a	7.69	-8,172	25f	>50	-7,356
26c	1.82	-8,108	38	>50	-7,355
17	0.48	-8,035	42a	>50	-7,251
25a	0.27	-8,012	26b	>50	-7,144
25d	18.85	-7,983	42b	>50	-7,052
30a	3.95	-7,97	25c	21.90	-6,678
35b	2.63	-7,726	3^c	0.06	-6,553
46	50.00	-7,644	47b	>50	-6,424
25g	4.67	-7,63	47a	>50	-6,148
35c	1.55	-7,605	25e	>50	-7,431
35a	2.48	-7,6			
25b	0.32	-7,484			

^aValues are the mean of at least three independent experiments. ^bDocking score was calculated by the software Gold and expressed as Kcal/mol. ^cReference 13

The higher the bulkiness in C2 of the thiazole, the lower the affinity for PI4KIIIα over PI4KIIIβ with the phenyl substituent (**25d**) representing the highest tolerated hindrance after which the inhibition of PI4KIIIβ is also compromised. The reason for the specificity of these compounds towards PI4KIIIβ seems to depend on the different opening (compared to the α isoform) of a specific loop that in the β isoform goes from ILE595 to ILE604 (Figure 7). In fact in this portion of protein, the sequence of PI4KIIIβ resembles more the sequence of PI3K (α, γ end δ) than the sequence of PI4KIIIα. In particular, the presence of a cysteine residue in position 30 of PI4KIIIα (corresponding to proline 597 in PI4KIIIβ) could change the fold of this “selectivity loop”.




Figure 7. Observed binding mode of **25a** (yellow sticks) superimposed to compound **1** (magenta sticks) into the binding site of PI4KIII β (green ribbons). The orange ribbons represent the modelled structure of PI4KIII α .

CONCLUSIONS

An increasing number of reports suggest a causal link between enterovirus infections and pulmonary exacerbations in CF patients. We report the discovery of a new class of multi-target agents active as broad-spectrum antivirals and correctors of the F508del-CFTR folding defect. To identify these drug candidates, we first carried out a virtual screening on the PI4KIII β (a host protein involved in enterovirus replication) catalytic site to select commercially available compounds: our choice was based on the best-predicted affinity for the target kinase and 2D similarity (in a few cases) to known CFTR correctors/potentiators. Among the selected compounds, hit **17** showed activity in cell-free PI4KIII β inhibition assay and cell-based enterovirus replication assays and was therefore considered a promising starting point for further structure-based optimization. A small collection of analogues of compound **17** was then designed, synthesized and biologically evaluated for their *i*) activity against panel of enteroviruses representative of all major groups; *ii*) inhibition of lipid kinases PI4KIII β , PI4KIII α and PI3K- α /p85 α ; *iii*) corrector/potentiator activity on F508del-CFTR.

1
2
3 Three compounds (**25a**, **25d**, **25g**) were finally identified as novel multi-target agents able to
4
5 act as broad-spectrum-antivirals (enterovirus family) and as correctors of F508del-CFTR
6
7 folding defect. These compounds represent a valuable starting point to develop a novel
8
9 polypharmacological approach for the treatment of closely-related pulmonary diseases such
10
11 as cystic fibrosis and enterovirus infections with a single pill.
12
13

14 15 16 17 **EXPERIMENTAL SECTION**

18 **Molecular modeling**

19 **Homology modeling**

20
21 The structure of PI4KIII β was built with the Prime⁵³ 38013 software on the basis of the
22
23 crystal structure 2CHZ using ClustalW for sequence alignment and knowledge-based as
24
25 building method. The structure of PI4KIII α was built with the online server 3D-JIGSAW⁵⁴ on
26
27 the basis of the crystal structure 4D0L.
28
29
30

31 **Molecular dynamics**

32
33 The structure of modelled PI4KIII β was aligned to the 2CHZ structure. Compound **1** was
34
35 extracted from 2CHZ and was manually introduced in the structure of PI4KIII β . A molecular
36
37 dynamics simulation of the resulting complex was performed using Desmond v40013. The
38
39 complex was neutralized using sodium counter ions. The complex and the counter ions were
40
41 immersed in a orthorhombic periodic SPC water bath that extended about 10 Å in each
42
43 direction. After an initial default relaxation protocol, a MD production run was performed for
44
45 10 ns with a time step of 2 fs.
46
47
48

49 **Virtual screening**

50
51 From the last 2 ns of the dynamics simulation 100 frames were extracted and clustered on
52
53 the basis of RMSD. Five clusters were generated. The protein representative of each cluster
54
55 was processed with the Schrödinger Suite 2014-3⁵⁵ Protein Preparation Wizard tool. On each
56
57
58
59
60

1
2
3 structure a grid was generated with the software Glide 65013 centering the grid on compound
4
5 **1**, then all PI4KIII β ligands available from the Pubchem database were docked with the SP
6
7 protocol. Structures were selected for virtual screening on the basis of their enrichment
8
9 factor. The Mid Asinex database was extracted from the ZINC database⁵⁶ and docked in the
10
11 binding site of compound **1** using compound **1** as the center of the grid. The software Glide
12
13 65013 with the SP protocol was used for High throughput docking. The best 1000
14
15 compounds in terms of the docking score were selected and submitted to one more docking
16
17 cycle of docking with the XP protocol. On the basis of the docking score and a visual
18
19 inspection 25 compounds were selected and 13 compounds were purchased.
20
21

22 **Ligand preparation**

23
24 Ligands were prepared with the LigPrep⁵⁷ tool available in the Schrödinger Suite 2015-4.
25
26 Ionization states were generated at pH 7.0 \pm 2.0 with Epik.
27
28

29 **Ligand docking**

30
31 The X-ray coordinates of PI4KIII β in complex with compound **1** were extracted from the
32
33 Protein Data Bank (PDB code 4D0L). The structure was then processed with the Schrödinger
34
35 Suite 2015-4 Protein Preparation Wizard tool.⁵⁸ The A Chain was selected, water molecules
36
37 were removed, and an exhaustive sampling of the orientations of groups, whose hydrogen
38
39 bonding network needs to be optimized, was performed. Finally, the protein structure was
40
41 refined to relieve steric clashes with a restrained minimization with the OPLS3 force field⁵⁹
42
43 until a final RMSD of 0.30 Å with respect to the input protein coordinates.
44
45

46
47 Docking studies were performed using Glide⁶⁰ 69017 with the SP protocol. The protein
48
49 structure, prepared as described above, was used to build the energy grid. The enclosing box
50
51 was centered on the cocrystallized ligand. All parameters were set to their default value. The
52
53 docking protocol was validated by redocking the cocrystallized ligand (compound **1**).
54
55
56
57
58
59
60

Chemistry

General. All commercially available chemicals were purchased from both Sigma- Aldrich and Alfa Aesar and, unless otherwise noted, used without any previous purification. Solvents used for work-up and purification procedures were of technical grade. Dry solvents used in the reactions were obtained by distillation of technical grade materials over appropriate dehydrating agents. Reactions were monitored by thin layer chromatography on silica gel-coated aluminium foils (silica gel on Al foils, SUPELCO Analytical, Sigma-Aldrich) at 254 and 365 nm. Where indicated, products were purified by silica gel flash chromatography on columns packed with Merck Geduran Si 60 (40-63 μm). ^1H and ^{13}C NMR spectra were recorded on BRUKER AVANCE 300 MHz and BRUKER AVANCE 400 MHz spectrometers. Chemical shifts (δ scale) are reported in parts per million relative to TMS. ^1H -NMR spectra are reported in this order: multiplicity and number of protons; signals were characterized as: *s* (singlet), *d* (doublet), *t* (triplet), *q* (quadruplet), *m* (multiplet), *bs* (broad signal). ESI-mass spectra were recorded on an API 150EX apparatus and are reported in the form of (m/z). Elemental analyses were performed on a Perkin-Elmer PE 2004 elemental analyzer. Melting points were taken using a Gallenkamp melting point apparatus and were uncorrected. All final compounds showed chemical purity $\geq 95\%$ as determined by elemental analysis data for C, H, and N (within 0.4% of the theoretical values).

Synthesis of N-(2-((3-acetylphenyl)amino)-4'-methyl-[4,5'-bithiazol]-2'-yl)acetamide (17).

Et_3N (84 μL , 0.60 mmol) was added to a stirred suspension of intermediate **24** (100 mg, 0.30 mmol) in dry DCM (4 mL) at 0 $^\circ\text{C}$. After 15 minutes acetyl chloride (32 μL , 0.45 mmol), diluted in dry DCM (0.5 mL), was added dropwise. The resulting solution was warmed to room temperature and stirred for 15 h. Next, H_2O and DCM were added and the aqueous phase was extracted twice with DCM. The combined organic phases were washed with brine,

1
2
3 dried over Na₂SO₄ and evaporated. The crude was purified by flash chromatography using
4 DCM/MeOH (98/2) as eluent to afford compound **17** as a yellow solid. Yield 77%; mp 244-
5 246 °C. MS (ESI) [M + H]⁺: 373.3 *m/z*. ¹H NMR (DMSO-*d*₆ 300 MHz): δ 2.14 (s, 3H), 2.51
6 2.51 (s, 3H), 2.63 (s, 3H), 6.95 (s, 1H), 7.48 (t, 1H, *J* = 7.9 Hz), 7.56 (d, 1H, *J* = 7.9 Hz), 7.77 (d,
7 1H, *J* = 7.9 Hz), 8.51 (s, 1H), 10.56 (s, 1H), 12.06 (s, 1H). ¹³C NMR (DMSO-*d*₆ 100.6 MHz):
8 δ 17.56, 22.92, 27.42, 103.25, 116.75, 120.52, 121.44, 121.65, 129.79, 138.11, 141.81,
9 143.21, 143.29, 155.58, 162.92, 168.73, 198.28. Anal. (C₁₇H₁₆N₄O₂S₂) C, H, N.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Synthesis of 1-(2-amino-4-methylthiazol-5-yl)ethanone (20). A solution of thiourea **18** (283
mg, 3.72 mmol) and 3-chloro-2,4-pentanedione **19** (419 μL, 3.72 mmol) in ethanol (20 mL)
was heated at reflux for 12 h, and then the reaction mixture was cooled down to 0 °C. The
precipitate was separated by filtration over a Buchner funnel and washed with cold ethanol
and ether to afford the product **20** as a white solid. Yield 95%. MS (ESI) [M + H]⁺: 157.2
m/z. ¹H NMR (DMSO-*d*₆ 400 MHz): δ 2.44 (s, 3H), 2.52 (s, 3H), 9.49 (bs, 2H).

Synthesis of 1-(2-amino-4-methylthiazol-5-yl)-2-bromoethanone (21). A suspension of
intermediate **20** (500 mg, 3.20 mmol) in 48% HBr solution in water (10 mL) was warmed to
60 °C. A solution of Br₂ (148 μL, 2.88 mmol) in 1,4-dioxane (10 mL) was added dropwise
and the reaction mixture was heated at 60 °C for 3 h. After cooling down to room
temperature, saturated aqueous NaHCO₃ solution and ethyl acetate were added and the
aqueous phase was extracted three times with ethyl acetate. The combined organic phases
were washed with brine, dried over Na₂SO₄ and concentrated under vacuum to obtain
compound **21**, used in the next step without any further purification. Yield 90%. MS (ESI)
[M + H]⁺: 235.0 *m/z*, [M + 2 + H]⁺: 237.1 *m/z*. ¹H NMR (DMSO-*d*₆ 400 MHz): δ 2.46 (s,
3H), 4.48 (s, 2H), 9.18 (bs, 2H).

Synthesis of 1-(3-acetylphenyl)thiourea (23). Benzoyl isothiocyanate (547 μL, 4.07 mmol)
was added dropwise to a solution of 3'-aminoacetophenone **22** (500 mg, 3.70 mmol) in dry

1
2
3 DCM (12 mL) and the mixture was stirred at room temperature for 12 h. The solvent of
4
5 reaction was evaporated, the residue was dissolved in THF/NaOH 1N (1/1, 15 mL) and the
6
7 mixture was refluxed for 3 h. After cooling to room temperature, H₂O and ethyl acetate were
8
9 added and the aqueous phase was extracted twice with ethyl acetate. The combined organic
10
11 phases were dried over Na₂SO₄ and evaporated. The resulting solid was crystallized from
12
13 ether. Yield 72%. MS (ESI) [M + H]⁺: 195.1 *m/z*. ¹H NMR (DMSO-*d*₆ 400 MHz): δ 2.57 (s,
14
15 3H), 7.46 (t, 1H, *J* = 7.9 Hz), 7.55 (bs, 2H), 7.70-7.73 (m, 2H), 8.03 (s, 1H), 9.89 (s, 1H).
16
17

18
19 *Synthesis of 1-(3-((2'-amino-4'-methyl-[4,5'-bithiazol]-2-yl)amino)phenyl)ethanone (24).*

20
21 Intermediates **21** (200 mg, 0.85 mmol) and **23** (165 mg, 0.85 mmol) were suspended in
22
23 ethanol (5 mL) and the mixture was heated at reflux for 1 h. Then saturated aqueous NaHCO₃
24
25 solution and ethyl acetate were added to the mixture and the aqueous phase was extracted
26
27 three times with ethyl acetate. The combined organic phases were washed with brine, dried
28
29 over Na₂SO₄ and concentrated under vacuum. Ether was added to the crude and the solid
30
31 obtained was filtered over a Buchner funnel, washed with ether and used in the following
32
33 step without any further purification. Yield 84%. MS (ESI) [M + H]⁺: 331.3 *m/z*. ¹H NMR
34
35 (DMSO-*d*₆ 300 MHz): δ 2.35 (s, 3H), 2.60 (s, 3H), 6.67 (s, 1H), 7.09 (bs, 2H), 7.46 (t, 1H, *J*
36
37 = 7.9 Hz), 7.55 (d, 1H, *J* = 7.9 Hz), 7.75 (d, 1H, *J* = 7.9 Hz), 8.40 (s, 1H), 10.55 (s, 1H).
38
39
40

41 *General Procedure for the Synthesis of Compounds 25a-g.*

42
43 *Method A (for 25a-c).* Et₃N (84 μL, 0.60 mmol) was added to a stirred suspension of
44
45 intermediate **24** (100 mg, 0.30 mmol) in dry DCM (4 mL) at 0 °C. After 15 minutes the
46
47 proper acyl chlorides or anhydride (0.45 mmol), diluted in dry DCM (0.5 mL), were added
48
49 dropwise. The resulting solution was warmed to room temperature and stirred for 12-15 h.
50
51 Next, H₂O and DCM were added and the aqueous phase was extracted twice with DCM. The
52
53 combined organic phases were washed with brine, dried over Na₂SO₄ and evaporated. The
54
55 crude was purified by flash chromatography using DCM/MeOH (98/2) as eluent.
56
57
58
59
60

1
2
3 *N*-(2-((3-acetylphenyl)amino)-4'-methyl-[4,5'-bithiazol]-2'-yl)propionamide (**25a**). Yield
4 75%; mp 244-246 °C. MS (ESI) [M + H]⁺: 387.1 *m/z*. ¹H NMR (DMSO-*d*₆ 400 MHz): δ 1.11
5 (t, 3H, *J* = 7.5 Hz), 2.45 (q, 2H, *J* = 7.5 Hz), 2.52 (s, 3H), 2.64 (s, 3H), 6.95 (s, 1H), 7.50 (t,
6 1H, *J* = 7.9 Hz), 7.57 (d, 1H, *J* = 7.9 Hz), 7.82 (d, 1H, *J* = 7.9 Hz), 8.47 (s, 1H), 10.56 (s,
7 1H), 12.03 (s, 1H). ¹³C NMR (DMSO-*d*₆ 100.6 MHz): δ 9.63, 17.56, 27.43, 28.69, 103.25,
8 116.72, 120.47, 121.49, 121.63, 129.82, 138.11, 141.82, 143.24, 143.33, 155.62, 162.94,
9 172.36, 198.28. Anal. (C₁₈H₁₈N₄O₂S₂) C, H, N.

10
11
12
13
14
15
16
17
18 *N*-(2-((3-acetylphenyl)amino)-4'-methyl-[4,5'-bithiazol]-2'-yl)pivalamide (**25b**). Yield 80%;
19 mp 232-234 °C. MS (ESI) [M + H]⁺: 415.4 *m/z*. ¹H NMR (DMSO-*d*₆ 400 MHz): δ 1.25 (s,
20 9H), 2.53 (s, 3H), 2.63 (s, 3H), 6.95 (s, 1H), 7.50 (t, 1H, *J* = 7.9 Hz), 7.57 (d, 1H, *J* = 7.9 Hz),
21 7.85 (d, 1H, *J* = 7.9 Hz), 8.43 (s, 1H), 10.57 (s, 1H), 11.79 (s, 1H). ¹³C NMR (DMSO-*d*₆
22 100.6 MHz): δ 16.45, 26.90, 27.05 (3x), 39.24, 102.80, 117.29, 120.96, 122.25, 122.46,
23 129.67, 138.02, 140.79, 142.33, 143.58, 156.98, 163.28, 176.67, 198.36. Anal.
24 (C₂₀H₂₂N₄O₂S₂) C, H, N.

25
26
27
28
29
30
31
32
33 *N*-(2-((3-acetylphenyl)amino)-4'-methyl-[4,5'-bithiazol]-2'-yl)-2,2,2-trifluoroacetamide
34 (**25c**). Yield 65%; mp 254-256 °C. MS (ESI) [M + H]⁺: 427.1 *m/z*. ¹H NMR (DMSO-*d*₆ 400
35 MHz): δ 2.54 (s, 3H), 2.63 (s, 3H), 7.14 (s, 1H), 7.50 (t, 1H, *J* = 7.9 Hz), 7.58 (d, 1H, *J* = 7.9
36 Hz), 7.75 (d, 1H, *J* = 7.9 Hz), 8.47 (s, 1H), 10.64 (s, 1H), 14.05 (s, 1H). ¹³C NMR (DMSO-*d*₆
37 100.6 MHz): δ 15.84, 27.41, 105.27, 115.80, 116.68, 120.50, 121.44, 121.79, 129.86, 138.12,
38 141.59, 143.31, 143.48, 155.43, 163.45, 168.73, 198.21. Anal. (C₁₇H₁₃F₃N₄O₂S₂) C, H, N.

39
40
41
42
43
44
45
46
47 *Method B (for 25d-f)*. Et₃N (84 μL, 0.60 mmol) was added to a stirred suspension of
48 intermediate **24** (100 mg, 0.30 mmol) in dry DCM (4 mL) at 0 °C. After 15 minutes the
49 proper acyl chlorides (0.45 mmol), diluted in dry DCM (0.5 mL), were added dropwise. The
50 resulting solution was warmed to room temperature and heated at reflux for 15 h. Next, H₂O
51 and DCM were added and the aqueous phase was extracted twice with DCM. The combined
52
53
54
55
56
57
58
59
60

1
2
3 organic phases were washed with brine, dried over Na₂SO₄ and evaporated. The crude was
4
5 purified by flash chromatography using DCM/MeOH (99/1) as eluent.
6

7 *N*-(2-((3-acetylphenyl)amino)-4'-methyl-[4,5'-bithiazol]-2'-yl)benzamide (**25d**). Yield 75%;
8
9 mp 199-200 °C. MS (ESI) [M + H]⁺: 435.3 *m/z*. ¹H NMR (DMSO-*d*₆ 400 MHz): δ 2.58 (s,
10 3H), 2.66 (s, 3H), 7.02 (s, 1H), 7.49-7.66 (m, 5H), 7.83 (d, 1H, *J* = 7.9 Hz), 8.12 (d, 2H, *J* =
11 7.8 Hz), 8.50 (s, 1H), 10.59 (s, 1H), 12.62 (s, 1H). ¹³C NMR (DMSO-*d*₆ 100.6 MHz): δ
12 17.42, 27.44, 103.60, 116.76, 121.52, 121.66, 128.61, 129.02, 129.06 (2x), 129.83 (2x),
13 132.59, 133.31, 138.12, 141.82, 143.23, 143.29, 155.72, 163.02, 167.79, 198.30. Anal.
14 (C₂₂H₁₈N₄O₂S₂) C, H, N.
15
16
17
18
19
20
21

22 *N*-(2-((3-acetylphenyl)amino)-4'-methyl-[4,5'-bithiazol]-2'-yl)-4-iodobenzamide (**25e**).
23 Yield 65%; mp 220-223 °C. MS (ESI) [M + H]⁺: 561.3 *m/z*. ¹H NMR (DMSO-*d*₆ 400 MHz):
24 δ 2.58 (s, 3H), 2.66 (s, 3H), 7.01 (s, 1H), 7.50 (t, 1H, *J* = 7.9 Hz), 7.58 (d, 1H, *J* = 7.9 Hz),
25 7.81 (d, 1H, *J* = 7.9 Hz), 7.88 (d, 2H, *J* = 8.5 Hz), 7.94 (d, 2H, *J* = 8.5 Hz), 8.50 (s, 1H),
26 10.58 (s, 1H), 12.70 (s, 1H). ¹³C NMR (DMSO-*d*₆ 100.6 MHz): δ 17.33, 27.45, 101.14,
27 103.65, 116.77, 121.03, 121.51, 121.67, 129.82, 130.44 (2x), 132.34, 137.95 (2x), 138.12,
28 141.81, 143.16, 143.29, 154.99, 163.02, 165.15, 198.30. Anal. (C₂₂H₁₇IN₄O₂S₂) C, H, N.
29
30
31
32
33
34
35
36
37

38 *N*-(2-((3-acetylphenyl)amino)-4'-methyl-[4,5'-bithiazol]-2'-yl)-4-methylbenzamide (**25f**).
39 Yield 72%; mp 243-244 °C. MS (ESI) [M + H]⁺: 449.2 *m/z*. ¹H NMR (DMSO-*d*₆ 400 MHz):
40 δ 2.40 (s, 3H), 2.58 (s, 3H), 2.66 (s, 3H), 7.01 (s, 1H), 7.36 (d, 2H, *J* = 8.0 Hz), 7.50 (t, 1H, *J*
41 = 7.9 Hz), 7.57 (d, 1H, *J* = 7.9 Hz), 7.82 (d, 1H, *J* = 7.9 Hz), 8.02 (d, 2H, *J* = 8.0 Hz), 8.50 (s,
42 1H), 10.58 (s, 1H), 12.55 (s, 1H). ¹³C NMR (DMSO-*d*₆ 100.6 MHz): δ 17.76, 21.55, 27.44,
43 103.55, 116.75, 120.52, 121.51, 121.66, 128.63 (2x), 129.62 (2x), 129.83, 131.51, 138.13,
44 141.42, 141.83, 143.26, 143.39, 155.78, 163.00, 168.09, 198.31. Anal. (C₂₃H₂₀N₄O₂S₂) C, H,
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60 N.

1
2
3 *Method C (for 25g)*. Et₃N (84 μL, 0.60 mmol) was added to a stirred suspension of
4 intermediate **24** (100 mg, 0.30 mmol) in dry DMF (4 mL) at 0 °C. After 15 minutes Boc
5 anhydride (0.60 mmol), diluted in dry DMF (0.5 mL), was added dropwise under vigorous
6 stirring. The resulting solution was warmed to room temperature and heated at 50 °C for 12
7 h. Next, H₂O and ethyl acetate were added and the aqueous phase was extracted twice with
8 ethyl acetate. The combined organic phases were washed with brine, dried over Na₂SO₄ and
9 evaporated. The crude was purified by flash chromatography using DCM/MeOH (98/2) as
10 eluent.
11
12
13
14
15
16
17
18
19

20
21 *Tert-butyl (2-((3-acetylphenyl)amino)-4'-methyl-[4,5'-bithiazol]-2'-yl)carbamate (25g)*.
22 Yield 63%; mp 235-236 °C. MS (ESI) [M + H]⁺: 431.5 *m/z*. ¹H NMR (DMSO-*d*₆ 400 MHz):
23 δ 1.50 (s, 9H), 2.47 (s, 3H), 2.63 (s, 3H), 6.92 (s, 1H), 7.49 (t, 1H, *J* = 7.9 Hz), 7.56 (d, 1H, *J*
24 = 7.9 Hz), 7.80 (d, 1H, *J* = 7.9 Hz), 8.50 (s, 1H), 10.54 (s, 1H), 11.39 (s, 1H). ¹³C NMR
25 (DMSO-*d*₆ 100.6 MHz): δ 17.51, 27.42, 28.37 (3x), 79.61, 103.15, 116.70, 120.03, 121.24,
26 121.65, 128.34, 138.55, 141.83, 143.41, 143.62, 155.67, 157.03, 168.39, 198.32. Anal.
27 (C₂₀H₂₂N₄O₃S₂) C, H, N.
28
29
30
31
32
33
34
35

36 *General Procedure for the Synthesis of Compounds 26a-d*. A solution of intermediate **24**
37 (100 mg, 0.30 mmol) and the proper isocyanate or isothiocyanate (0.45 mmol) in pyridine (2
38 mL) was heated at reflux for 12-18 h. The mixture was cooled to room temperature and
39 diluted with ethyl acetate. The organic phase was washed with saturated aqueous NH₄Cl
40 solution and brine, dried over Na₂SO₄, and concentrated under vacuum. The crude was
41 purified by flash chromatography using DCM/MeOH (98/2) as eluent.
42
43
44
45
46
47
48

49 *1-(2-((3-acetylphenyl)amino)-4'-methyl-[4,5'-bithiazol]-2'-yl)-3-phenylurea (26a)*. Yield
50 59%; mp 236-239 °C. MS (ESI) [M + H]⁺: 450.3 *m/z*. ¹H NMR (DMSO-*d*₆ 400 MHz): δ 2.51
51 (s, 3H), 2.64 (s, 3H), 6.92 (s, 1H), 7.05 (t, 1H, *J* = 7.4 Hz), 7.33 (t, 2H, *J* = 7.4 Hz), 7.48-7.52
52 (m, 3H), 7.57 (d, 1H, *J* = 7.8 Hz), 7.80 (d, 1H, *J* = 7.8 Hz), 8.47 (s, 1H), 8.99 (s, 1H), 10.52
53
54
55
56
57
58
59
60

(s, 1H), 10.55 (s, 1H). ^{13}C NMR (DMSO- d_6 100.6 MHz): δ 17.29, 27.43, 102.86, 116.73, 119.02 (2x), 121.46, 121.62, 123.17, 128.20, 129.39 (2x), 129.83, 138.10, 139.17, 139.43, 141.83, 143.36, 151.90, 155.68, 162.92, 198.35. Anal. ($\text{C}_{22}\text{H}_{19}\text{N}_5\text{O}_2\text{S}_2$) C, H, N.

1-(2-((3-acetylphenyl)amino)-4'-methyl-[4,5'-bithiazol]-2'-yl)-3-phenylthiourea (**26b**).

Yield 65%; mp 217-218 °C. MS (ESI) $[\text{M} + \text{H}]^+$: 466.4 m/z . ^1H NMR (DMSO- d_6 400 MHz): δ 2.54 (s, 3H), 2.64 (s, 3H), 6.99 (s, 1H), 7.07 (t, 1H, $J = 7.4$ Hz), 7.32 (t, 2H, $J = 7.4$ Hz), 7.48 (t, 1H, $J = 7.8$ Hz), 7.57 (d, 1H, $J = 7.8$ Hz), 7.69 (d, 1H, $J = 7.8$ Hz), 7.73 (d, 2H, $J = 7.4$ Hz), 8.57 (s, 1H), 10.20 (s, 1H), 10.59 (s, 1H), 12.68 (s, 1H). ^{13}C NMR (DMSO- d_6 100.6 MHz): δ 17.30, 27.60, 103.92, 116.74, 121.50, 121.77, 122.48, 127.80, 128.20 (2x), 128.84 (2x), 129.81, 138.12, 138.30, 139.89, 141.73, 142.96, 155.60, 163.20, 176.12, 198.28. Anal. ($\text{C}_{22}\text{H}_{19}\text{N}_5\text{OS}_3$) C, H, N.

1-(2-((3-acetylphenyl)amino)-4'-methyl-[4,5'-bithiazol]-2'-yl)-3-cyclohexylurea (**26c**).

Yield 69%; mp 197-198 °C. MS (ESI) $[\text{M} + \text{H}]^+$: 456.4 m/z . ^1H NMR (DMSO- d_6 400 MHz): δ 1.14-1.38 (m, 6H), 1.52-1.56 (m, 1H), 1.64-1.68 (m, 2H), 1.80-1.83 (m, 2H), 2.45 (s, 3H), 2.61 (s, 3H), 6.53 (d, 1H, $J = 7.8$ Hz), 6.84 (s, 1H), 7.48 (t, 1H, $J = 7.8$ Hz), 7.56 (d, 1H, $J = 7.8$ Hz), 7.85 (d, 1H, $J = 7.8$ Hz), 8.41 (s, 1H), 10.09 (s, 1H), 10.52 (s, 1H). ^{13}C NMR (DMSO- d_6 100.6 MHz): δ 17.53, 24.67, 25.58, 27.41 (2x), 33.12 (2x), 48.44, 102.47, 116.67, 120.57, 121.45, 121.57, 129.81, 138.07, 141.85, 143.61, 147.01, 155.38, 157.13, 162.82, 198.29. Anal. ($\text{C}_{22}\text{H}_{25}\text{N}_5\text{O}_2\text{S}_2$) C, H, N.

1-(2-((3-acetylphenyl)amino)-4'-methyl-[4,5'-bithiazol]-2'-yl)-3-cyclohexylthiourea (**26d**).

Yield 52%; mp 234-236 °C. MS (ESI) $[\text{M} + \text{H}]^+$: 472.3 m/z . ^1H NMR (DMSO- d_6 400 MHz): δ 1.24-1.37 (m, 6H), 1.52-1.56 (m, 1H), 1.63-1.67 (m, 2H), 1.85-1.93 (m, 2H), 2.49 (s, 3H), 2.63 (s, 3H), 6.96 (s, 1H), 7.47 (t, 1H, $J = 7.8$ Hz), 7.56 (d, 1H, $J = 7.8$ Hz), 7.85 (d, 1H, $J = 7.8$ Hz), 8.41 (s, 1H), 9.55 (s, 1H), 10.57 (s, 1H), 11.41 (s, 1H). ^{13}C NMR (DMSO- d_6 100.6 MHz): δ 17.84, 24.38, 25.56, 27.50 (2x), 31.80 (2x), 52.62, 103.47, 116.67, 121.45, 121.58,

1
2
3 121.74, 129.79, 137.20, 138.11, 141.65, 142.93, 146.01, 163.00, 171.90, 198.11. Anal.
4
5 (C₂₂H₂₅N₅OS₃) C, H, N.

6
7 *Synthesis of phenyl (5-acetyl-4-methylthiazol-2-yl)carbamate (27).* 1-(2-amino-4-
8 methylthiazol-5-yl)ethanone **20** (1000 mg, 6.40 mmol) was added to a suspension of NaH
9 60% dispersion in mineral oil (768 mg, 19.20 mmol) in DMF (15 mL) at 0 °C. Diphenyl
10 carbonate (3428 mg, 16.0 mmol) was added while cooling and the reaction mixture was
11 stirred for additional 30 minutes at room temperature. H₂O and ethyl acetate were added and
12 the aqueous phase was extracted three times with ethyl acetate. The combined organic phases
13 were washed twice with an aqueous solution of LiCl (5% w/w) and brine, dried over Na₂SO₄
14 and concentrated under vacuum. Ether was added to the crude and the white solid obtained
15 was filtered over a Buchner funnel, washed with ether and used in the following step without
16 any further purification. Yield: 67%. MS (ESI) [M + H]⁺: 277.2 *m/z*. ¹H NMR (DMSO-*d*₆ 400
17 MHz): δ 2.52 (s, 3H), 2.57 (s, 3H), 7.27-7.34 (m, 3H), 7.44-7.48 (m, 2H), 12.71 (s, 1H).
18
19
20
21
22
23
24
25
26
27
28
29
30

31
32 *General Procedure for the Synthesis of Compounds 28a,b.* The proper amine (1.09 mmol)
33 was added to a solution of intermediate **27** (300 mg, 1.09 mmol) in dry THF (15 mL). The
34 mixture was heated at 50 °C for 5-6 h, after which H₂O and ethyl acetate were added and the
35 reaction mixture was cooled down to room temperature. The aqueous phase was extracted
36 twice with ethyl acetate, the combined organic phases were washed with brine, dried over
37 Na₂SO₄ and concentrated under vacuum. The crude was purified by flash chromatography
38 using DCM/MeOH (97/3) as eluent.
39
40
41
42
43
44
45
46

47 *1-(5-acetyl-4-methylthiazol-2-yl)-3-(4-chlorobenzyl)urea (28a).* Yield: 61%. MS (ESI) [M
48 + H]⁺: 324.2 *m/z*. ¹H NMR (DMSO-*d*₆ 300 MHz): δ 2.49 (s, 3H), 2.62 (s, 3H), 4.34 (d, 2H, *J*
49 = 5.7 Hz), 7.18 (bs, 1H), 7.33 (d, 2H, *J* = 8.2 Hz), 7.40 (d, 2H, *J* = 8.2 Hz), 11.02 (s, 1H).
50
51
52
53
54
55
56
57
58
59
60

1
2
3 *1-(5-acetyl-4-methylthiazol-2-yl)-3-(adamantan-1-yl)urea (28b)*. Yield: 93%. MS (ESI) [M
4 + H]⁺: 334.5 *m/z*. ¹H NMR (DMSO-*d*₆ 300 MHz): δ 1.64-1.65 (m, 6H), 1.93-1.95 (m, 6H),
5 2.05-2.06 (m, 3H), 2.49 (s, 3H), 2.63 (s, 3H), 6.42 (s, 1H), 10.21 (s, 1H).
6
7

8
9
10 *General Procedure for the Synthesis of Compounds 29a,b*. A suspension of the proper
11 intermediate **28a,b** (0.62 mmol) in 48% HBr solution in water (2 mL) was warmed to 60 °C.
12 A solution of Br₂ (42 μL, 0.81 mmol) in 1,4-dioxane (2 mL) was added dropwise and the
13 reaction mixture was heated at 60 °C for 3 h. After cooling down to room temperature,
14 saturated aqueous NaHCO₃ solution and ethyl acetate were added and the aqueous phase was
15 extracted three times with ethyl acetate. The combined organic phases were washed with
16 brine, dried over Na₂SO₄ and concentrated under vacuum. Intermediates **29a,b** were used in
17 the next step without any further purification.
18
19
20
21
22
23
24
25
26

27 *1-(5-(2-bromoacetyl)-4-methylthiazol-2-yl)-3-(4-chlorobenzyl)urea (29a)*. Yield: 88%. MS
28 (ESI) [M + H]⁺: 402.4 *m/z*, [M + 2 + H]⁺: 404.3 *m/z*, [M + 4 + H]⁺: 406.3 *m/z*. ¹H NMR
29 (DMSO-*d*₆ 300 MHz): δ 2.49 (s, 3H), 4.34 (d, 2H, *J* = 5.7 Hz), 4.46 (bs, 2H), 7.18 (bs, 1H),
30 7.36 (d, 2H, *J* = 8.2 Hz), 7.43 (d, 2H, *J* = 8.2 Hz), 11.23 (s, 1H).
31
32
33
34
35

36 *1-(adamantan-1-yl)-3-(5-(2-bromoacetyl)-4-methylthiazol-2-yl)urea (29b)*. Yield: 92%.
37 MS (ESI) [M + H]⁺: 412.3 *m/z*, [M + 2 + H]⁺: 414.3 *m/z*. ¹H NMR (DMSO-*d*₆ 300 MHz): δ
38 1.65-1.67 (m, 6H), 1.93-1.95 (m, 6H), 2.06-2.08 (m, 3H), 2.49 (s, 3H), 4.49 (bs, 2H), 6.43 (s,
39 1H), 10.31 (s, 1H).
40
41
42
43
44

45 *General Procedure for the Synthesis of Compounds 30a,b*. A suspension of intermediate **23**
46 (100 mg, 0.51 mmol) and the proper compound **29a,b** (0.51 mmol) in ethanol (6 mL) was
47 heated at reflux for 1 h. After cooling down to room temperature, saturated aqueous NaHCO₃
48 solution, H₂O and ethyl acetate were added and the aqueous phase was extracted three times
49 with ethyl acetate. The combined organic phases were washed with brine, dried over Na₂SO₄
50
51
52
53
54
55
56
57
58
59
60

and concentrated under vacuum. The crude was purified by flash chromatography using DCM/MeOH (97/3) as eluent.

1-(2-((3-acetylphenyl)amino)-4'-methyl-[4,5'-bithiazol]-2'-yl)-3-(4-chlorobenzyl)urea

(30a). Yield 78%; mp 240-241 °C. MS (ESI) $[M + H]^+$: 498.2 *m/z*. ¹H NMR (DMSO-*d*₆ 400 MHz): δ 2.47 (s, 3H), 2.61 (s, 3H), 4.34 (d, 2H, *J* = 5.7 Hz), 6.86 (s, 1H), 7.09 (bs, 1H), 7.33 (d, 2H, *J* = 8.2 Hz), 7.40 (d, 2H, *J* = 8.2 Hz), 7.48 (t, 1H, *J* = 7.8 Hz), 7.55 (d, 1H, *J* = 7.8 Hz), 7.83 (d, 1H, *J* = 7.8 Hz), 8.42 (s, 1H), 10.51 (s, 1H), 10.53 (s, 1H). ¹³C NMR (DMSO-*d*₆ 100.6 MHz): δ 17.50, 27.41, 42.73, 102.57, 116.67, 120.01, 121.45, 121.58, 128.78 (2x), 129.54 (2x), 129.82, 131.89, 138.08, 139.25, 141.84, 143.21, 143.55, 154.41, 157.67, 162.84, 198.30. Anal. (C₂₃H₂₀ClN₅O₂S₂) C, H, N.

1-(2-((3-acetylphenyl)amino)-4'-methyl-[4,5'-bithiazol]-2'-yl)-3-(adamantan-1-yl)urea

(30b). Yield 83%; mp 227-230 °C. MS (ESI) $[M + H]^+$: 508.5 *m/z*. ¹H NMR (DMSO-*d*₆ 400 MHz): δ 1.65-1.67 (m, 6H), 1.93-1.95 (m, 6H), 2.05-2.06 (m, 3H), 2.49 (s, 3H), 2.63 (s, 3H), 6.78 (s, 1H), 6.97 (s, 1H), 7.49 (t, 1H, *J* = 7.8 Hz), 7.56 (d, 1H, *J* = 7.8 Hz), 7.81 (d, 1H, *J* = 7.8 Hz), 8.46 (s, 1H), 10.21 (bs, 1H), 10.59 (s, 1H). ¹³C NMR (DMSO-*d*₆ 100.6 MHz): δ 16.39, 27.47, 29.29 (3x), 36.32 (3x), 41.73 (3x), 51.16, 103.55, 116.76, 119.09, 121.49, 121.63, 129.83, 138.08, 139.61, 141.76, 142.58, 152.13, 158.27, 163.05, 198.32. Anal. (C₂₆H₂₉N₅O₂S₂) C, H, N.

Synthesis of N-(5-acetyl-4-methylthiazol-2-yl)acetamide (31). Intermediate **20** (1000 mg, 6.40 mmol) was suspended in THF/DCM (3/2, 12 mL) and the mixture was cooled down to 0 °C. Pyridine (1.3 mL) was added, followed by the dropwise addition of acetyl chloride (683 μ L, 9.60 mmol). The reaction mixture was stirred at 0 °C for 3 h. Next, H₂O and ethyl acetate were added and the aqueous phase was extracted three times with ethyl acetate. The combined organic phases were washed three times with saturated aqueous NH₄Cl solution and brine, dried over Na₂SO₄ and concentrated under vacuum. Intermediate **31** was used in

1
2
3 the next step without any further purification. Yield: 87%. MS (ESI) $[M + H]^+$: 199.3 m/z . 1H
4 NMR (DMSO- d_6 400 MHz): δ 2.17 (s, 3H), 2.46 (s, 3H), 2.56 (s, 3H), 12.44 (s, 1H).
5
6

7 *Synthesis of N-(5-(2-bromoacetyl)-4-methylthiazol-2-yl)acetamide (32)*. A solution of Br₂
8 (388 μ L, 7.6 mmol) in 1,4-dioxane (8.6 mL) was added dropwise to a stirred solution of
9 intermediate **31** (1200 mg, 6.05 mmol) in 1,4-dioxane (23 mL). The mixture was heated at 50
10 °C for 22 h. After cooling down to room temperature, saturated aqueous NaHCO₃ solution
11 and ethyl acetate were added and the aqueous phase was extracted three times with ethyl
12 acetate. The combined organic phases were washed with brine, dried over Na₂SO₄ and
13 concentrated under vacuum. The crude was purified by flash chromatography using
14 DCM/acetone (95/5) as eluent. Yield 84%. MS (ESI) $[M + H]^+$: 277.3 m/z , $[M + 2 + H]^+$:
15 279.4 m/z . 1H NMR (DMSO- d_6 400 MHz): δ 2.11 (s, 3H), 2.46 (s, 3H), 4.52 (bs, 2H), 12.44
16 (s, 1H).
17
18
19
20
21
22
23
24
25
26
27
28

29 *General Procedure for the Synthesis of Intermediates 34a-d*. Benzoyl isothiocyanate (547
30 μ L, 4.07 mmol) was added dropwise to a solution of the appropriate aniline **33a-d** (3.70
31 mmol) in dry DCM (12 mL) and the mixture was stirred at room temperature for 12 h. The
32 solvent of reaction was evaporated, the solid was dissolved in THF/NaOH 1N (1/1, 15 mL)
33 and the mixture was refluxed for 2 h. Next, H₂O and ethyl acetate were added and the
34 aqueous phase was extracted three times with ethyl acetate. The combined organic phases
35 were dried over Na₂SO₄ and evaporated. Crystallization from ether afforded intermediates
36 **34a-d**.
37
38
39
40
41
42
43
44
45
46

47 *1-phenylthiourea (34a)*. Yield 72%. MS (ESI) $[M + H]^+$: 153.1 m/z . 1H NMR (DMSO- d_6
48 300 MHz): δ 7.09-7.14 (m, 1H), 7.30-7.42 (m, 6H), 9.67 (s, 1H).
49
50

51 *1-(3-hydroxyphenyl)thiourea (34b)*. Yield 74%. MS (ESI) $[M + H]^+$: 169.2 m/z . 1H NMR
52 (DMSO- d_6 300 MHz): δ 6.50-6.54 (m, 1H), 6.74-6.77 (m, 1H), 6.87-6.88 (m, 1H), 7.09 (t,
53 1H, $J = 8.0$ Hz), 7.32-7.36 (bs, 2H), 9.45 (s, 1H), 9.58 (s, 1H).
54
55
56
57
58
59
60

1
2
3 *1-(3-methoxyphenyl)thiourea (34c)*. Yield 75%. MS (ESI) [M + H]⁺: 183.3 *m/z*. ¹H NMR
4 (DMSO-*d*₆ 300 MHz): δ 3.74 (s, 3H), 6.67-6.70 (m, 1H), 6.90-6.94 (m, 1H), 7.11-7.13 (m,
5 1H), 7.22 (t, 1H, *J* = 8.1 Hz), 7.46-7.49 (bs, 2H), 9.73 (s, 1H).
6
7

8
9
10 *1-(4-acetylphenyl)thiourea (34d)*. Yield 78%. MS (ESI) [M + H]⁺: 195.4 *m/z*. ¹H NMR
11 (DMSO-*d*₆ 300 MHz): δ 2.49 (s, 3H), 7.79-7.83 (m, 2H), 7.87-8.03 (m, 4H), 10.69 (s, 1H).
12
13

14 *General Procedure for the Synthesis of Compounds 35a-d*. A solution of intermediate **32**
15 (50 mg, 0.18 mmol) and the proper thiourea **34a-d** (0.18 mmol) in ethanol (2.5 mL) was
16 heated at reflux for 1 h. After cooling down to room temperature, saturated aqueous NaHCO₃
17 solution, H₂O and ethyl acetate were added and the aqueous phase was extracted three times
18 with ethyl acetate. The combined organic phases were washed with brine, dried over Na₂SO₄
19 and concentrated under vacuum. The crude was purified by flash chromatography using
20 DCM/MeOH (97/3) as eluent.
21
22
23
24
25
26
27
28

29
30 *N-(4'-methyl-2-(phenylamino)-[4,5'-bithiazol]-2'-yl)acetamide (35a)*. Yield 85%; mp 187-
31 189 °C. MS (ESI) [M + H]⁺: 331.1 *m/z*. ¹H NMR (CDCl₃ 400 MHz): δ 2.19 (s, 3H), 2.57 (s,
32 3H), 6.58 (s, 1H), 7.10-7.12 (m, 1H), 7.36-7.42 (m, 4H), 8.18 (s, 1H), 11.94 (s, 1H). ¹³C
33 NMR (CDCl₃ 100.6 MHz): δ 17.00, 23.07, 102.63, 118.50 (2x), 121.29, 123.32, 129.54 (2x),
34 140.15, 142.70, 143.31, 156.66, 164.76, 167.92. Anal. (C₁₅H₁₄N₄OS₂) C, H, N.
35
36
37
38
39

40
41 *N-(2-((3-hydroxyphenyl)amino)-4'-methyl-[4,5'-bithiazol]-2'-yl)acetamide (35b)*. Yield
42 71%; mp 148-149 °C. MS (ESI) [M + H]⁺: 347.2 *m/z*. ¹H NMR (acetone-*d*₆ 400 MHz): δ
43 2.26 (s, 3H), 2.51 (s, 3H), 6.52-6.54 (m, 1H), 6.78 (s, 1H), 7.15-7.18 (m, 2H), 7.27 (s, 1H),
44 8.45 (s, 1H), 9.30 (s, 1H), 10.85 (s, 1H). ¹³C NMR (acetone-*d*₆ 100.6 MHz): δ 16.51, 21.90,
45 101.85, 104.58, 108.75, 109.12, 120.91, 129.81, 142.39, 143.14, 143.95, 155.28, 158.21,
46 163.25, 167.72. Anal. (C₁₅H₁₄N₄O₂S₂) C, H, N.
47
48
49
50
51

52
53
54 *N-(2-((3-methoxyphenyl)amino)-4'-methyl-[4,5'-bithiazol]-2'-yl)acetamide (35c)*. Yield
55 74%; mp 118-119 °C. MS (ESI) [M + H]⁺: 361.2 *m/z*. ¹H NMR (DMSO-*d*₆ 400 MHz): δ 2.14
56
57
58
59
60

(s, 3H), 2.48 (s, 3H), 3.82 (s, 3H), 6.53 (d, 1H, $J = 8.2$ Hz), 6.90 (s, 1H), 6.98 (s, 1H, $J = 8.2$ Hz), 7.20 (t, 1H, $J = 8.2$ Hz), 7.71 (s, 1H), 10.42 (s, 1H), 12.06 (s, 1H). ^{13}C NMR (DMSO- d_6 100.6 MHz): δ 17.53, 22.93, 55.52, 102.71, 105.60, 107.78, 109.65, 120.81, 130.02, 142.77, 142.90, 143.12, 155.59, 160.40, 162.96, 168.75. Anal. ($\text{C}_{16}\text{H}_{16}\text{N}_4\text{O}_2\text{S}_2$) C, H, N.

N-(2-((4-acetylphenyl)amino)-4'-methyl-[4,5'-bithiazol]-2'-yl)acetamide (**35d**). Yield 79%; mp 233-235 °C. MS (ESI) $[\text{M} + \text{H}]^+$: 373.2 m/z . ^1H NMR (DMSO- d_6 400 MHz): δ 2.15 (s, 3H), 2.51 (s, 3H), 2.54 (s, 3H), 7.07 (s, 1H), 7.77 (d, 2H, $J = 8.7$ Hz), 7.97 (d, 2H, $J = 8.7$ Hz), 10.79 (s, 1H), 12.11 (s, 1H). ^{13}C NMR (DMSO- d_6 100.6 MHz): δ 17.49, 22.93, 26.82, 104.38, 116.33 (2x), 120.33, 130.34, 130.42 (2x), 143.41, 145.48, 152.60, 155.63, 162.37, 168.83, 196.63. Anal. ($\text{C}_{17}\text{H}_{16}\text{N}_4\text{O}_2\text{S}_2$) C, H, N.

Synthesis of 1-(3-acetylphenyl)guanidine (36). Nitric acid (164 μL , 3.70 mmol) was added to a solution of 3'-aminoacetophenone **22** (500 mg, 3.70 mmol) in ethanol (10 mL), followed by addition of a solution of cyanamide (778 mg, 18.5 mmol) in a minimal amount of water. The mixture was heated at reflux for 24 h and concentrated in vacuum. After cooling to 0 °C, ether was added and the precipitate was separated by filtration over a Buchner funnel. Then saturated aqueous NaHCO_3 solution and ethyl acetate were added to the solid and the aqueous phase was extracted three times with ethyl acetate. The combined organic phases were washed with brine, dried over Na_2SO_4 and concentrated under vacuum to obtain compound **36**, used in the next step without any further purification. Yield 73%. MS (ESI) $[\text{M} + \text{H}]^+$: 178.2 m/z . ^1H NMR (DMSO- d_6 300 MHz): δ 2.57 (s, 3H), 7.49 (bs, 3H), 7.58-7.60 (m, 2H), 7.76-7.78 (m, 1H), 7.88 (s, 1H), 9.67 (bs, 1H).

Synthesis of 1-(3-((4-(2-amino-4-methylthiazol-5-yl)-1H-imidazol-2-yl)amino)phenyl)ethanone (37). A solution of intermediate **36** (150 mg, 0.85 mmol) in ethanol (5 mL) was added dropwise to a solution of compound **21** (200 mg, 0.85 mmol) and Et_3N (118 μL , 0.85 mmol) in ethanol (10 mL). The mixture was heated at reflux for 12 h,

1
2
3 after which H₂O and ethyl acetate were added. The organic phase was washed with saturated
4
5 aqueous NH₄Cl solution and brine, dried over Na₂SO₄ and concentrated under vacuum to
6
7 obtain compound **37**, used in the next step without any further purification. Yield 82%. MS
8
9 (ESI) [M + H]⁺: 314.3 *m/z*. ¹H NMR (DMSO-*d*₆ 400 MHz): δ 2.37 (s, 3H), 2.64 (s, 3 H), 7.09
10
11 (s, 1 H), 7.65 (t, 1H, *J* = 7.9 Hz), 7.79 (d, 1H, *J* = 7.9 Hz), 7.94 (d, 1H, *J* = 7.9 Hz), 8.04 (s,
12
13 1H), 9.50 (bs, 2H), 10.56 (s, 1H), 12.01 (s, 1H).
14
15

16
17 *Synthesis of N-(5-(2-((3-acetylphenyl)amino)-1H-imidazol-4-yl)-4-methylthiazol-2-*
18
19 *yl)acetamide (38)*. Et₃N (89 μL, 0.64 mmol) was added to a stirred suspension of
20
21 intermediate **37** (100 mg, 0.32 mmol) in dry DCM (4.5 mL) at 0 °C. After 15 minutes acetyl
22
23 chloride (34 μL, 0.48 mmol), diluted in dry DCM (0.5 mL), was added dropwise. The
24
25 resulting solution was warmed to room temperature and stirred for 8 h. Next, H₂O and DCM
26
27 were added and the aqueous phase was extracted three times with DCM. The combined
28
29 organic phases were washed with brine, dried over Na₂SO₄ and evaporated. The crude was
30
31 purified by flash chromatography using DCM/MeOH (97/3) as eluent. Yield 47%; mp 222-
32
33 225 °C. MS (ESI) [M + H]⁺: 356.2 *m/z*. ¹H NMR (DMSO-*d*₆ 400 MHz): δ 2.14 (s, 3H), 2.37
34
35 (s, 3H), 2.65 (s, 3H), 7.11 (s, 1H), 7.66 (t, 1H, *J* = 7.9 Hz), 7.79 (d, 1H, *J* = 7.9 Hz), 7.94 (d,
36
37 1H, *J* = 7.9 Hz), 8.05 (s, 1H), 10.56 (s, 1H), 11.84 (s, 1H), 12.06 (s, 1H). ¹³C NMR (DMSO-
38
39 *d*₆ 100.6 MHz): δ 17.56, 22.56, 27.42, 116.75, 119.9, 120.51, 121.65, 127.71, 129.79, 138.10,
40
41 140.81, 143.21, 143.28, 154.98, 162.89, 168.63, 198.32. Anal. (C₁₇H₁₇N₅O₂S) C, H, N.
42
43
44

45
46 *Synthesis of 1-(3-acetylphenyl)-3-(3-nitrophenyl)thiourea (40)*. 3'-aminoacetophenone **22**
47
48 (300 mg, 2.22 mmol) was added to a solution of 3-nitrophenyl isothiocyanate (400 mg, 2.22
49
50 mmol) in dry DCM (6.50 mL). The solution was stirred at room temperature for 18 h. The
51
52 precipitate was separated by filtration over a Buchner funnel and washed with ether,
53
54 affording compound **40** as a white solid. Yield 88%. MS (ESI) [M + H]⁺: 316.2 *m/z*. ¹H NMR
55
56 (DMSO-*d*₆ 400 MHz): δ 2.58 (s, 3H), 7.52 (t, 1H, *J* = 7.9 Hz), 7.63 (t, 1H, *J* = 8.1 Hz), 7.76-
57
58
59
60

1
2
3 7.78 (m, 2H), 7.92 (d, 1H, $J = 7.9$ Hz), 7.98 (d, 1H, $J = 8.1$ Hz), 8.07 (s, 1H), 8.56 (s, 1H),
4
5 10.31 (bs, 1H), 10.33 (bs, 1H).
6

7 *Synthesis of 1-(3-acetylphenyl)-3-(3-aminophenyl)thiourea (41)*. Iron powder (1490 mg,
8
9 26.67 mmol), water (7 mL) and concentrated HCl (4 drops) were added to a solution of
10
11 compound **40** (400 mg, 1.27 mmol) in ethanol (35 mL). After heating at reflux for 2 h, the
12
13 mixture was filtrated hot, washed with ethanol and concentrated in vacuum. The crude was
14
15 purified by flash chromatography using DCM/MeOH (98/2) as eluent. Yield 75%. MS (ESI)
16
17 $[M + H]^+$: 286.1 m/z . 1H NMR ($CDCl_3$, 400 MHz): δ 2.58 (s, 3H), 3.20-3.51 (bs, 2H), 6.59-
18
19 6.62 (m, 2H), 6.68 (d, 1H, $J = 7.7$ Hz), 7.18 (t, 1H, $J = 7.8$ Hz), 7.45 (t, 1H, $J = 7.8$ Hz),
20
21 7.74-7.79 (m, 2H), 7.95 (s, 1H), 8.05 (s, 1H), 8.38 (s, 1H).
22
23
24

25 *General Procedure for the Synthesis of Compounds 42a,b*. The proper acyl chloride (0.53
26
27 mmol) was added to a stirred solution of intermediate **41** (100 mg, 0.35 mmol) and pyridine
28
29 (56 μ L, 0.70 mmol) in dry THF (4 mL). The resulting solution was stirred at room
30
31 temperature for 2 h, after which H_2O and ethyl acetate were added and the aqueous phase was
32
33 extracted three times with ethyl acetate. The combined organic phases were washed with
34
35 brine, dried over Na_2SO_4 and concentrated under vacuum. The crude was purified by flash
36
37 chromatography using DCM/MeOH (99/1) as eluent.
38
39

40
41 *N-(3-(3-(3-acetylphenyl)thioureido)phenyl)acetamide (42a)*. Yield 67%; mp 171-173 $^{\circ}C$.
42
43 MS (ESI) $[M + H]^+$: 328.1 m/z . 1H NMR ($DMSO-d_6$, 400 MHz): δ 2.04 (s, 3H), 2.57 (s, 3H),
44
45 7.16-7.19 (m, 1H), 7.25 (t, 1H, $J = 7.9$ Hz), 7.35-7.37 (m, 1H), 7.47 (t, 1H, $J = 7.9$ Hz), 7.71-
46
47 7.79 (m, 3H), 8.07 (bs, 1H), 9.89 (s, 1H), 9.96 (s, 1H), 9.98 (s, 1H). ^{13}C NMR ($DMSO-d_6$
48
49 100.6 MHz): 23.43, 27.23, 116.20, 117.10, 119.38, 123.64, 124.55, 128.90, 128.98, 129.04,
50
51 136.97, 139.64, 140.17, 140.52, 168.56, 180.27, 197.97. Anal. ($C_{17}H_{17}N_3O_2S$) C, H, N.
52
53

54 *N-(3-(3-(3-acetylphenyl)thioureido)phenyl)pivalamide (42b)*. Yield 69%; mp 157-159 $^{\circ}C$.
55
56 MS (ESI) $[M + H]^+$: 370.3 m/z . 1H NMR ($DMSO-d_6$, 300 MHz): δ 1.22 (s, 9H), 2.57 (s, 3H),
57
58
59
60

1
2
3 7.14 (d, 1H, $J = 7.9$ Hz), 7.25 (t, 1H, $J = 7.9$ Hz), 7.42-7.50 (m, 2H), 7.71-7.80 (m, 3H), 8.06
4 (bs, 1H), 9.25 (s, 1H), 9.89 (s, 1H), 9.94 (s, 1H). ^{13}C NMR (DMSO- d_6 100.6 MHz): δ 27.23,
5 27.65 (3x), 30.90, 116.29, 117.14, 119.35, 123.68, 124.71, 128.88, 128.92, 129.10, 137.43,
6 139.65, 140.20, 140.52, 176.96, 180.27, 197.97. Anal. ($\text{C}_{20}\text{H}_{23}\text{N}_3\text{O}_2\text{S}$) C, H, N.

7
8
9
10
11 *Synthesis of 1-(4-(dimethylamino)phenyl)thiourea (44).* Benzoyl isothiocyanate (543 μL ,
12 4.04 mmol) was added dropwise to a solution of N,N-Dimethyl-p-phenylenediamine **43** (500
13 mg, 3.67 mmol) in dry DCM (12 mL) and the mixture was stirred at room temperature for 12
14 h. The solvent of reaction was evaporated, the solid was dissolved in THF/NaOH 1N (1/1, 15
15 mL) and the mixture was refluxed for 3 h. Next, H_2O and ethyl acetate were added and the
16 aqueous phase was extracted three times with ethyl acetate. The combined organic phases
17 were dried over Na_2SO_4 and evaporated. The resulting solid was crystallized from ether.
18 Yield 80%. MS (ESI) $[\text{M} + \text{H}]^+$: 196.4 m/z . ^1H NMR (DMSO- d_6 300 MHz): δ 2.88 (s, 6H),
19 6.69 (d, 2H, $J = 8.7$ Hz), 7.08 (d, 2H, $J = 8.7$ Hz), 7.55 (bs, 2H), 9.58 (s, 1H).
20
21
22
23
24
25
26
27
28
29
30
31

32 *Synthesis of N^2 -(4-(dimethylamino)phenyl)-4'-methyl-[4,5'-bithiazole]-2,2'-diamine (45).*
33 Intermediates **21** (150 mg, 0.64 mmol) and **44** (125 mg, 0.64 mmol) were suspended in
34 ethanol (6 mL) and the mixture was heated at reflux for 30 minutes. Then saturated aqueous
35 NaHCO_3 solution and ethyl acetate were added to the mixture and the aqueous phase was
36 extracted three times with ethyl acetate. The combined organic phases were washed with
37 brine, dried over Na_2SO_4 and concentrated under vacuum. Ether was added to the crude and
38 the solid obtained was filtered over a Buchner funnel, washed with ether and used in the
39 following step without any further purification. Yield 77%. MS (ESI) $[\text{M} + \text{H}]^+$: 332.2 m/z .
40 ^1H NMR (DMSO- d_6 400 MHz): δ 2.47 (s, 3H), 2.86 (s, 6H), 6.72 (s, 1H), 6.76 (d, 2H, $J = 8.8$
41 Hz), 7.43 (d, 2H, $J = 8.8$ Hz), 9.21 (bs, 2H), 10.58 (s, 1H).
42
43
44
45
46
47
48
49
50
51
52
53

54 *N-(2-((4-(dimethylamino)phenyl)amino)-4'-methyl-[4,5'-bithiazol]-2'-yl)propionamide*
55 (**46**). Et_3N (84 μL , 0.60 mmol) was added to a stirred suspension of intermediate **45** (100 mg,
56
57
58
59
60

0.30 mmol) in dry DCM (4 mL) at 0 °C. After 15 minutes propionyl chloride (39 μ L, 0.45 mmol), diluted in dry DCM (0.5 mL), was added dropwise. The resulting solution was warmed to room temperature and stirred for 8 h. Next, H₂O and DCM were added and the aqueous phase was extracted twice with DCM. The combined organic phases were washed with brine, dried over Na₂SO₄ and evaporated. The crude was purified by flash chromatography using DCM/MeOH (98/2) as eluent. Yield 68%; mp 222-224 °C. MS (ESI) [M + H]⁺: 388.3 *m/z*. ¹H NMR (DMSO-*d*₆ 400 MHz): δ 1.10 (t, 3H, *J* = 7.6 Hz), 2.43 (q, 2H, *J* = 7.6 Hz), 2.47 (s, 3H), 2.86 (s, 6H), 6.73 (s, 1H), 6.76 (d, 2H, *J* = 8.8 Hz), 7.43 (d, 2H, *J* = 8.8 Hz), 9.90 (s, 1H), 11.98 (s, 1H). ¹³C NMR (DMSO-*d*₆ 100.6 MHz): δ 8.78, 16.88, 30.43, 40.46 (2x), 101.02, 113.74 (2x), 120.67 (2x), 121.44, 131.54, 143.10, 144.02, 147.72, 155.12, 165.29, 171.76. Anal. (C₁₈H₂₁N₅OS₂) C, H, N.

Biology

Antiviral assays – materials and methods Assay preparation

Enterovirus (EV):

Rhabdomyosarcoma (RD) cells, Vero cells and Hela-Rh cells, subcultured in cell growth medium [MEM Rega3 (Cat. N°19993013; Invitrogen) supplemented with 10% FCS (Integro), 5 ml 200 mM L-glutamine (25030024) and 5 mL 7.5% sodium bicarbonate (25080060)] at a ratio of 1:4 and grown for 7 days in 150 cm² tissue culture flasks (Techno Plastic Products), were harvested and seeded in a 96-well plate at a cell density of 20 000 cells/well in assay medium (MEM Rega3, 2% FCS, 5 ml L-glutamine and 5 ml sodium bicarbonate) to perform standardized antiviral assay against EV71 and EVD68, CV and PV, RV02 and RV14, respectively.

Antiviral activity and cytotoxicity determinations

1
2
3 Compounds were prepared as DMSO stock solution with a final compound concentration of
4
5 10mM. The compound profiling setup was performed employing a Freedom EVO200 liquid
6
7 handling platform (Tecan). The evaluation of the cytostatic/cytotoxic as well as the antiviral
8
9 effect of each compound was performed in parallel within one run. Three 8-step 1-to-5
10
11 dilution series were prepared (starting from 100 μ M) in assay medium added to empty wells
12
13 (picornaviruses: 96-well microtiter plates, Falcon, BD) or in the medium present on top of
14
15 pre-seeded cells. Subsequently, 50 μ L of a 4x virus dilution in assay medium (assay medium
16
17 supplemented with 15 ml $MgCl_2$ 1M (Sigma, M1028) in case of RV) was added followed by
18
19 50 μ L of cell suspension. The assay plates were returned to the incubator for 2-3
20
21 (picornavirus, 35°C for RV) days, a time at which maximal cytopathic effect (CPE) for
22
23 picornaviruses is observed.
24
25
26

27
28 For the evaluation of cytostatic/cytotoxic effects and for the evaluation of the antiviral effect
29
30 in case of PV, CV, RV, the assay medium was replaced with 75 μ L of a 5% MTS (Promega)
31
32 solution in phenol red-free medium and incubated for 1.5 hours (37°C, 5% CO_2 , 95-99%
33
34 relative humidity). Absorbance was measured at a wavelength of 498 nm (Safire2, Tecan)
35
36 and optical densities (OD values) were converted to percentage of untreated controls.
37
38

39
40 Analysis of the raw data, quality control of each individual dose-response curve and
41
42 calculation, if possible, of the EC_{50} , EC_{90} and CC_{50} values was performed employing
43
44 ViroDM, a custom-made data processing software package. The EC_{50} and EC_{90} (values
45
46 derived from the dose-response curve) represent the concentrations at which respectively
47
48 50% and 90% inhibition of viral replication would be observed. The CC_{50} (value derived
49
50 from the dose-response curve) represents the concentration at which the metabolic activity of
51
52 the cells would be reduced to 50 % of the metabolic activity of untreated cells.
53
54
55
56
57
58
59
60

1
2
3 The EC_{50} , EC_{90} and $CC_{50} \pm SD$ were, whenever possible, calculated respectively as the
4
5 median of all the EC_{50} , EC_{90} or CC_{50} values derived from the 3 individual dose-response
6
7 curves. The selectivity index (SI), indicative of the therapeutic window of the compound, was
8
9 calculated as CC_{50}/EC_{50} . No further statistical analysis was performed.
10
11

12 13 14 **CFTR assays**

15
16
17 The effects of compounds on CFTR biogenesis and function were measured using
18
19 newly developed assays exploiting CFTR fusion probes with anion-sensitive YFP⁶¹ and pH-
20
21 sensitive pHTomato.⁶² Lipofectamine transfection was used for transient transfection of
22
23 HEK293 cells. Cells plated in 96-well plates were incubated with the YFP-CFTR- or CFTR-
24
25 pHTomato- encoding plasmid⁴⁸ using Lipofectamine 2000 (Life Technologies), according to
26
27 manufacturer's instructions. Following transfection, cell plates were returned to the 37°C
28
29 incubator for 24 h. Plates were further incubated at 30°C for 24 h prior to imaging, with or
30
31 without additional drug treatment.
32
33

34
35
36 All imaging was carried out using ImageXpress (ImageXpress Micro XLS, Molecular
37
38 Devices); an image-acquisition system equipped with wide-field inverted fluorescence
39
40 microscope and fluidics robotics. Images were obtained with a 20X objective, using
41
42 excitation/emission filters 472 ± 30 nm and 520 ± 35 nm, for YFP-CFTR and 531 ± 20 nm
43
44 and 592 ± 20 nm for CFTR-pHTomato. In the latter assay, eGFP and Hoechst nuclear stain
45
46 images were also acquired for each well, using excitation/emission filters 472 ± 30 nm and
47
48 520 ± 35 nm, and 377 ± 25 nm and 447 ± 30 nm, respectively. For each plate, the laser
49
50 intensity and exposure were optimized to achieve the highest possible fluorescence whilst
51
52 avoiding both photobleaching and saturation (illumination intensity 100-150/225 cd, and
53
54 exposure 0.1 – 0.2 s)
55
56
57
58
59
60

1
2
3 For the YFP-CFTR assays, before imaging, cells were washed twice with 100 μ l standard
4 buffer (140 mM NaCl, 4.7 mM KCl, 1.2 mM MgCl₂, 5 mM HEPES, 2.5 mM CaCl₂, 1 mM
5 Glucose, pH 7.4). Images were taken for 150 s at a frequency of 0.5 Hz. 50 μ L extracellular
6 Γ (as standard buffer with 140 mM NaCl replaced with 300 mM NaI; resulting in 100 mM
7 final [Γ]) was added at 20 s, and activating compounds (50 μ M Forskolin alone or together
8 with 10 μ M compounds for acute treatment) were added at 60 s.
9

10
11
12 For the CFTR-pHTomato assay, before imaging, cells were washed twice with 100 μ L
13 standard buffer (as above). During imaging, extracellular pH was changed using addition of
14 50 μ L pH 6 buffer (as standard buffer, with 5 mM HEPES replaced with 10 mM MES: final
15 [MES] 3.3 mM, \sim pH 6.5), and 50 μ L pH 9 buffer (as standard buffer, with 5 mM HEPES
16 replaced with 100 mM Tris: final [Tris] 25 mM, \sim pH 8.8). Two pHTomato images
17 (acquisition frequency 0.5 Hz) were taken in each condition. To account for variation in
18 transfection efficiency the pHTomato fluorescence was normalized using average
19 fluorescence intensity of a soluble eGFP, co-expressed in the cytosol. Because the rise in
20 pHTomato fluorescence falls largely within the 6.5 to 8.8 pH range,⁶³ the change in
21 fluorescence obtained upon increasing extracellular pH ($\Delta F_{\text{membrane}}$) was used as an estimate
22 of membrane-exposed CFTR.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 **In vitro kinase inhibition assays**

46
47 Recombinant full length, HIS6-tagged PI4KIII β was purchased from ProQinase (Germany);
48 recombinant full length, GST tagged PI4KIII α was from Life Technologies. Recombinant
49 full length, HIS6-tagged (PI3K- α) and full length, myc-tagged (p85 α) PI3K- α /p85 α was
50 purchased from ProQinase (Germany).
51
52
53
54
55

56
57 Assay conditions:
58
59
60

1
2
3 PI4KIII β , PI4KIII α and PI3K- α /p85 α reactions were performed in 10 μ L using 20 mM Tris-
4 HCl pH 7.5, 0.125 mM EGTA, 2 mM DTT, 0.04% Triton, 3 mM MgCl₂, 3 mM MnCl₂, 20
5 μ M ATP, 0.01 μ Ci γ -P33 ATP, 200 μ M Pi:3PS, 10% DMSO, 0.4 ng/ μ L of PI4KIII β , 16
6 ng/ μ L of PI4KIII α and 7.6 ng/ μ L of PI3K- α /p85 α . All reactions were performed at 30 °C for
7 10 min. Reactions were stopped by adding 5 μ L of phosphoric acid 0.8%. Aliquots (10 μ L)
8 were then transferred into a P30 Filtermat (PerkinElmer), washed five times with 0.5 %
9 phosphoric acid and four times with water for 5 min. The filter was dried and transferred to a
10 sealable plastic bag, and scintillation cocktail (4 mL) was added. Spotted reactions were read
11 in a scintillation counter (Trilux, Perkinelmer). IC₅₀ values were obtained according to
12 Equation (1), where v is the measured reaction velocity, V is the apparent maximal velocity
13 in the absence of inhibitor, I is the inhibitor concentration, and IC₅₀ is the 50% inhibitory
14 concentration.

$$v = V / \{1 + (I / IC_{50})\} \quad (1)$$

Lipidic substrate preparation:

35
36 PI: phosphatidylinositol (Sigma); PS: 2-Oleoyl-1-palmitoyl-sn-glycero-3-phospho-L-serine
37 (Sigma). PI and PS were dissolved in chloroform/methanol 9:1 and mixed at a 1:3 ratio. After
38 chloroform/methanol evaporation, water was added to 1:62.5 w/v and the mixture sonicated
39 to clarity.
40
41
42
43
44

Kinase Panel:

45
46 All Tyrosine- and Serine/Threonine kinase reactions were performed according to
47 manufacturer's instructions, using 10-50 ng of enzyme. Details on the nature of the substrates
48 and their concentration are reported elsewhere.⁶³ For some kinases, NP-40 or BSA was
49 added. All reactions were performed in 10 μ L at 30 °C for 10 min using protein low-binding
50
51
52
53
54
55
56
57
58
59
60

1
2
3 tubes. Reactions were stopped, transferred to filter and counted as reported in Ref. 59.
4
5 PI4KIII β was purchased from Proqinase. Reactions were performed according to the
6
7 manufacturer's instructions and detected using ADP-Glo™ Lipid Kinase Assay (Promega).
8
9

10 **Supporting Information.** 3D coordinates of the PI4KIII β and PI4KIII α homology models.
11
12 Molecular formula strings. This material is available free of charge via the Internet at
13
14 <http://pubs.acs.org>.
15
16

17 18 AUTHOR INFORMATION

19 20 21 **Corresponding Author**

22
23 *Prof. Marco Radi, P4T Group, Dipartimento di Scienze degli Alimenti e del Farmaco,
24
25 Università degli Studi di Parma, Viale delle Scienze, 27/A, 43124 Parma, Italy. E-mail:
26
27 marco.radi@unipr.it; phone: +39 0521 906080; fax: +39 0521 905006
28
29

30
31 *Prof. Johan Neyts, Laboratory of Virology and Experimental Chemotherapy, Rega
32
33 Institute for Medical Research, KU Leuven, Minderbroedersstraat 10, 3000, Leuven,
34
35 Belgium. E-mail: johan.neyts@kuleuven.be; phone: +32 163 21893; fax: +32 163 37340
36
37

38 39 **Author Contributions**

40
41 The manuscript was written through contributions of all authors. All authors have given
42
43 approval to the final version of the manuscript.
44
45

46 47 ACKNOWLEDGMENT

48
49 Work in the P4T group has been supported by the Chiesi Foundation (Bando Dottorati di
50
51 Ricerca 2014) and by the University of Parma (to MR). Work in the IGM laboratory has been
52
53 supported by the Italian Cancer Research Association (AIRC), Grant IG15868. EL's work
54
55 was supported by the Cystic Fibrosis Trust, (CFT Project No RS31). SL is funded by the
56
57
58
59
60

1
2
3 China Scholarship Council (CSC) (Grant 201403250056) and KL is supported by the
4
5 European Union as a Marie-Sklodowska Curie European Training Network ANTIVIRALS
6
7 (GA 642434).
8
9

10 ABBREVIATIONS

11
12 CF, cystic fibrosis; CFTR, cystic fibrosis transmembrane conductance regulator; COPD,
13
14 chronic obstructive pulmonary disorder; CPE, cytopathic effect; CVB3, coxsackievirus B3;
15
16 DC-SIGN, dendritic cell-specific intercellular adhesion molecule-3 grabbing nonintegrin;
17
18 ECHO11, Echovirus 11; EV, enterovirus; F508del, deletion of Phe 508; RV, human
19
20 rhinovirus; HTD, high-throughput docking; PI, phosphatidylinositol; PI4K,
21
22 phosphatidylinositol 4-kinase; PIP₂, phosphatidylinositol 4,5-bisphosphate; PIP₃,
23
24 phosphatidylinositol 3,4,5-trisphosphate; PI4P, phosphatidylinositol 4-phosphate; PV1,
25
26 poliovirus 1; TGN, trans-Golgi network.
27
28
29
30

31 REFERENCES

32
33 (1) Adams, M. J.; King, A. M. Q.; Carstens, E. B. Ratification Vote on Taxonomic
34
35 Proposals to the International Committee on Taxonomy of Viruses (2013). *Arch. Virol.* **2013**,
36
37 *158*, 2023–2030.
38
39

40
41 (2) Tapparel, C.; Siegrist, F.; Petty, T. J.; Kaiser, L. Picornavirus and Enterovirus Diversity
42
43 with Associated Human Diseases. *Infect. Genet. Evol.* **2013**, *14*, 282-293.
44
45

46
47 (3) Sabanathan, S.; Tan, L. V.; Thwaites, L.; Wills, B.; Qui, P. T.; Rogier van Doorn, H.
48
49 Enterovirus 71 Related Severe Hand, Foot and Mouth Disease Outbreaks in South-East Asia:
50
51 Current Situation and Ongoing Challenges. *J. Epidemiol. Community Health* **2014**, *68*, 500-
52
53 502.
54

55
56 (4) Esposito, S.; Bosis, S.; Niesters, H.; Principi, N. Enterovirus D68 Infection. *Viruses*
57
58 **2015**, *7*, 6043-6050.
59
60

1
2
3 (5) Leigh, R.; Proud, D. Virus-Induced Modulation of Lower Airway Diseases:
4 Pathogenesis and Pharmacologic Approaches to Treatment. *Pharmacol. Ther.* **2015**, *148*,
5 185-198.
6
7

8
9
10 (6) Wat, D. Impact of Respiratory Viral Infections on Cystic Fibrosis. *Postgrad. Med. J.*
11 **2003**, *79*, 201-203.
12
13

14
15 (7) (a) Waters, V.; Ratjen, F. Pulmonary Exacerbations in Children with Cystic Fibrosis.
16 *Ann. Am. Thorac. Soc.* **2015**, *12* (Suppl. 2), S200-S206. (b) Singanayagam, A.; Joshi, P. V.;
17 Mallia, P.; Johnston, S. L. Viruses Exacerbating Chronic Pulmonary Disease: The Role of
18 Immune Modulation. *BMC Med.* **2012**, *10*, 27. (c) Goffard, A.; Lambert, V.; Salleron, J.;
19 Herwegh, S.; Engelmann, I.; Pinel, C.; Pin, I.; Perrez, T.; Prévotat, A.; Dewilde, A.; Delhaes,
20 L. Virus and Cystic Fibrosis: Rhinoviruses Are Associated with Exacerbations in Adult
21 Patients. *J. Clin. Virol.* **2014**, *60*, 147-153. (d) Esther, C. R.; Lin, F. C.; Kerr, A.; Miller, M.
22 B.; Gilligan, P. H. Respiratory Viruses Are Associated with Common Respiratory Pathogens
23 in Cystic Fibrosis. *Pediatr. Pulmonol.* **2014**, *49*, 926-931.
24
25
26
27
28
29
30
31
32
33
34

35
36 (8) van Ewijk, B. E.; van der Zalm, M. M.; Wolfs, T. F. W.; Fleer, A.; Kimpen, J. L. L.;
37 Wilbrink, B.; van der Ent, C. K. Prevalence and Impact of Respiratory Viral Infections in
38 Young Children With Cystic Fibrosis: Prospective Cohort Study. *Pediatrics* **2008**, *122*, 1171-
39 1176.
40
41
42
43
44

45
46 (9) Debing, Y.; Neyts, J.; Delang, L. The Future of Antivirals: Broad-Spectrum Inhibitors.
47 *Curr. Opin. Infect. Dis.* **2015**, *28*, 596-602.
48
49

50
51 (10) de Chasse, B.; Meyniel-Schicklin, L.; Aublin-Gex, A.; André, P.; Lotteau, V. New
52 Horizons for Antiviral Drug Discovery from Virus-Host Protein Interaction Networks. *Curr.*
53 *Opin. Virol.* **2012**, *2*, 606-613.
54
55
56
57
58
59
60

1
2
3 (11) Delang, L.; Paeshuyse, J.; Neyts, J. The Role of Phosphatidylinositol 4-Kinases and
4 Phosphatidylinositol 4-Phosphate during Viral Replication. *Biochem. Pharmacol.* **2012**, *84*,
5 1400-1408.
6
7

8
9
10 (12) Hsu, N. Y.; Ilnytska, O.; Belov, G.; Santiana, M.; Chen, Y. H.; Takvorian, P. M.; Pau,
11 C.; van der Schaar, H.; Kaushik-Basu, N.; Balla, T.; Cameron, C. E.; Ehrenfeld, E.; van
12 Kuppeveld, F. J. M.; Altan-Bonnet, N. Viral Reorganization of the Secretory Pathway
13 Generates Distinct Organelles for RNA Replication. *Cell* **2010**, *141*, 799-811.
14
15

16
17 (13) Arita, M.; Kojima, H.; Nagano, T.; Okabe, T.; Wakita, T.; Shimizu, H.
18 Phosphatidylinositol 4-kinase III Beta Is a Target of Enviroxime-Like Compounds for
19 Antipoliiovirus Activity. *J. Virol.* **2011**, *85*, 2364-2372.
20
21

22 (14) van der Schaar, H. M.; Leyssen, P.; Thibaut, H. J.; de Palma, A.; van der Linden, L.;
23 Lanke, K. H. W.; Lacroix, C.; Verbeken, E.; Conrath, K.; MacLeod, A. M.; Mitchell, D. R.;
24 Palmer, N. J.; van de Poël, H.; Andrews, M.; Neyts, J.; van Kuppeveld, F. J. M. A Novel,
25 Broad-Spectrum Inhibitor of Enterovirus Replication That Targets Host Cell Factor
26 Phosphatidylinositol 4-Kinase III β . *Antimicrob. Agents Chemother.* **2013**, *57*, 4971-4981.
27
28

29 (15) Di Paolo, G.; De Camilli, P. Phosphoinositides in Cell Regulation and Membrane
30 Dynamics. *Nature* **2006**, *443*, 651-657.
31
32

33 (16) Clayton, E. L.; Minogue, S.; Waugh, M. G. Mammalian Phosphatidylinositol 4-
34 Kinases as Modulators of Membrane Trafficking and Lipid Signaling Networks. *Prog. Lipid*
35 *Res.* **2013**, *52*, 294-304.
36
37

38 (17) Balla, T. Phosphoinositides: Tiny Lipids With Giant Impact on Cell Regulation.
39 *Physiol. Rev.* **2013**, *93*, 1019-1137.
40
41
42
43
44
45
46
47
48
49
50
51

1
2
3 (18) Himmel, B.; Nagel, G. Protein Kinase-Independent Activation of CFTR by
4 Phosphatidylinositol Phosphates. *EMBO Rep.* **2004**, *5*, 85-90.
5
6

7
8 (19) Balla, A.; Balla, T. Phosphatidylinositol 4-Kinases: Old Enzymes with Emerging
9 Functions. *Trends Cell Biol.* **2006**, *16*, 351-361.
10
11

12
13 (20) Altan-Bonnet, N.; Balla, T. Phosphatidylinositol 4-Kinases: Hostages Harnessed to
14 Build Panviral Replication Platforms. *Trends Biochem. Sci.* **2012**, *37*, 293-302.
15
16

17
18 (21) Boura, E.; Nencka, R. Phosphatidylinositol 4-Kinases: Function, Structure, and
19 Inhibition. *Exp. Cell Res.* **2015**, *337*, 136-145.
20
21

22
23 (22) Mejdrová, I.; Chalupská, D.; Kögler, M.; Šála, M.; Plačková, P.; Baumlová, A.;
24 Hřebabecký, H.; Procházková, E.; Dejmek, M.; Guillon, R.; Strunin, D.; Weber, J.; Lee, G.;
25 Birkus, G.; Mertlíková-Kaiserová, H.; Boura, E.; Nencka, R. Highly Selective
26 Phosphatidylinositol 4-Kinase III β Inhibitors and Structural Insight into Their Mode of
27 Action. *J. Med. Chem.* **2015**, *58*, 3767-3793.
28
29
30
31
32
33

34
35 (23) MacLeod, A. M.; Mitchell, D. R.; Palmer, N. J.; Van de Poël, H.; Conrath, K.;
36 Andrews, M.; Leyssen, P.; Neyts, J. Identification of a Series of Compounds with Potent
37 Antiviral Activity for the Treatment of Enterovirus Infections. *ACS Med. Chem. Lett.* **2013**, *4*,
38 585-589.
39
40
41
42
43
44

45 (24) LaMarche, M. J.; Borawski, J.; Bose, A.; Capacci-Daniel, C.; Colvin, R.; Dennehy,
46 M.; Ding, J.; Dobler, M.; Drumm, J.; Gaither, L. A.; Gao, J.; Jiang, X.; Lin, K.; McKeever,
47 U.; Puyang, X.; Raman, P.; Thohan, S.; Tommasi, R.; Wagner, K.; Xiong, X.; Zabawa, T.;
48 Zhu, S.; Wiedmann, B. Anti-Hepatitis C Virus Activity and Toxicity of Type III
49 Phosphatidylinositol-4-Kinase Beta Inhibitors. *Antimicrob. Agents Chemother.* **2012**, *56*,
50 5149-5156.
51
52
53
54
55
56
57
58
59
60

1
2
3 (25) Balla, A.; Tuymetova, G.; Toth, B.; Szentpetery, Z.; Zhao, X.; Knight, Z. A.; Shokat,
4 K.; Steinbach, P. J.; Balla, T. Design of Drug-Resistant Alleles of Type-III
5 Phosphatidylinositol 4-Kinases Using Mutagenesis and Molecular Modeling. *Biochemistry*
6 **2008**, *47*, 1599-1607.

7
8
9
10
11
12 (26) Knight, Z. A.; Gonzalez, B.; Feldman, M. E.; Zunder, E. R.; Goldenberg, D. D.;
13 Williams, O.; Loewith, R.; Stokoe, D.; Balla, A.; Toth, B.; Balla, T.; Weiss, W. A.; Williams,
14 R. L.; Shokat, K. M. A Pharmacological Map of the PI3-K Family Defines a Role for
15 p110alpha in Insulin Signaling. *Cell* **2006**, *125*, 733-747.

16
17
18
19
20
21 (27) Anighoro, A.; Bajorath, J.; Rastelli, G. Polypharmacology: Challenges and
22 Opportunities in Drug Discovery: Miniperspective. *J. Med. Chem.* **2014**, *57*, 7874-7887.

23
24
25
26
27 (28) Reddy, A. S.; Zhang, S. Polypharmacology: Drug Discovery for the Future. *Expert*
28 *Rev. Clin. Pharmacol.* **2013**, *6*, 41-47.

29
30
31
32 (29) Bowers, K. J.; Chow, E.; Xu, H.; Dror, R. O.; Eastwood, M. P.; Gregersen, B. A.;
33 Klepeis, J. L.; Kolossvary, I.; Moraes, M. A.; Sacerdoti, F. D.; Salmon, J. K.; Shan, Y.;
34 Shaw, D. E. Scalable Algorithms for Molecular Dynamics Simulations on Commodity
35 Clusters. Proceedings of the 2006 ACM/IEEE conference on Supercomputing; ACM: New
36 York, 2006.

37
38
39
40
41 (30) Wang, Y.; Suzek, T.; Zhang, J.; Wang, J.; He, S.; Cheng, T.; Shoemaker, B. A.;
42 Gindulyte, A.; Bryant, S. H. PubChem BioAssay: 2014 update. *Nucleic Acids Res.* **2014**, *42*
43 (Database issue), D1075-1082.

44
45
46
47 (31) <http://www.asinex.com/> accessed on February 14, 2014.

48
49
50
51
52 (32) Pedemonte, N.; Sonawane, N. D.; Taddei, A.; Hu, J.; Zegarra-Moran, O.; Suen, Y. F.;
53 Robins, L. I.; Dicus, C. W.; Willenbring, D.; Nantz, M. H.; Kurth, M. J.; Galiotta, L. J.;
54
55
56
57
58
59
60

1
2
3 Verkman, A. S. Phenylglycine and Sulfonamide correctors of Defective Delta F508 and
4
5 G551D Cystic Fibrosis Transmembrane Conductance Regulator Chloride-Channel Gating.
6
7 *Mol. Pharmacol.* **2005**, *67*, 1797-1807.
8
9

10 (33) Pedemonte, N.; Lukacs, G. L.; Du, K.; Caci, E.; Zegarra-Moran, O.; Galiotta, L. J.;
11
12 Verkman, A. S. Small-molecule Correctors of Defective DeltaF508-CFTR Cellular
13
14 Processing Identified by High-Throughput Screening. *J. Clin. Invest.* **2005**, *115*, 2564-2571.
15
16

17 (34) Burke, J. E.; Inglis, A. J.; Perisic, O.; Masson, G. R.; McLaughlin, S. H.; Rutaganira,
18
19 F.; Shokat, K. M.; Williams, R. L. Structures of PI4KIII β Complexes Show Simultaneous
20
21 Recruitment of Rab11 and Its Effectors. *Science* **2014**, *344*, 1035-1038.
22
23

24 (35) Yoo, C. L.; Yu, G. J.; Yang, B.; Robins, L. I.; Verkman, A. S.; Kurth, M. J. 4'-Methyl-
25
26 4,5'-Bithiazole-Based Correctors of Defective Δ F508-CFTR Cellular Processing. *Bioorg.*
27
28 *Med. Chem. Lett.* **2008**, *18*, 2610-2614.
29
30

31 (36) Donald, M. B.; Rodriguez, K. X.; Shay, H.; Phuan, P. W.; Verkman, A. S.; Kurth, M.
32
33 J. Click-Based Synthesis of Triazolobithiazole Δ F508-CFTR Correctors for Cystic Fibrosis.
34
35 *Bioorg. Med. Chem.* **2012**, *20*, 5247-5253.
36
37

38 (37) Haydon, D. J.; Bennett, J. M.; Brown, D.; Collins, I.; Galbraith, G.; Lancett, P.;
39
40 Macdonald, R.; Stokes, N. R.; Chauhan, P. K.; Sutariya, J. K.; Nayal, N.; Srivastava, A.;
41
42 Beanland, J.; Hall, R.; Henstock, V.; Noula, C.; Rockley, C.; Czaplowski, L. Creating an
43
44 Antibacterial with in Vivo Efficacy: Synthesis and Characterization of Potent Inhibitors of
45
46 the Bacterial Cell Division Protein FtsZ with Improved Pharmacological Properties. *J. Med.*
47
48 *Chem.* **2010**, *53*, 3927-3936.
49
50

51 (38) De Lucca, G. V.; Kim, U. T.; Vargo, B. J.; Duncia, J. V.; Santella, J. B.; Gardner, D.
52
53 S.; Zheng, C.; Liauw, A.; Wang, Z.; Emmett, G.; Wacker, D. A.; Welch, P. K.; Covington,
54
55
56
57
58
59
60

1
2
3 M.; Stowell, N. C.; Wadman, E. A.; Das, A. M.; Davies, P.; Yeleswaram, S.; Graden, D. M.;
4
5 Solomon, K. A.; Newton, R. C.; Trainor, G. L.; Decicco, C. P.; Ko, S. S. Discovery of CC
6
7 Chemokine Receptor-3 (CCR3) Antagonists with Picomolar Potency. *J. Med. Chem.* **2005**,
8
9 *48*, 2194-2211.

10
11
12 (39) Diab, S.; Teo, T.; Kumarasiri, M.; Li, P.; Yu, M.; Lam, F.; Basnet, S. K. C.; Sykes, M.
13
14 J.; Alberecht, H.; Milne, R.; Wang, S. Discovery of 5-(2-(Phenylamino)pyrimidin-4-
15
16 yl)thiazol-2(3H)-one Derivatives as Potent Mnk2 Inhibitors: Synthesis, SAR Analysis and
17
18 Biological Evaluation. *ChemMedChem* **2014**, *9*, 962-972.

19
20
21
22 (40) Sperandio, O.; Petitjean, M.; Tuffery, P. wwwLigCSRre: a 3D Ligand-Based Server for
23
24 Hit Identification and Optimization. *Nucleic Acids Res.* **2009**, *37* (Web Server), W504-W509.

25
26
27
28 (41) (a) Radi, M.; Falchi, F.; Garbelli, A.; Samuele, A.; Bernardo, V.; Paolucci, S.;
29
30 Baldanti, F.; Schenone, S.; Manetti, F.; Maga, G.; Botta, M. Discovery of the First Small
31
32 Molecule Inhibitor of Human DDX3 Specifically Designed to Target the RNA Binding Site:
33
34 Towards the next Generation HIV-1 Inhibitors. *Bioorg. Med. Chem. Lett.* **2012**, *22*, 2094-
35
36 2098. (b) Bloom, J. D.; Digrandi, M. J.; Dushin, R. G.; Curran, K. J.; Ross, A. A.; Norton, E.
37
38 B.; Terefenko, E.; Jones, T. R.; Feld, B.; Lang, S. A. Thiourea Inhibitors of Herpes Viruses.
39
40 Part 1: Bis-(Aryl)thiourea Inhibitors of CMV. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 2929-
41
42 2932.

43
44
45
46 (42) Borrok, M. J.; Kiessling, L. Non-Carbohydrate Inhibitors of the Lectin DC-SIGN. *J.*
47
48 *Am. Chem. Soc.* **2007**, *129*, 12780-12785.

49
50
51 (43) Ren, X. X.; Ma, L.; Liu, Q. W.; Li, C.; Huang, Z.; Wu, L.; Xiong, S. D.; Wang, J. H.;
52
53 Wang, H. B. The Molecule of DC-SIGN Captures Enterovirus 71 and Confers Dendritic
54
55 Cell-Mediated Viral Trans-Infection. *Viol. J.* **2014**, *11*, 47.

1
2
3 (44) Ye, L.; Hu, B.; El-Badri, F.; Hudson, B. M.; Phuan, P. W.; Verkman, A. S.; Tantillo,
4 S. J.; Kurth, M. J. Δ F508-CFTR Correctors: Synthesis and Evaluation of Thiazole-Tethered
5 Imidazolones, Oxazoles, Oxadiazoles, and Thiadiazoles. *Bioorg. Med. Chem. Lett.* **2014**, *24*,
6 5840-5844.
7
8

9
10
11
12 (45) Ford Siltz, L. A.; Viktorova, E. G.; Zhang, B.; Kouiyavskaya, D.; Dragunsky, E.;
13 Chumakov, K.; Isaacs, L.; Belov, G. A. New Small-Molecule Inhibitors Effectively Blocking
14 Picornavirus Replication. *J. Virol.* **2014**, *88*, 11091-11107.
15
16
17

18
19
20 (46) Leong, W. F.; Chow, V. T. Transcriptomic and Proteomic Analyses of
21 Rhabdomyosarcoma Cells Reveal Differential Cellular Gene Expression in Response to
22 Enterovirus 71 Infection. *Cell. Microbiol.* **2006**, *8*, 565-580.
23
24
25
26

27 (47) Van Goor, F.; Hadida, S.; Grootenhuys, P. D. J.; Burton, B.; Stack, J. H.; Straley, K. S.;
28 Decker, C. J.; Miller, M.; McCartney, J.; Olson, E. R.; Wine, J. J.; Frizzell, R. A.; Ashlock,
29 M.; Negulescu, P. A. Correction of the F508del-CFTR Protein Processing Defect in Vitro by
30 the Investigational Drug VX-809. *Proc. Natl. Acad. Sci. U. S. A.* **2011**, *108*, 18843–18848.
31
32
33
34
35
36

37 (48) Langron, E.; Simone, M. I.; Delalande, C. M. S.; Reymond, J.-L.; Selwood, D. L.;
38 Vergani, P. Improved Fluorescence Assays to Measure the Defects Associated with F508del-
39 CFTR Allow Identification of New Active Compounds. *Br. J. Pharmacol.* doi:
40 10.1111/bph.13715.
41
42
43
44
45

46 (49) Van Goor, F.; Hadida, S.; Grootenhuys, P. D. J.; Burton, B.; Cao, D.; Neuberger, T.;
47 Turnbull, A.; Singh, A.; Joubran, J.; Hazlewood, A.; Zhou, J.; McCartney, J.; Arumugam, V.;
48 Decker, C.; Yang, J.; Young, C.; Olson, E. R.; Wine, J. J.; Frizzell, R. A.; Ashlock, M.;
49 Negulescu, P. Rescue of CF Airway Epithelial Cell Function in Vitro by a CFTR Potentiator,
50 VX-770. *Proc. Natl. Acad. Sci. U. S. A.* **2009**, *106*, 18825–18830.
51
52
53
54
55
56
57
58
59
60

1
2
3 (50) Friesner, R. A.; Banks, J. L.; Murphy, R. B.; Halgren, T. A.; Klicic, J. J.; Mainz, D. T.;
4
5 Repasky, M. P.; Knoll, E. H.; Shaw, D. E.; Shelley, M.; Perry, J. K.; Francis, P.; Shenkin, P.
6
7 S. Glide: A New Approach for Rapid, Accurate Docking and Scoring. 1. Method and
8
9 Assessment of Docking Accuracy. *J. Med. Chem.* **2004**, *47*, 1739–1749.
10
11

12 (51) Warren, G. L.; Andrews, C. W.; Capelli, A.-M.; Clarke, B.; LaLonde, J.; Lambert, M.
13
14 H.; Lindvall, M.; Nevins, N.; Semus, S. F.; Senger, S.; Tedesco, G.; Wall, I. D.; Woolven, J.
15
16 M.; Peishoff, C. E.; Head, M. S. A Critical Assessment of Docking Programs and Scoring
17
18 Functions. *J. Med. Chem.* **2006**, *49*, 5912–5931.
19
20
21

22 (52) Manetti, F.; Falchi, F.; Crespan, E.; Schenone, S.; Maga, G.; Botta, M. N-(Thiazol-2-
23
24 Yl)-2-Thiophene Carboxamide Derivatives as Abl Inhibitors Identified by a Pharmacophore-
25
26 Based Database Screening of Commercially Available Compounds. *Bioorg. Med. Chem. Lett.*
27
28 **2008**, *18*, 4328–4331.
29
30
31

32 (53) Jacobson, M. P.; Pincus, D. L.; Rapp, C. S.; Day, T. J. F.; Honig, B.; Shaw, D. E.;
33
34 Friesner, R. A. A Hierarchical Approach to All-Atom Protein Loop Prediction. *Proteins:*
35
36 *Struct. Funct. Bioinf.* **2004**, *55*, 351-367.
37
38

39 (54) Bates, P. A.; Kelley, L. A.; MacCallum, R. M.; Sternberg, M. J. E. Enhancement of
40
41 Protein Modeling by Human Intervention in Applying the Automatic Programs 3D-JIGSAW
42
43 and 3D-PSSM. *Proteins: Struct. Funct. Bioinf.* **2001**, Suppl. 5, 39-46.
44
45
46

47 (55) *Schrödinger Release 2014-3: Maestro*, version 9.9, Schrödinger, LLC, New York,
48
49 NY, 2014.
50
51

52 (56) Irwin, J. J.; Sterling, T.; Mysinger, M. M.; Bolstad, E. S.; Coleman, R. G. ZINC: a
53
54 Free Tool to Discover Chemistry for Biology. *J. Chem. Inf. Model.* **2012**, *52*, 1757-1768.
55
56
57
58
59
60

1
2
3 (57) *Schrödinger Release 2015-4: LigPrep*, version 3.6, Schrödinger, LLC, New York, NY,
4
5 2015.

6
7
8 (58) (a) *Schrödinger Release 2015-4: Schrödinger Suite 2015-4 Protein Preparation*
9
10 *Wizard*; Epik version 3.4, Schrödinger, LLC, New York, NY, 2015; (b) Impact version 6.9,
11
12 Schrödinger, LLC, New York, NY, 2015; Prime version 4.2, Schrödinger, LLC, New York,
13
14 NY, 2015.

15
16
17 (59) Harder, E.; Damm, W.; Maple, J.; Wu, C.; Reboul, M.; Xiang, J.Y.; Wang, L.;
18
19 Lupyan, D.; Dahlgren, M.K.; Knight, J.L.; Kaus, J.W.; Cerutti, D.S.; Krilov, G.; Jorgensen,
20
21 W.L.; Abel, R.; Friesner, R.A. OPLS3: A Force Field Providing Broad Coverage of Drug-
22
23 like Small Molecules and Proteins. *J. Chem. Theory Comput.* **2016**, *12*, 281-296.

24
25
26 (60) *Small-Molecule Drug Discovery Suite 2015-4: Glide*, version 6.9, Schrödinger, LLC,
27
28 New York, NY, 2015.

29
30
31 (61) Galiotta, L. J. V.; Haggie, P. M.; Verkman, A. S. Green Fluorescent Protein-Based
32
33 Halide Indicators with Improved Chloride and Iodide Affinities. *FEBS Lett.* **2001**, *499*, 220-
34
35 224.

36
37
38 (62) Li, Y.; Tsien, R. W. pHTomato, a Red, Genetically Encoded Indicator That Enables
39
40
41 Multiplex Interrogation of Synaptic Activity. *Nat. Neurosci.* **2012**, *15*, 1047-1053.

42
43
44 (63) Tassini, S.; Castagnolo, D.; Scalacci, N.; Kissova, M.; Armijos-Rivera, J.I.; Giagnorio,
45
46 F.; Maga, G.; Costantino, G.; Crespan, E.; Radi, M. A multicomponent pharmacophore
47
48 fragment-decoration approach to identify selective LRRK2-targeting probes. *Med. Chem.*
49
50 *Commun.* **2016**, *7*, 484-494.
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table of Contents Graphic

