

1 **Prenatal diagnosis and management of vasa**
2 **previa in twin pregnancies: A case series**
3 **and systematic review**
4
5
6
7
8
9

10 Eric JAUNIAUX¹,MD,PhD
11 Yaakov MELCER²,MD
12 Ron MAYMON²,MD
13
14
15
16

17 ¹*Academic Department of Obstetrics and Gynaecology, UCL Institute for*
18 *Women’s Health, University College London (UCL), London, UK;*
19 ²*Department of Obstetrics and Gynecology, Assaf Harofeh Medical Center,*
20 *Zerifin, Israel, affiliated with the Sackler School of Medicine, Tel-Aviv University,*
21 *Tel-Aviv, Israel*
22
23
24
25
26
27
28
29

30 The authors report no conflict of interest
31

32 No funding was obtained for this study.
33
34
35
36

37 **Corresponding author:** *Professor Eric Jauniaux, Academic Department of*
38 *Obstetrics and Gynaecology, Institute for Women’s Health, University College*
39 *London, 86-96 Chenies Mews, London WC1E 6HX, UK. Telephone numbers:*
40 *+44/207/3908113 Fax: +44/207/3908115*
41 *E-mail: e.jauniaux@ucl.ac.uk*
42
43
44

45 **Word count:** 3208
46
47

48 **Condensation:** Management strategies in twin pregnancies presenting with
49 vasa praevia should be tailored on the basis of chorionicity and changes in
50 cervical length.
51
52

53 **Short title:** Management of vasa previa in twins
54
55

56 **Key Words:** Velamentous; umbilical cord; vasa previa; twins; placenta.
57
58
59
60
61
62
63
64
65

Abstract

Background: Twin pregnancies are at higher risks of velamentous cord insertion (VCI) and vasa previa (VP). In-vitro fertilization (IVF) is an additional direct risk factor of abnormal cord insertion and thus the incidence of VP is likely to increase over the next decades.

Objective: Our aim was to evaluate the specific challenges associated with the prenatal diagnosis and management of VP in twins.

Study design: We searched our database for twin pregnancies diagnosed with VP and managed antenatally using measurements of cervical length (CL) and performed a systematic review of articles which correlated prenatal diagnosis of VP in twins and pregnancy outcome. **Data sources:** PubMed and MEDLINE were searched for article published between the first prenatal ultrasound description of VP in 1987 and 20 October 2016 using medical subject heading (MeSH) terms, keywords and their combination including “vasa previa”, “velamentous insertion of the cord”, “velamentous vessel”, “membranous umbilical vessels”, “twin pregnancy”, “multiple pregnancy”, “placenta previa” “In vitro fertilization pregnancy”, “succenturiate lobe”, “low-lying placenta” and “bilobate placenta” as related to “sonography”, “ultrasound diagnosis”, “prenatal diagnosis” and “colour Doppler”. **Study eligibility criteria:** The primary eligibility criteria were articles which correlated prenatal ultrasound imaging of VP and pregnancy outcome. The secondary outcome was the use of CL in the management of twin pregnancies diagnosed antenatally with VP.

Study appraisal and synthesis methods: The final selection included 3 case

1 report series, 9 retrospective cohort studies and 1 retrospective case-control
2
3 study of VP diagnosed prenatally and confirmed at birth in twin pregnancies.
4
5

6
7 **Results:** The search of our databases identified six cases of dichorionic-
8
9 diamniotic (DCDA) twins and 1 case of monochorionic-diamniotic (MCDA) twins
10 diagnosed prenatally with VP between 22 and 29 weeks and managed using
11 CL. Two cases were delivered by emergency because of rapid changes in CL in
12 one and bleeding on placenta praevia in the other at 33 weeks and 30 weeks,
13 respectively. The systematic review identified data on 56 cases. The incidence
14 of twin pregnancies diagnosed antenatally with VP in the cohort and case-
15 control studies was 11.0%. Data on chorionicity was available in only 34 cases
16 and CL measurements were used by only the authors of two case reports and
17 four cohort studies. VCI was the most common additional ultrasound findings in
18 twins presenting with VP in both our series and the systematic review.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34
35 **Conclusions:** VP is associated with specific prenatal and obstetric
36 complications with different outcomes in singletons compared to twins. Data on
37 the diagnosis and management of VP in twin pregnancies are limited but there
38 is enough evidence to warrant guidelines for targeted screening. To enable the
39 development of efficient management protocols tailored to the need of individual
40 cases, future studies of the screening, diagnosis and management of VP should
41 be prospective and multicentric with detailed data on twins including chorionicity
42 and use of CL.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Introduction

A vasa previa (VP) describes an aberrant chorionic vessel directly connected to the umbilical cord circulation but running between the amniotic and the chorionic layers of the placental free membranes below the fetal presenting part.¹ VP are classically separated into 2 different categories: type I which describes vessels that connect a velamentous cord to the main placental mass and type II which describes vessels running between one or more accessory lobes of the placenta.¹ The classic presentation of undiagnosed VP in labour is the presence of painless vaginal bleeding. This occurs mainly when the cervix is effaced and dilated and the membranes rupture spontaneously or are ruptured artificially. It can lead to rapid fetal exsanguination and death¹⁻⁵ and the perinatal mortality of undiagnosed VP has been reported to be as high as 60%.⁶

The incidence of VP has been reported to range between in 1 in 1200 to 1 in 5000 pregnancies but the condition may have been under-reported.¹⁻⁵ A recent systematic review of 10 cohort studies found that the incidence of VP is 0.6 per 1000 pregnancies.⁷ Approximately 90% of women presenting with a VP also have a velamentous cord insertion (VCI) and 3-4% of women presenting with a VCI have a VP.⁸ Unlike the single umbilical artery cord which is routinely recorded by midwives at birth, these cord anomalies are only recorded when associated with perinatal complications and the placental is examined by a pathologist.

The pathophysiology of VP and VCI is uncertain but it has been linked to an abnormal development of the definitive placenta. Both cord anomalies are more common in bilobated placenta, succenturiate or accessory placenta and

1 low-lying placenta.¹⁻⁸ VCI and VP type I are also more common in twin
2 pregnancies and monochorionicity doubles the risk for VCI compared to
3 dichorionicity.^{11,12} Maternal smoking has been recently reported as a risk factor
4 for VCI and thus potentially of VP.¹³ Assisted reproduction technology (ART)
5 and in vitro fertilization (IVF) in particular is associated with a higher incidence
6 of abnormally shaped placenta, placenta previa and cord insertion outside the
7 placental chorionic plate.^{7,14} It has been hypothesized these placental and cord
8 anomalies could be due to the inadequate orientation of the IVF blastocyst at
9 the time of implantation or to a higher incidence of vanishing twins in IVF than in
10 spontaneous twins.^{10,14-18} The incidence of VP has been reported to be as high
11 as one in 300 pregnancies conceived after ART.^{7,14} These data suggest that the
12 incidence of VP in both singleton and multiple pregnancy gestations is likely to
13 increase in the next decade with the increase use of ART worldwide.

14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Recent national guidelines, expert reviews and decision analysis study on the management of VP have recommended delivery of all women presenting with VP between 34-36 weeks of gestation but none provide with a specific algorithm for twins.^{3-6,8,19} Twin pregnancies are associated with specific obstetric complications and in particular with a higher incidence of premature delivery.²⁰ Like singletons, twins presenting with VP have very poor outcomes when antenatal diagnosis is not made before the onset of labour.²¹ Previous studies and reviews have not identified twin pregnancies diagnosed with VP during pregnancy as requiring a different management strategy. We present a case series and a systematic review of the literature of twin pregnancies diagnosed prenatally with VP. Our aim is to evaluate the role of ultrasound

1 imaging in optimizing the management of individual cases of twins diagnosed
2
3 antenatally with VP and in supporting the development of the corresponding
4
5 clinical guidelines.
6
7
8
9

10 11 12 **Material and Methods**

13 14 15 ***Case series***

16
17
18 The location of the placental insertion of the umbilical cord is included in our
19
20 routine mid-trimester transabdominal fetal detailed anatomy ultrasound protocol
21
22 in targeted cases i.e. in cases of low-lying placenta, bilobate placenta,
23
24 succenturiate lobed placenta, multiple pregnancies and IVF conceptions. When
25
26 a VCI is identified on transabdominal ultrasound, a transvaginal scan (TVS) is
27
28 performed to exclude a VP. Our units have collaborated on the use of cervical
29
30 length (CL) in the management of twin pregnancies^{22,23} and are using similar
31
32 protocols based on national and international guidelines for the prenatal
33
34 diagnosis of VP. Asymptomatic women presenting with VP are followed-up
35
36 every 2 weeks, from 26-28 weeks until delivery, for fetal growth and wellbeing
37
38 and with TVS for CL and VP position. The timing of delivery is scheduled
39
40 according to changes in CL and/or clinical symptoms (mainly uterine
41
42 contractions and/or vaginal bleeding), following a course of corticosteroids.
43
44 When the CL remains stable with normal fetal development and there are no
45
46 clinical symptoms delivery is planned at around 36 weeks.
47
48
49
50
51
52
53
54

55 56 ***Systematic review information sources and search strategy***

57
58 The search protocol was designed a priori and registered on PROSPERO
59
60
61
62
63
64
65

1 (#50164). We performed an electronic search of the literature in PubMed and
2
3 MEDLINE and selected relevant studies that have been published between the
4
5 first prenatal ultrasound description of VP by Gianopoulos *et al.*²⁴ in 1987 and
6
7 20 October 2016. The search was achieved using medical subject heading
8
9 (MeSH) terms, keywords and their combination including “vasa previa”,
10
11 “velamentous insertion of the cord”, “velamentous vessel”, “membranous
12
13 umbilical vessels”, “twin pregnancy”, “multiple pregnancy”, “placenta previa” “In
14
15 vitro fertilization pregnancy”, “succenturiate lobe”, “low-lying placenta” and
16
17 “bilobate placenta” as related to “sonography”, “ultrasound diagnosis”, “prenatal
18
19 diagnosis” and “colour Doppler”.
20
21
22
23
24

25 We included all cohort studies and case reports that described the
26
27 prenatal diagnosis of VP in women presenting with a twin pregnancy (Figure 1).
28
29

30 ***Systematic review eligibility criteria***

31
32 The primary eligibility criteria were articles which correlated prenatal ultrasound
33
34 imaging of VP and pregnancy outcome. The secondary outcome was the use of
35
36 CL in the management of twin pregnancies diagnosed antenatally with VP. Two
37
38 authors (EJ and RM) independently assessed inclusion criteria, data extraction
39
40 and analysis.
41
42
43
44

45 ***Systematic review study selection***

46
47 The initial database search provided 44 reports and cross-referencing provided
48
49 an additional two reports, making a total of 43 records after removal of three
50
51 duplicates (Figure 1). Out of the 43 records screened, 13 did not included data
52
53 on prenatal ultrasound imaging of VP and were therefore excluded. After the
54
55 second selection, letters with no description of the case, commentaries and
56
57
58
59
60
61
62
63
64
65

1 reviews were excluded. A further six reports where antenatal ultrasound was
2 performed but cases of VP were only diagnosed at delivery were further
3 excluded leaving 13 reports for the final analysis.
4
5
6

7 ***Statistical analysis***

8 Numerical data from the case series and systematic review were analysed
9 using the StatGraphic data analysis and statistical software package
10 (Manugistics, Rockville, MD).
11
12
13
14
15
16
17
18
19
20

21 **Results**

22 ***Case series***

23 The search of our databases identified seven sets of twins diagnosed prenatally
24 with VP, followed-up using our standard protocol and confirmed at birth by
25 detailed pathologic examination. Table 1 presents the clinical data and outcome
26 of these cases. There were six cases of dichorionic-diamniotic (DCDA) twins
27 and 1 case of monochorionic-diamniotic (MCDA) twins (case 7). Five DCDA
28 cases were conceived following IVF. The mean (SD) gestational at time of
29 diagnosis was 26+5 (3+1) weeks. In five cases, a VCI was detected on
30 transabdominal ultrasound at the mid-trimester examination or during a fetal
31 growth scan follow-up. In the other two cases, the diagnosis of VP was made at
32 the time of the first CL measurement. In five cases, the CL remained >25 mm
33 and the patients were managed conservatively without hospitalization. In one
34 case, the CL was found to be 10 mm at 32+6 weeks (case 5) and an
35 emergency CS performed the next day because of vaginal bleeding. One case
36 was complicated by placenta previa accreta (case 6) for which an emergency
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 caesarean delivery was performed at 30+3 weeks because of major vaginal
2 bleeding. In the case of MCDA twins (case 7), the cord of the presenting twin
3 (fetus 1) was inserted half-way between the lower placental edge and the
4 cervix. On TVS, there was a large VP above the internal os with non-pulsatile
5 flow on Doppler imaging (Figures 2 A & B). This venous VP could be traced
6 back to the area of the VCI with colour mapping. No VP arteries were found.
7
8 The CL was stable on follow-up scans and there was no ultrasound evidence of
9 twin-to-twin transfusion syndrome (TTTS) but a 20% discrepancy in estimated
10 fetal weight was noted from 32+4 weeks and an elective caesarean delivery
11 was performed at 35+1 weeks. The mean (SD) gestational at the time of
12 delivery was 34.2 (2.1) weeks.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

28 ***Systematic review report characteristics***

29 We found data on 56 cases of VP diagnosed prenatally in twin pregnancies,
30 confirmed at birth and included in 3 case report series,²⁵⁻²⁷ 9 retrospective
31 cohort studies^{6,17,28-34} and 1 retrospective case-control study³⁵ (Table 2). The
32 incidence of twin pregnancies diagnosed with VP on ultrasound during
33 pregnancy for which ultrasound data and post-natal confirmation was available
34 in the cohort and case-control studies was 11.0% (48 cases of twins for 438
35 pregnancies diagnosed with VP).
36
37
38
39
40
41
42
43
44
45
46
47

48 ***Systematic review synthesis of results***

49 Out of the 34 cases for which data on chorionicity were available, there were 19
50 DCDA and 15 MCDA twin pregnancies. In one case report, the diagnosis of VP
51 was made at the time of laser photocoagulation of communicating anastomoses
52 in a MCDA twin pregnancy complicated by twin-twin transfusion syndrome
53
54
55
56
57
58
59
60
61
62
63
64
65

1 (TTTS).²⁷ The authors of two case reports^{26,27} and of four cohorts³¹⁻³⁴ described
2 the use of CL measurements in their management of twin pregnancies
3
4 presenting with VP on antenatal ultrasound.
5
6
7

8 Only two authors of cohort studies included perinatal data on individual
9 twin pregnancies that were diagnosed by ultrasound antenatally.^{29,31} Detailed
10 measurements of CL were only provided in two cases in a cohort series³¹ and in
11 one case report.²⁷ Table 3 displays the baseline perinatal characteristics of 12
12 cases of VP diagnosed antenatally and verified by clinical documentation at
13 birth and/or a pathology report. In seven cases, there was information on
14 additional ultrasound findings including a low-lying placenta and VCI in three
15 cases^{25,26} and VCI alone in two cases.^{17,28} The mean (SD) gestational at the
16 time of diagnosis was 23+6 (7+1) weeks. Information on the mode of delivery
17 was available in eight cases including five elective and three emergency
18 caesarean sections, two for premature rupture of the membranes^{27,28} and one
19 for preeclampsia.¹⁷ The mean (SD) gestational at time of delivery was 32+6
20 (3+5) weeks. The neonatal outcome of the corresponding 24 fetuses including
21 two stillbirths and two neonatal deaths.^{29,31}
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 Discussion

47 *Main findings*

48 Our review indicates a total incidence of 11.0% of twin pregnancies in cohort
49 studies on the prenatal diagnosis of VP. Our case series and systematic review
50 confirm that like for singletons, the outcome for twin fetuses and neonates is
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 excellent when the VP is diagnosed antenatally. Determination of chorionicity
2 and serial measurements of CL can provide individualised management
3
4 pathways for twin pregnancies with VP and in particular in the timing of delivery
5
6 or the need for emergent caesarean section before a planned delivery around
7
8
9
10 36 weeks.
11

12 13 14 15 16 ***Comparison with existing literature*** 17

18 A 97% survival rate has been reported in cases of VP diagnosis prenatally
19 compared to only 44% when the diagnosis was made during delivery.⁶ A recent
20
21 systematic review of ultrasound in the diagnosis of VP including two prospective
22
23 and six retrospective cohort studies out of which six had a poor methodology
24
25 found prenatal detection rates ranging between 53% (10/19) to 100% for a total
26
27 of 442,633 women, including 138 cases of VP.³⁶ Four out of the eight studies
28
29 used TVS for primary assessment, while the remaining four studies used
30
31 transabdominal ultrasound and only used TVS when a VP was suspected on
32
33 the transabdominal scan. The gestational age at the time of diagnosis was
34
35 reported in five studies and ranged from 18+0 to 26+6 weeks. The data of twin
36
37 pregnancies were not analyzed separately by the authors. The accuracy of
38
39 ultrasound at 9-13 weeks' gestation in screening for VP was evaluated
40
41 prospectively in case-control study.³⁷ However, the definitive placenta is only
42
43 fully formed by 10-11 weeks of gestation¹ making it difficult in many cases to
44
45 predict the final position of the umbilical cord and to identify placental
46
47 morphologic anomalies in particular in twins. The performance of ultrasound in
48
49 screening for and diagnosing VP considered more efficient at around mid-
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 gestation^{3-5,36} and the mean gestational age at diagnosis in twins, in our series
2
3 and in the literature, was 26+5 and 23+6 weeks, respectively.
4

5
6 Cases of VP in both singletons and twins described in the literature
7
8 present a pulsatile flow velocity waveforms indicative of an arterial previa
9
10 vessel.^{3-5,6,31} One of our case (case 7) presented with venous VP and the
11
12 arteries were not visible on TVS color mapping (Figure 2B). We found no similar
13
14 case described in the international literature. A venous VP could be easily
15
16 mistaken for “free loop” of cord³ and thus in particular in this case it may be
17
18 useful to trace the route of the vessel to the VIC and to demonstrate that it does
19
20 not “move” with changing maternal position and/or between ultrasound
21
22 examinations.
23
24
25
26

27
28 Several studies have suggested that the majority of cases of VP can be
29
30 suspected prenatally by routinely evaluating the placental cord insertion in
31
32 pregnancies with low-lying placenta and the placental morphology.^{4,6-8, 29,38} The
33
34 incidence of succenturiate lobes and other morphological placental anomalies is
35
36 higher in twin pregnancies.^{1,11,39,40} A VCI of one of the umbilical cords is eight
37
38 times more common in twins than in singletons.³⁹ The odd ratio for VP in
39
40 pregnancies presenting with a bilobated placenta or with succenturiate lobe has
41
42 been evaluated to be 22.11 in twin pregnancies.³⁹ Five of the seven cases of
43
44 our series (Table 1) and five out of the eight cases described in details in the
45
46 literature (Table 3) presented with a VCI. One cases in our series and three in
47
48 the literature also presented a low-lying placenta or placenta previa. Thus, the
49
50 placental cord insertion should be documented at the second trimester scan in
51
52 all twin pregnancies.
53
54
55
56
57
58
59
60
61
62
63
64
65

1 The largest study to date on the perinatal outcome of VP is based on a
2
3 retrospective multicentric cohort of 155 women with VP diagnosed at the mid-
4
5 trimester scan and it included 50 (32.3%) cases associated with a bilobated or
6
7 succenturiate lobed placentas and 95 (61%) with low-lying placenta.⁶ Only
8
9 seven cases (4.5%) were twin pregnancies and the authors provided with no
10
11 data on chorionicity and individual outcome. Overall only half of the cohort
12
13 studies that are included in the present review provide detailed data on
14
15 chorionicity (Table 2). Similarly, to singleton pregnancies, a ruptured VP in the
16
17 presenting twin of a DC pair will lead to acute hemorrhage in the corresponding
18
19 twin. By contrast, in MC twins, the acute hemorrhage in the presenting twin may
20
21 also lead to acute blood shift and rapid exsanguination in the co-twin due to the
22
23 presence of inter-twin placental vascular anastomoses. Van Steenis *et al.*
24
25 reported recently a case of double fatal outcome in MC twins and reviewed
26
27 similar cases published in the international literature since the first case reports
28
29 in 1939.⁴¹ Including their case, there were three other recent cases reported in
30
31 the last five years in the international literature but surprisingly none were
32
33 diagnosed prenatally⁴²⁻⁴⁴ and were, therefore, not included in the present
34
35 review.
36
37
38
39
40
41
42
43
44
45
46

47 ***Clinical implications***

48
49 The ultimate management goal of confirmed VP prenatally is to deliver before
50
51 labor and in particular before the rupture of membranes while minimizing the
52
53 impact of iatrogenic prematurity. There is no consensus about the time of
54
55 delivery in cases of confirmed VP and the currently low prevalence of prenatal
56
57
58
59
60
61
62
63
64
65

1 diagnosis of this condition precludes any prospective trials to evaluate the ideal
2 timing. Data from a decision analysis study comparing 11 strategies for delivery
3 timing in a patient with VP found that delivery between 34 and 36 weeks
4 balances the risk of premature rupture of the membranes and subsequent fetal
5 hemorrhage and death versus the risks of prematurity.¹⁹ This study did not
6 differentiate between singleton and twin pregnancies. A cost-utility analysis
7 based on a decision-analytic model comparing relevant strategies and life-long
8 outcomes for mother and infant has shown that screening all twin pregnancies
9 for VP with transvaginal ultrasound is cost-effective and has incremental cost-
10 effectiveness ratio of \$5488 per quality-adjusted life-years gained.⁴⁵ We suggest
11 that in twins diagnosed with VP with uncomplicated antenatal course, to
12 minimize the impact of prematurity the optimal timing for scheduled caesarean
13 delivery is 36 weeks.

14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34 A recent systematic review and meta-analysis of the association between
35 placental implantation abnormalities including placenta previa, placenta accreta,
36 VP, and VIC and preterm delivery in singleton gestations has found a perinatal
37 death rate risk ratio of 4.52 (95% CI, 2.77-7.39) for VP.² The associated
38 complications of prematurity are, in many cases, the result of iatrogenic preterm
39 birth in an effort to prevent stillbirth. Antenatal hospitalization to allow for closer
40 surveillance for signs of labor in a unit with appropriate neonatal facilities has
41 also been proposed for all pregnancies presenting with VP from 30-32 weeks of
42 gestation^{3-5,31} but the evidence is weak and based on low-quality evidence.⁴ No
43 specific data are available for twins pregnancies diagnosed with VP.

1 Twin pregnancies are at higher risks of preterm delivery and a recent
2
3 retrospective cohort study of 441 women with twin pregnancies has found that
4
5 integration of serial measurements of CL using a stepwise algorithm in
6
7 asymptomatic women with twin gestations can improve the detection of women
8
9 at risk of preterm birth.⁴⁶ Women with VP in a MC twin pregnancy also require a
10
11 regular follow-up to assess the risks of TTTS and thus determining the twin
12
13 chorionicity is pivotal in the management of these high-risk pregnancies.
14
15
16
17
18
19
20

21 ***Strengths and weaknesses***

22
23 This study provides data supporting the need to screen for VP in twin
24
25 pregnancies in particular those resulting from IVF. A recent systematic review
26
27 and meta-analysis has found no clear evidence of benefit from planned
28
29 caesarean section for term twin pregnancies with leading cephalic
30
31 presentation⁴⁷ and thus for women opting for a vaginal birth it is essential to
32
33 have exclude a VP antenatally. Our data also support the determination of
34
35 chorionicity and use of transvaginal sonographic assessment of CL to evaluate
36
37 the individual risk of premature delivery in twins presenting with VP. Overall our
38
39 study is the first to highlights the need to incorporate these parameters in local
40
41 protocols and national guidelines in the perinatal management of twins. The
42
43 weakness of the study is its retrospective nature and the lack of specific details
44
45 on twins in most cohort series. In twins diagnosed with VP antenatally in the
46
47 literature, information on chorionicity was available in only 34 out of 56 (60.7%)
48
49 cases and CL measurements used by only six authors out of 13 thus limiting the
50
51 number of cases available for analysis.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Conclusions

VP is associated with specific prenatal and obstetric complications with different outcomes in singletons compared to twins. It is important to make the diagnosis in the second trimester to allow preventative measures to be taken and avoid subsequent perinatal complications. There is enough evidence to warrant guidelines for targeted screening with TVS and colour Doppler in twin pregnancies to rule out VP, as the relative obstetric risk is significantly higher than in singletons. In twin pregnancies, the ultrasound follow-up should also be tailored according to chorionicity, changes in cervical length and clinical symptoms. Outpatient management is possible if there is no evidence of cervical shortening on TVS and there are no symptoms of bleeding or preterm uterine activity. Multicentric prospective VP screening studies with detailed data on twins including their chorionicity are required to develop more efficient management protocols tailored to the need of individual cases.

References

1. Benirschke K, Burton GJ, Baergen RN. Pathology of the human placenta, 6th edition, Springer-Verlag, Berlin, 2012.
2. Vahanian SA, Lavery JA, Ananth CV, Vintzileos A. Placental implantation abnormalities and risk of preterm delivery: a systematic review and metaanalysis. *Am J Obstet Gynecol* 2015; 213: S78-90.
3. Silver BM. Abnormal placentation: Placenta previa, vasa previa and placenta accreta. *Obstet Gynecol* 2015; 126: 654-668.
4. Society of Maternal-Fetal (SMFM) Publications Committee, Sinkey RG, Odibo AO, Dashe JS. #37: Diagnosis and management of vasa previa. *Am J Obstet Gynecol* 2015; 213: 615-619.
5. Gagnon R, Morin L, Bly S, Butt K, Cargil YM, Denis N, Hietala-Coyle MA, Lim KI, Ouellet A, Racicot MH, Salem S, Hudon L, Basso M, Bos H, Delisle MF, Farine D, Grabowska K, Menticoglou S, Mundle W, Murphy-Kaulbeck L, Ouellet A, Pressey T, Roggensack A; Diagnostic Imaging Committee; Maternal Fetal Medicine Committee. SOGC CLINICAL PRACTICE GUIDELINE: guidelines for the management of vasa previa. *Int J Gynaecol Obstet* 2010; 108: 85-89.
6. Oyelese Y, Catanzarite V, Prefumo F, Lashley S, Schachter M, Tovbin Y, Goldstein V, Smulian JC. Vasa previa: the impact of prenatal diagnosis on outcomes. *Obstet Gynecol* 2004; 103: 937-942.
7. Ruitter L, Kok N, Limpens J, Derks JB, de Graaf IM, Mol B, Pajkrt E. Incidence of and risk indicators for vasa praevia: a systematic review. *BJOG* 2016; 123: 1278-1287.

- 1 8. Vintzileos AM, Ananth CV, Smulian JC. Using ultrasound in the clinical
2 management of placental implantation abnormalities. *Am J Obstet*
3
4
5
6 *Gynecol* 2015; 213: S70-77.
7
- 8 9. Räisänen S, Georgiadis L, Harju M, Keski-Nisula L, Heinonen S. Risk
9 factors and adverse pregnancy outcomes among births affected by
10
11 velamentous umbilical cord insertion: a retrospective population-based
12
13 register study. *Eur J Obstet Gynecol Reprod Biol* 2012; 165: 231-234.
14
15
- 16 10. Ebbing C, Kiserud T, Johnsen SL, Albrechtsen S, Rasmussen S.
17
18 Prevalence, risk factors and outcomes of velamentous and marginal cord
19
20 insertions: a population-based study of 634,741 pregnancies. *PLoS One*
21
22
23 2013; 8: e70380.
24
25
- 26 11. Hubinont C, Lewi L, Bernard P, Marbaix E, Debiève F, Jauniaux E.
27
28 Anomalies of the placenta and umbilical cord in twin gestations. *Am J*
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
Obstet Gynecol 2015; 213: S91-102.
12. Costa-Castro T, Zhao DP, Lipa M, Haak MC, Oepkes D, Severo M,
Montenegro N, Matias A, Lopriore E. Velamentous cord insertion in
dichorionic and monochorionic twin pregnancies - Does it make a
difference? *Placenta* 2016; 42: 87-92.
13. Suzuki S, Kato M. Clinical significance of pregnancies complicated by
velamentous umbilical cord insertion associated with other umbilical
cord/placental abnormalities. *J Clin Med Res* 2015; 7: 853-856.
14. Schachter M, Tovbin Y, Arieli S, Friendler S, Ron-El R, Sherman D. In
vitro fertilization is a risk factor for vasa previa. *Fertil Steril* 2002; 78:
642–643.

- 1 15. Jauniaux E, Englert Y, Vanesse M, Hiden M, Wilkin P. Pathologic
2 features of placentas from singleton pregnancies obtained by in vitro
3 fertilization and embryo transfer. *Obstet Gynecol* 1990; 76: 61–64.
4
5
6
7
- 8 16. Delbaere I, Goetgeluk S, Derom C, De Bacquer D, De Sutter P,
9 Temmerman M. Umbilical cord anomalies are more frequent in twins
10 after assisted reproduction. *Hum Reprod* 2007; 22: 2763-2767.
11
12
13
14
- 15 17. Baulies S, Maiz N, Muñoz A, Torrents M, Echevarría M, Serra B.
16 Prenatal ultrasound diagnosis of vasa praevia and analysis of risk
17 factors. *Prenat Diagn* 2007; 27: 595-599.
18
19
20
21
22
- 23 18. Jauniaux E, Ben-Ami I, Maymon R. Do assisted-reproduction twin
24 pregnancies require additional antenatal care? *Reprod Biomed Online*
25 2013; 26: 107-119.
26
27
28
29
- 30 19. Robinson BK, Grobman WA. Effectiveness of timing strategies for
31 delivery of individuals with vasa previa. *Obstet Gynecol* 2011; 117: 542-
32 549.
33
34
35
36
- 37 20. Fuchs F, Senat MV. Multiple gestations and preterm birth. *Semin Fetal*
38 *Neonatal Med* 2016; 21: 113-120.
39
40
41
- 42 21. Fung TY, Lau TK. Poor perinatal outcome associated with vasa previa: is
43 it preventable? A report of three cases and review of the literature.
44 *Ultrasound Obstet Gynecol* 1998; 12: 430-433.
45
46
47
48
- 49 22. Maymon R, Herman A, Jauniaux E, Frenkel J, Ariely S, Sherman D.
50 Transvaginal sonographic assessment of cervical length changes during
51 triplet gestation. *Hum Reprod* 2001; 16: 956-960.
52
53
54
55
56
- 57 23. Shulman A, Jauniaux E, Holmes A, Maschiach S, Maymon R. Vaginal
58
59
60
61
62
63
64
65

- 1 sonography of the cervix for the prediction of "time to delivery" in ART
2
3
4 twins gestations. *Twin Res* 2002; 5: 255-259.
- 5
6 24. Gianopoulos J, Carver T, Tomich PG, Karlman R, Gadwood K. Diagnosis
7
8 of vasa previa with ultrasonography. *Obstet Gynecol* 1987; 69: 488-491.
9
- 10
11 25. Raga F, Ballester MJ, Osborne NG, Bonilla-Musoles F. Role of color flow
12
13 Doppler ultrasonography in diagnosing velamentous insertion of the
14
15 umbilical cord and vasa previa. A report of two cases. *J Reprod Med*
16
17 1995; 40: 804-808.
18
19
- 20
21 26. Gandhi M, Cleary-Goldman J, Ferrara L, Ciorica D, Saltzman D,
22
23 Rebarber A. The association between vasa previa, multiple gestations,
24
25 and assisted reproductive technology. *Am J Perinatol* 2008; 25: 587-589.
26
27
- 28
29 27. Chmait RH, Chavira E, Kontopoulos EV, Quintero RA. Third trimester
30
31 fetoscopic laser ablation of type II vasa previa. *J Matern Fetal Neonatal*
32
33 *Med* 2010; 23: 459-462.
34
- 35
36 28. Nomiya M, Toyota Y, Kawano H. Antenatal diagnosis of velamentous
37
38 umbilical cord insertion and vasa previa with color Doppler imaging.
39
40 *Ultrasound Obstet Gynecol* 1998; 12: 426-429.
41
- 42
43 29. Lee W, Lee VL, Kirk JS, Sloan CT, Smith RS, Comstock CH. Vasa
44
45 previa: prenatal diagnosis, natural evolution, and clinical outcome. *Obstet*
46
47 *Gynecol* 2000; 95: 572-576.
48
- 49
50 30. Smorgick N, Tovbin Y, Ushakov F, Vaknin Z, Barzilay B, Herman A,
51
52 Maymon R. Is neonatal risk from vasa previa preventable? The 20-year
53
54 experience from a single medical center. *J Clin Ultrasound* 2010; 38:
55
56 118-122.
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
31. Bronsteen R, Whitten A, Balasubramanian M, Lee W, Lorenz R, Redman M, Goncalves L, Seubert D, Bauer S, Comstock C. Vasa previa: clinical presentations, outcomes, and implications for management. *Obstet Gynecol* 2013; 122: 352-357.
 32. Rebarber A, Dolin C, Fox NS, Klauser CK, Saltzman DH, Roman AS. Natural history of vasa previa across gestation using a screening protocol. *J Ultrasound Med* 2014; 33: 141-147.
 33. Swank ML, Garite TJ, Maurel K, Das A, Perlow JH, Combs CA, Fishman S, Vanderhoeven J, Nageotte M, Bush M, Lewis D; Obstetrix Collaborative Research Network. Vasa previa: diagnosis and management. *Am J Obstet Gynecol* 2016; 215: 223.e1-e6.
 34. Catanzarite V, Cousins L, Daneshmand S, Schwendemann W, Casele H, Adamczak J, Tith T, Patel A. Prenatally diagnosed vasa previa: A single-institution series of 96 cases. *Obstet Gynecol* 2016; 128: 1153-1161.
 35. Francois K, Mayer S, Harris C, Perlow JH. Association of vasa previa at delivery with a history of second-trimester placenta previa. *J Reprod Med* 2003; 48: 771-774.
 36. Ruiter L, Kok N, Limpens J, Derks JB, de Graaf IM, Mol BW, Pajkrt E. Systematic review of accuracy of ultrasound in the diagnosis of vasa previa. *Ultrasound Obstet Gynecol* 2015; 45: 516-522.
 37. Hasegawa J, Nakamura M, Sekizawa A, Matsuoka R, Ichizuka K, Okai T. Prediction of risk for vasa previa at 9-13 weeks' gestation. *J Obstet Gynaecol Res* 2011; 37: 1346-1351.
 38. Sepulveda W, Rojas I, Robert JA, Schnapp C, Alcalde JL. Prenatal

- 1 detection of velamentous insertion of the umbilical cord: a prospective
2 color Doppler ultrasound study. *Ultrasound Obstet Gynecol* 2003; 21:
3 564-569.
4
5
6
7
8 39. Suzuki S, Igarashi M, Inde Y, Miyake H. Abnormally shaped placentae in
9 twin pregnancy. *Arch Gynecol Obstet* 2010; 281: 65-69.
10
11
12 40. Gavriil P, Jauniaux E, Leroy F. Pathologic examination of placentas from
13 singleton and twin pregnancies obtained after in vitro fertilization and
14 embryo transfer *Pediatr Pathol* 1993; 13: 453-462.
15
16
17
18 41. van Steenis A, Zhao DP, Steggerda SJ, Kist WJ, Haak MC, Oepkes D,
19 Lopriore E. Double fatal outcome after ruptured vasa previa in
20 monozygotic twins: case report and review of the literature. *J Matern
21 Fetal Neonatal Med* 2016; 29: 2523-2526.
22
23
24
25 42. Papathanasiou D, Witlox R, Oepkes D, Walther FJ, Bloemenkamp KW,
26 Lopriore E. Monozygotic twins with ruptured vasa previa: double
27 trouble! *Fetal Diagn Ther* 2010; 28: 48-50.
28
29
30
31 43. Gillissen A, Sueters M, van Lith JM, Walther FJ, Lopriore E. Acute
32 hemorrhage in monozygotic twins with ruptured velamentous vessels:
33 anemic twin resuscitated by its co-twin through placental vascular
34 anastomoses? *Fetal Diagn Ther* 2013; 34: 56-58.
35
36
37
38 44. Zahumensky J, Gojis O, Kiss I, Kucera E. Velamentous insertion of the
39 umbilical cord of twin B as a cause of vasa previa in monozygotic
40 diamniotic twins. *Int J Gynaecol Obstet* 2013; 122: 261-262.
41
42
43
44 45. Cipriano LE, Barth WH Jr, Zaric GS. The cost-effectiveness of targeted
45 or universal screening for vasa praevia at 18-20 weeks of gestation in
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Ontario. BJOG 2010; 117: 1108-1118.

- 2
3
4 46. Melamed N, Pittini A, Hirsch L, Yogev Y, Korzeniewski SJ, Romero R,
5
6 Barrett J. Do serial measurements of cervical length improve the
7
8 prediction of preterm birth in asymptomatic women with twin gestations?
9
10 Am J Obstet Gynecol 2016;215:616.e1-616.e14.
11
12
13 47. Hofmeyr GJ, Barrett JF, Crowther CA. Planned caesarean section for
14
15 women with a twin pregnancy. Cochrane Database Syst Rev.
16
17 2015:CD006553.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure legends

Fig 1: Flow diagram showing the selection of reports included in the review.

Fig 2: Transvaginal ultrasound scans: A) showing a large VP (arrow) above the internal os of the cervix (Cx) on color mapping; B) with non-pulsatile flow on Doppler imaging (arrow). Consent was obtained from the patient for the use of these images.

Table 1. Prenatal ultrasound data and outcome of 7 cases of twin pregnancies

diagnosed with VP

Case No.	GA at diagnosis (wks + days)	Other US findings	Chorionicity-amnionicity	CL (mm) at last TVS (GA wks + days)	GA at delivery (wks + days)	Outcome Mode of delivery Fetal sex & weight (g)
1	28+2	None	DCDA	30 (32+3)	36+1	EICS M2810/M2850
2	29+1	None	DCDA	33 (32+5)	34+0	EICS F2026/F2030
3	29+2	VCI & SUA x2	DCDA	38 (34+2)	35+0	EICS M2040/M2240
4	22+0	VCI	DCDA	34 (35+5)	36+2	EICS F2270/M2370
5	29+2	VCI	DCDA	10 (32+6)	33+0	EmCS M1720/F1730
6	25+4	VCI & PP	DCDA	30 (28+2)	30+3	EmCS F1240/M1240
7	23+2	VCI	MCDA	30 (34+0)	35+1	EICS F2910/F2400

VCI = velamentous cord insertion; PP = placenta previa; SUA = single umbilical artery; DCDA = dichorionic-diamniotic; MCDA = monochorionic-diamniotic; EICS = elective caesarean section; EmCS = emergency caesarean section; F = female; M = male.

Table 2. Characteristics of the 13 reports included in the review

Author (year)	Type of study	No. of cases of VP diagnosed antenatally and confirmed at birth	No. of twin pregnancies with VP	Chorionicity
Raga <i>et al.</i> ²⁵ (1995)	Case report	2	1	DCDA
Nomiyama <i>et al.</i> ²⁸ (1998)	Cohort study of VCI (n=6)	2	1	DCDA
Lee <i>et al.</i> ²⁹ (2000)	Cohort study of VP	18	2	N/A
Francois <i>et al.</i> ³⁵ (2003)	Case-control study of VP	13	1	N/A
Oyelese <i>et al.</i> ⁶ (2004)	Multicenter cohort study of VP	155	7	N/A
Baulies <i>et al.</i> ¹⁷ (2007)	Cohort study of VP	9	2	2 DCDA
Ghandi <i>et al.</i> ²⁶ 2008	Case report	3	3	3 DCDA
Chmait <i>et al.</i> ²⁷ (2010)	Case report	2	1	MCDA
Smorgick <i>et al.</i> ³⁰ (2010)	Cohort study of VP	19	3	3 DCDA
Bronsteen <i>et al.</i> ³¹ (2013)	Cohort study of VP	56	7	N/A
Rebarber <i>et al.</i> ³² (2014)	Cohort study of VP	29	5	3 DCDA 2 MCDA
Swank <i>et al.</i> ³³ (2016)	Multicenter cohort study of VP	47	5	N/A
Catanzarite <i>et al.</i> ³⁴ (2016)	Cohort study of VP	92	18	6 DCDA 12 MCDA

VCI = velamentous cord insertion; VP= vasa previa; DCDA= dichorionic-diamniotic; MCDA = monochorionic-diamniotic; N/A= not available.

Table 3. Baseline perinatal characteristics of 12 VP cases included in the review

Author (year)	GA at diagnosis (wks)	Other US findings	Chorionicity-amnionity	CL (mm) at last TVS (wks)	GA at delivery (wks +days)	Outcome Mode of delivery Fetal sex & weight (g)
Raga <i>et al.</i> ²⁵ (1995)	34	VCI&LLP	DCDA	N/A	38	EICS M2810/F?
Nomiyama <i>et al.</i> ²⁸ (1998)	18	VCI	DCDA	N/A	31	EmCS F2026/F2030
Lee <i>et al.</i> ²⁹ (2000)	N/A N/A	N/A N/A	N/A N/A	N/A N/A	N/A 27+1	1 Stillbirth 1 post-natal death
Baulies <i>et al.</i> ¹⁷ (2007)	35 21	None VCI	DCDA DCDA	N/A N/A	36 34	EmCS F3050/M2210 EICS F1669/M1990
Ghandi <i>et al.</i> ²⁶ (2008)	29 18 19	None VCI&LLP VCI&LLP	DCDA DCDA DCDA	N/A N/A N/A	34 34 35	EICS EICS EICS
Chmait <i>et al.</i> ²⁷ (2010)	22	TTTS	MCDA	11 (30+5)	34+3	EmCS F2656/1240
Bronsteen <i>et al.</i> ³¹ (2013)	N/A N/A	N/A N/A	N/A N/A	25 (24) 15 (28)	27 31	1 post-natal death Stillbirth

GA = Gestational age; wks = weeks; VCI = velamentous cord insertion; LLP = low-lying placenta; SUA = single umbilical artery; DCDA = dichorionic-diamniotic; MCDA = monochorionic-diamniotic; TTTS = twin-twin transfusion syndrome; EICS = elective caesarean section; EmCS = emergency caesarean section; F = female; M = male; N/A = not available

Figure 1 Flow Chart




Figure 2A US
[Click here to download high resolution image](#)




Figure 2B US
[Click here to download high resolution image](#)

