# Mutations in noncoding regions of *GJB1* are a major cause of X-linked CMT

Pedro J. Tomaselli, MD, MSc\* Alexander M. Rossor, MRCP, PhD\* Alejandro Horga, MD, PhD Zane Jaunmuktane, FRCPath Aisling Carr, MRCP, PhD Paola Saveri, BSc Giuseppe Piscosquito, MD Davide Pareyson, MD Matilde Laura, MD, PhD Julian C. Blake, FRCP Roy Poh, PhD James Polke, PhD Henry Houlden, FRCP, PhD Mary M. Reilly, FRCP, FRCPI, MD

Correspondence to Dr. Reilly: m.reilly@ucl.ac.uk

# Supplemental data at Neurology.org

#### ABSTRACT

**Objective:** To determine the prevalence and clinical and genetic characteristics of patients with X-linked Charcot-Marie-Tooth disease (CMT) due to mutations in noncoding regions of the gap junction  $\beta$ -1 gene (*GJB*1).

**Methods:** Mutations were identified by bidirectional Sanger sequence analysis of the 595 bases of the upstream promoter region, and 25 bases of the 3' untranslated region (UTR) sequence in patients in whom mutations in the coding region had been excluded. Clinical and neurophysiologic data were retrospectively collected.

**Results:** Five mutations were detected in 25 individuals from 10 kindreds representing 11.4% of all cases of CMTX1 diagnosed in our neurogenetics laboratory between 1996 and 2016. Four pathogenic mutations, c.-17G>A, c.-17+1G>T, c.-103C>T, and c.-146-90\_146-89insT were detected in the 5'UTR. A novel mutation, c.\*15C>T, was detected in the 3' UTR of *GJB1* in 2 unrelated families with CMTX1 and is the first pathogenic mutation in the 3'UTR of any myelin-associated CMT gene. Mutations segregated with the phenotype, were at sites predicted to be pathogenic, and were not present in the normal population.

**Conclusions:** Mutations in noncoding DNA are a major cause of CMTX1 and highlight the importance of mutations in noncoding DNA in human disease. Next-generation sequencing platforms for use in inherited neuropathy should therefore include coverage of these regions. *Neurology*® 2017;88:1445-1453

#### GLOSSARY

**CMT** = Charcot-Marie-Tooth disease; **CMTX** = X-linked Charcot-Marie-Tooth disease; **CV** = conduction velocity; **Cx32** = connexin 32; **EVS** = Exome Sequencing Project; **ExAC** = Exome Aggregation Consortium; **GJB1** = gap junction  $\beta$ -1 gene; **IRES** = internal ribosomal entry site; **ORF** = open reading frame; **UTR** = untranslated region.

Mutations in the gap junction  $\beta$ -1 gene (*GJB1*) encoding the transmembrane channel protein, connexin 32 (Cx32), are the most common cause of X-linked Charcot-Marie-Tooth disease (CMTX) and the second commonest cause of Charcot-Marie-Tooth disease (CMT) overall.<sup>1</sup> The Cx32 protein is widely expressed in human tissues, including myelinating Schwann cells in the peripheral nervous system.<sup>2</sup> In the peripheral nervous system, Cx32 is found in the non-compact myelin of the paranodes and incisures, where it allows the movement of small molecules and ions between the multiple concentric myelinated layers of the Schwann cell and the axon membrane.<sup>2,3</sup> *GJB1* exists as 2 transcripts that are regulated by 2 tissue-specific promoters (P1 and P2), allowing differential expression of these transcripts in neuronal and non-neuronal tissue.<sup>4-6</sup> The transcriptional machinery in neuronal tissue requires the P2 promoter and other elements located in the 5' untranslated region (UTR) for efficient Cx32 expression. Mutations in the 5' UTR region have previously been described by our group and others as causative of

<sup>\*</sup>These authors contributed equally to this work.

From the MRC Centre for Neuromuscular Diseases (P.J.T., A.M.R., A.H., A.C., M.L., M.M.R.), Department of Neuropathology (Z.J.), and Department of Neurogenetics (R.P., J.P., H.H.), National Hospital for Neurology and Neurosurgery, UCL Institute of Neurology, Queen Square, London, UK; Clinic of Central and Peripheral Degenerative Neuropathies Unit (P.S., G.P., D.P.), Department of Clinical Neurosciences, IRCCS Foundation, C. Besta Neurological Institute, Milan, Italy; Department of Clinical Neurophysiology (J.C.B.), Norfolk and Norwich University Hospital, Norfolk, UK.

Go to Neurology.org for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article. The Article Processing Charge was paid by the Wellcome Trust and Medical Research Council.

This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (CC BY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Black symbols = affected; empty symbols = unaffected; dot symbols = affected by history; diagonal line = deceased; arrow = index case.

CMTX1 and have been shown to impair P2mediated transcription of  $GJB1.^7$  Mutations in the 3' UTR region are a rare cause of hereditary diseases overall; however, as this region often contains mRNA regulatory elements, mutations in the 3' UTR may affect normal translation.<sup>8</sup> In this study, we sought to determine the frequency and phenotype of CMTX1 due to mutations in the 5' and 3' UTR noncoding regions of GJB1.

**METHODS Patients.** Patients harboring mutations in the 5' and 3' UTR of *GJB1* were identified from the CMT database of the National Hospital for Neurology and Neurosurgery, Queen Square, London, United Kingdom. In some patients in whom Sanger sequencing of the coding region of *GJB1* was negative, further screening of the 5' and 3' UTR was performed because of

the strong clinical suspicion of CMT1X on the basis of a lack of male-to-male transmission, more severely affected males, and a predominantly demyelinating polyneuropathy. The remaining patients were identified from diagnostic samples submitted for testing of G/B1 to the neurogenetics laboratory of The National Hospital of Neurology and Neurosurgery after routine screening of both the coding and noncoding regions of G/B1 was adopted. The clinical and neurophysiologic data were collected retrospectively for all identified patients with mutations in the 5' and 3' UTR of G/B1.

**Statistical analysis.** Statistical analysis was performed using a 2tailed Student unpaired *t* test (Excel; Microsoft, Redmond, WA).

**Genetics analysis.** Genetic testing was performed in the National Hospital of Neurology and Neurosurgery Neurogenetics Laboratory. Additional targeted genetic testing was performed in selected cases (appendix e-1 at Neurology. org). Mutations were identified by bidirectional Sanger sequence analysis of *GJB1* including 595 bases upstream of the ATG start codon, the coding region, and 25 bases of 3' UTR

Table 1	Clinical	featur	es of pati	ients with X	linked Charcot-Ma	arie-Too	oth diseas	e due to mutati	ons in the 5′ and 3′ un	translated	d regions of GJ	81		
ID	Mutation	Sex	Age at onset, y	Age at evaluation, y	Presenting symptom	Split hand	APB/ FDIO/ ADM	Ankle dorsiflexion/ plantar flexion	Reflexes UL/LL	Plantars	Pinprick UL/LL	Vibration UL/LL	Other	CMTES
1-I.2	c17G>A	F	55	61	Unsteadiness	No	5/4+/5	4+/5	Present/ankles absent	↑ <b>/</b> ↓	N/2″ below knee	N/knee	Pain, deafness left ear, Horner syndrome, CSF protein 0.49, abnormal MRI	13
1-II.2	c17G>A	F	22	42	Unsteadiness	No	4/4/4-	4+/5	Brisk/knees brisk and ankles absent	Mute	Mid-forearm/ mid-calf	N/CM	Pain, scoliosis, Raynaud disease, normal MRI	14
1-II.4	c17G>A	F	30	45	Difficulty walking	Yes	3/2/4-	1/4	Brisk/ankles absent	Mute	Wrist/mid-calf	N/knee	Scoliosis, pain	10
1-III.3	c17G>A	М	8	24	Tremor and difficulty writing	Yes	4-/4-/4	2/5	Triceps reduced, biceps and supinator absent/ knees reduced and ankles absent	↓/↓	N/N	N/N	Tremor, pain	7
1-III.5	c17G>A	F	15	25	Difficulty walking and leg pain	No	4-/4-/4	4/5	Present/ankles absent	↓/↓	N/N	N/N	Tremor, pain	5
1-IV.2	c17G>A	F	8	16	Clumsiness and leg pain	No	5/5/5	5/5	Reduced/ankles absent	$\downarrow/\downarrow$	N/N	N/N	Pain	_
2-11.2	c17G>A	F		58	Asymptomatic	No	5/5/5	5/5	Present/present	$\downarrow/\downarrow$	N/N	N/N	-	0
2-III.1	c17G>A	М	18	33	Difficulty walking	Yes	3/4/4	4/5	Present/ankles absent	$\downarrow/\downarrow$	N/ankle	N/N	Tremor	8
2-111.3	c17G>A	М	26	34	Clumsy hands, difficulty walking	No	4/4+/4	4/5	Present/ankles absent	↑ <b>/</b> ↓	Mid-forearm/ mid-calf	N/ankle	Positive sensory symptoms, CSF protein 0.45	10
2-111.4	c17G>A	М	32	33	"Floppy" walking	Yes	3/4/4	2/4	Present/ankles absent	$\downarrow/\downarrow$	N/N	N/N	Normal MRI	-
3-111.2	c103C>T	М	<10	39	Foot drop	Yes	0/2/3	0/5	Present/ankles absent	↓/↓	2″ Above wrist/ below knee	Wrist on right and MCP on left/ASIS on right and CM on left	Tremor	_
3-111.5	c103C>T	М	15	37	Clumsiness	Yes	2/3/4	1/4-	Absent/absent	Mute	N/mid-foot	N/ankle	Abnormal MRI	9
4-111.1	c17G>A	М	<10	34	Frequent falls	Yes	1/4/4	3/5	Triceps reduced, biceps and supinator absent/ ankles absent	Mute	N/N	N/ankle	Tremor, pain	8
4-III.3	c17G>A	М	<10	32	Difficulty walking	Yes	2/4/4-	2/5	Present/ankles absent	?/↑	N/above ankle	N/ankle	Tremor	11
4-III.5	c17G>A	F		31	Asymptomatic "Family gait"	Yes	4/4/4	4/5	Present/right ankle reduced and left absent	↓/↓	N/mid-foot	N/N	Tremor	9
5-1.2	c.*15C>T	F		58	Asymptomatic	No	5/5/5	5/5	Present/present	$\downarrow/\downarrow$	N/N	N/N	-	0
5-II.1	c.*15C>T	F		29	Asymptomatic	No	5/5/5	5/5	Present/present	$\downarrow/\downarrow$	N/N	N/N	-	0
5-11.2	c.*15C>T	М	5	21	Difficulty walking	No	4-/4-/ 4-	3/5	Absent/ankles absent	↓/↓	N/N	N/N	Pain	15
6-III.1	c 17+1G>T	М	15	35	Difficulty walking	Yes	2/4/ <sup>a</sup>	0/5	Supinator absent/ absent	$\downarrow/\downarrow$	N/feet	N/knee	Normal MRI	13
6-111.2	c 17+1G>T	М	15	24	Difficulty walking	_	-	-	-	-	-	-	-	-

Table 1	Continu	ed												
₽	Mutation	Sex	Age at onset, y	Age at evaluation, y	Presenting symptom	Split hand	APB/ FDIO/ ADM	Ankle dorsiflexion/ plantar flexion	Reflexes UL/LL	Plantars	Pinprick UL/LL	Vibration UL/LL	Other	CMTES
7-111.1	c103C>T	Σ	$\stackrel{<}{\sim}$ 10	38	Difficulty walking	Yes	0/0/3	0/1	Present/ankles absent	Mute	Mid-palm/upper calf	N/right ASIS and left knee	1	16
8-II.1	c.146- 90_146- 89intT	Σ	<10	24	Difficulty walking	I	I	I	Absent/absent	1/1	I	I	I	I
9-IV.2	c17G>A	ш	12	30	Difficulty walking		4/4-/4	4-/5	Present/ankles absent	†/†	Fingers/N	N/ankle	Ι	00
10-111.3	c.*15C>T	ш		71	Asymptomatic	No	5/5/5	4+/5	Present/present	†/†	N/N	N/CM	I	4
10-IV.3	c.*15C>T	Σ	14	45	Poor balance	No	4/4/4	1/2	Absent/absent	$\uparrow/\uparrow$	N/ankle	N/ankle	I	18
Abbreviatio Marie-Tooth Split-hand v	וחs: ↑ = exter ר Examinatio vas defined	nsor pl in Scor bv the	antar res  ~e; FDIO = predomir	ponse; ↓ = fl€ = first dorsal ant clinical a	exor plantar respon: interosseous; LL = and electrophysiolo.	se; ADM = lower l aic invol	l = abduct imbs; MCI lvement o	or digiti minimi; , <sup>2</sup> = metacarpop f APB over ADM	APB = abductor pollicis halangeal; N = normal; 1 and FDIO in the hand.	brevis; A{ UL = upț	SIS = anterior su oer limbs.	perior iliac spine; CM =	costal margin; CMTES	= Charcot-

sequence. Conditions and primers are available in appendix e-1. In silico analysis was performed with the aid of AlamutVisual (Interactive Biosoftware, Rouen, France), which includes the splice-prediction tools SpliceSiteFinder-like, MaxEntScan, NNSPLICE, GeneSplicer, and Human Splicing Finder.

**Standard protocol approvals, registrations, and patient consents.** This study was approved by the research ethics committee of the National Hospital for Neurology and Neurosurgery. All patients consented to publication of their clinical details.

**RESULTS Demographics.** A total of 25 individuals from 10 kindreds with mutations in the 5' and 3' UTR of GIB1 were identified (figure 1), of whom 14 were male and 11 female. The age at onset was reported to be less than 10 years in 7 male participants and 1 female participant with a range of 5-32 years in male participants and 8-55 years in female participants. Four patients from family 1 (1-I.2, 1-II.2, 1-II.4, and 1-III.3) and all 4 patients from family 2 have been reported previously.9 There was no male-tomale transmission in any of the pedigrees. From 1996 to 2016, 194 patients with mutations in the open reading frame of GIB1 were identified. Mutations in the 5' and 3' UTR therefore represent 11.4% of patients with CMTX1 identified in our neurogenetics laboratory.

Clinical features. The clinical details of the cohort are summarized in table 1. The most common presenting complaint was difficulty walking. Male participants were more severely affected than female participants. The mean Charcot-Marie-Tooth Examination Score was  $6.30 \pm 5.31$ (range 0-14, n = 10) for female participants and  $11.5 \pm 3.81$  (range 7–18, n = 10), for male participants (p = 0.021). One woman (2-II.2, aged 58) harboring the c.-17G>A mutation and 3 women (5-I.2, 5-II.1, and 10-III.3, aged 58, 29, and 71, respectively) harboring the c.\*15C>T mutation were asymptomatic but all had abnormal nerve conduction studies (table 2). Atypical presentations in our cohort included the following: patient 1-I.2 from family 1 (c.-17G>A) presented with late-onset CMTX (age 55), unilateral deafness, and Horner syndrome, characterized by miosis and ptosis. The proband (1-II.2) had mild scoliosis and her son (1-III.3) presented at age 8 years with hand tremor and difficulty writing. Three patients (1-I.2, 2-III.3, and 4-III.3) from families 1, 2, and 4 harboring the c.-17G>A mutation had unilateral extensor plantar responses. Postural tremor was present in 5 out of 6 male participants and 2 out of 7 female participants harboring the c.-17G>A mutation. Twelve patients had a split-hand, characterized by disproportionate involvement of the abductor

<sup>a</sup> Data unavailable

Table 2	N	europhysiologic	data															
			Sensory n	erve action	potentials						Compound	I muscle act	tion potentia	ls				
			Median (F	III-W)	Ulnar (FV-	·W)	Radial (for	earm-W)	Sural (calf	-ankle)	Median SE	on APB	Ulnar SE d	on ADM	C peronea EDB	l SE on	P tibial SE	on AH
ID	Sex	Age at examination, y	Amp, μV (≥8)	CV, m/s (≥50)	Amp, μV (≥5)	CV, m/s (≥50)	Amp, μV (≥20)	CV, m/s (≥50)	Amp, μV (≥5)	CV, m/s (≥40)	Amp, mV (≥5)	CV, m/s (≥50)	Amp, mV (≥8)	CV, m/s (≥50)	Amp, mV (≥2.5)	CV, m/s (≥40)	Amp, mV (≥4)	CV, m/s (≥40)
1- 1.2	F	61	3	43.5	5	50.5	16	55	Abs	_	5.8	44	9.5	56	1.1	35	7.4	_
1-II.2	F	45	4	40.5	3	46.5	16	47	Abs	—	1.6	32	7.7	51	Abs	-	0.6	-
1-II.4	F	41	3	40	5	57	13	46	Abs	_	6.2	50	9.1	63	0.2	37	9.1	35
1-III.3	М	25	Abs	Abs	Abs	_	Abs	_	Abs	-	0.5	32	7.5	39	Abs	Abs	0.9	38
2-11.2	F	58	9.8	50	6.5	57.1	Abs	_	Abs	_	6.7	53.7	11.3	60.3	0.8	43.7	1.5	37.9
2-111.1	М	41	2.6	41.7	3.5	39.1	11	37.6	3	40.5	2.5	35.5	5.7	41.1	0.8	32.9	4.2	35.2
2-111.3	М	34	3	38.5	1	40.5	_	_	1	31.5	0.9	42	5.2	42	0.1	23	4.1	35
2-111.4	М	33	2.7	36.7	5.2	40	6	38.5	3	31.4	0.6	34.7	4.1	39.1	0.1	33.5	0.5	27.9
3-111.2	М	39	Abs	_	Abs	_	Abs	_	_	_	0.2	23	2.6	33	1	41	_	_
3-111.5	М	37	2	38	Abs	_	13	48	Abs	—	1.5	36	2.5	40	Abs	_	0.2	27
4-III.1	М	34	-	—	—	—	Abs	—	Abs	—	0.5	—	6.5	—	0.1	27	—	-
4-III.3	М	32	Abs	—	Abs	_	5	—	_	_	1.3	27	3.6	32	Abs	_	-	-
4-III.5	F	31	Abs	-	Abs	-	14	—	-	-	8.6	48	7.8	38	Abs	-	-	-
5-1.2	F	58	8	47	5	50	16	50	6	41	5.5	58	9.4	57	3.9	46	12	53
5-II.1	F	29	4	50	5	48	20	50	Abs	-	5.9	53	10.8	60	6.6	45	12.8	45
5-II.2	М	21	Abs	_	Abs	_	Abs	—	-	_	1.9	34	1.4	37	-	_	0.6	32
6-III.1	М	30	Abs	-	Abs	-	-	—	Abs	-	1.9	32.5	9.3	41.1	Abs	-	2.7	34.4
7-III.1	М	48	Abs	_	Abs	_	9	38	Abs	_	0.3	_	4.5	36	Abs	_	-	-
8-II.1	М	Unknown	-	-	-	—	-	-	-	—	1.3	30	-	-	-	-	-	-
9-IV.2	F	30	3	38	3	37	15	43	Abs	_	4.8	40	8.3	45	0.1	32	2.4	32
10-III.3	F	71	Abs	-	Abs	-	9	45.5	Abs	-	8.1	50	9.3	57	1.9	37	-	-
10-IV.3	М	45	Abs	_	Abs	_	Abs	_	Abs	_	5.3	38	3.4	35	0.1	22	_	_

Abbreviations: — = not recorded; Abs = absent; ADM = abductor digiti minimi; AH = abductor hallucis; Amp = amplitude; APB = abductor pollicis brevis; C = common; CV = conduction velocity; EDB = extensor digitorum brevis; FIII = finger III; ID = family/patient identification; P = posterior; SE = surface electrode; W = wrist.

pollicis brevis compared with the first dorsal interosseous and abductor digiti minimi muscles.

**Neurophysiology.** Nerve conduction studies were available in 22 individuals (13 male and 9 female) and in all cases demonstrated a motor and sensory neuropathy (table 2). In male participants, the mean ulnar motor nerve conduction velocity (CV) was  $37.78 \pm 3.43$  m/s (range 32–42 m/s), whereas in female participants, the mean ulnar CV was  $54.15 \pm 8.09$  m/s (range 38–63 m/s), p < 0.0001 (tables 2 and e-1). There was a discrepancy between the median and ulnar compound muscle action potentials' amplitude, with the former being significantly reduced compared to the latter (table e-1). This finding is in accordance with the clinical observation of the split-hand.<sup>10</sup>

**MRI.** Brain MRI was performed in 5 patients (3 male and 2 female) and revealed a lesion of the corpus callosum in 1 female patient (1-I.2) with no vascular risk factors or clinical features suggestive of multiple sclerosis (appendix e-1).

**Neuropathology.** Sural nerve biopsy was available from 2 patients (1-II.2 and 2-III.3) and revealed a significant reduction in myelinated nerve fiber density and thin myelin sheaths (appendix e-1). There were occasional regeneration clusters and mild endoneurial edema. There were no inflammatory cells. These findings are similar to coding GJB1 mutation patients.

Genetic analysis. Five distinct mutations in the 5' and 3' UTR of GIB1 were identified (table 1). The position of the mutations relative to the GJB1 open reading frame (ORF) region is shown in figure 2. The nomenclature used in this study is based on current recommendations of the Human Genome Variation Society (HGVS).<sup>11</sup> In table e-2, there is an overview of all mutations in the 5' and 3' UTR regions of GIB1 and the corresponding nomenclature based on counting directly from the ATG translation initiation codon, which has been previously used to describe a number of mutations. The previously reported mutations9,12-15 c.-103C>T and c.-17G>A were detected in 2 (3 and 7) and 4 unrelated families (1, 2, 4, and 9), respectively, and segregated with the phenotype in all family members tested. Three novel mutations were identified: c.-146-90\_-146-89insT in family 8, c.-17+1G>T in family 6, and c.\*15C>T in families 5 and 10. The genome conservation scores are shown in table e-3; they were assessed using PhiloP, which were accessed through the UCSC Genome Browser (GRCh37/hg19).16 These 3 novel mutations segregate with the phenotype and are predicted to be pathogenic using AlamutVisual (Interactive Biosoftware) software. They are not

present in the NHLBI Exome Sequencing Project (EVS), Exome Aggregation Consortium (ExAC), dbSNP, or the 1000 Genome (1000genomes) databases.<sup>17–20</sup> ExAC and EVS only include the ORF and  $\pm 50$  bp of intronic sequences. The 2 new variants within the 5' UTR region were not detected in 100 controls. The new variant in the 3' UTR is not present in ExAC. The scores of the in silico splicing analysis for c.\*15C>T are shown in table e-4.

**DISCUSSION** In this study, we describe 2 new pathogenic mutations in the 5' UTR and a likely pathogenic mutation in the downstream 3' UTR region of GJB1. The evidence for the pathogenicity of these mutations is largely indirect and based on a typical CMT X1 phenotype, segregation within family members, in silico splice prediction analysis, and for the 3' UTR mutation, the presence of the same mutation in an unrelated individual with the same phenotype. All patients included in this study had a clinical or neurophysiologic phenotype typical for CMTX1 due to mutations in the G/B1 ORF region, characterized by a slowly progressive, predominantly length-dependent neuropathy, in which male participants were more severely affected than female participants and with an earlier age at onset.<sup>21</sup> In male participants, the motor CVs were in the intermediate range and slower than in female participants, as has been described previously.<sup>22</sup> Evidence suggests that loss of Cx32 channel function is the underlying pathomechanism responsible for CMTX1 due to coding mutations in GJB1.23

The nerve-specific 5' UTR of GJB1 is located immediately upstream of the start codon, adjacent to the P2 promoter. The P2 promoter contains binding sites for the neuron-specific transcription factors SOX10 and EGR2 that strongly activate Cx32 expression in the peripheral nervous system.<sup>7,24</sup> EGR2 has 3 proposed binding sites (E1, E2, and E3) within the P2 promoter, whereas SOX10 has 2 P2 binding sites (S1 and S2) (figure 2). Several mutations located within the core of the S2 SOX10 binding site have previously been described, a number of which have been shown to impair SOX10-mediated transcription of G/B1, resulting in a significant reduction in Cx32 expression.7,12,25,26 The novel c.-146-90\_-146-89insT mutation is located within the E3 EGR2 binding site. The E2 and E3 binding sites of promoter P2 are responsible for the majority of EGR2-mediated transcriptions of GIB1.24 It is therefore likely that this mutation results in reduced Cx32 expression as observed for mutations within the SOX10 binding site.



(A) Structural organization of *GJB1*. (B) Base numbering at each junction between regions according to the Human Genome Variation Society. (C) *GJB1* has 2 tissue-specific promoters (P1 and P2) that are alternatively spliced. In liver and pancreas, *GJB1* transcription is driven via promoter 1 (P1) upstream of the noncoding exon, exon 1a, whereas in neural tissue it is driven via the nerve-specific promoter 2 (P2) upstream of noncoding exon 1b.<sup>4,6</sup> The P1- and P2-expressed mRNAs have different 5' untranslated regions (UTRs) but an identical open reading frame (ORF) region and 3' UTR. (D) The EGR2 (E1, E2, and E3) and SOX10 (S1 and S2) binding sites of the P2 promoter region that function synergistically to regulate Cx32 expression in the nervous system. <sup>a</sup> Variants included in this study. <sup>b</sup> Novel variants.

The c.-17G>A mutation was identified in 4 different families (1, 2, 4, and 9) and has previously been reported by our group.<sup>9</sup> This mutation is located in the last base of exon 1b, which is one of the most highly conserved bases in splice-site consensus sequences. In silico splice site analysis predicted that this mutation may reduce the efficiency of splicing at this intron/exon boundary, leading to the inclusion of intron 1 and a mutant transcript.<sup>9</sup> The second novel c.-17+1G>T mutation in family 6 affects the adjacent base and is predicted to be pathogenic by the same mechanism.

We identified the c.-103C>T mutation in 2 different families (3 and 7). This mutation has previously been reported in unrelated families.<sup>12,14,15,27</sup> It is located within exon Ib, downstream of the P2 promoter, and lies within the internal ribosomal entry site (IRES) of the peripheral nerve specific mRNA transcript. The mutation is predicted to prevent translation of *GJB1* mRNA.<sup>28</sup> Taken together, our study and previous reports provide strong evidence that the c.-103C>T is pathogenic. Of note, the c.-102G>A variant (reported as c.-458G>A), affecting the adjacent base, did not segregate in a large family with CMT, suggesting that not every variant of an IRES element is pathogenic.<sup>29</sup>

Despite being located in a less conserved region, the mutation in the 3' UTR region, c.15C>T\*, segregated in 2 unrelated families with a typical CMTX1 phenotype. This specific 3' UTR region contains sequences that are predicted to act as regulatory elements critical in Cx32 translational activation/ repression, mRNA stability, micro-RNA binding, and transcript localization.<sup>8</sup> Although our understanding of the role of these sequences is poor, in silico splicing analysis predicts that this variant may create a 5' donor splice site leading to aberrant splicing within the 3' UTR. This in turn may affect mRNA stability, leading to downregulation of GJB1 expression.

The clinical and electrophysiologic findings of the patients described in this article with point mutations within S2 SOX10 and E3 EGR2 are indistinguishable from patients with mutations in the ORF of

*GJB1*. Previous in vitro analysis of the c.-146-27T>C (c.-529T>C) mutation in the E3 region or deletion of the S2 region have demonstrated a partial loss of promoter activity.<sup>7,24</sup> The indistinguishable clinical phenotype of the patients included in this study from patients with complete loss of function mutations in *GJB1* suggests that the noncoding mutations described cause complete loss of function.

In this study, we describe 5 pathogenic mutations, 3 of which are novel, in noncoding regions of *GJB1*, which are predicted to result in loss of function by a combination of transcription factor binding, disruption of mRNA translation, and altered mRNA stability. The search for these noncoding mutations was largely driven by the recognition of the classical phenotype of CMTX1 in the absence of mutations in the coding regions of *GJB1*. The large number of noncoding mutations in *GJB1* (11.4% of our cohort of 219 *GJB1* patients) is of interest and highlights the importance of mutations in noncoding DNA in human disease and the need to include noncoding regions of *GJB1* in targeted inherited neuropathy gene panels.

The study also raises the possibility that these types of mutations may be a more frequent cause of other inherited neurologic conditions than has been previously appreciated including in the not infrequent situation where next-generation sequencing identifies a heterozygous mutation for a gene known to cause recessive disease. How frequent similar noncoding mutations will be in other hereditary neuropathies and other inherited neurologic diseases has yet to be determined.

#### AUTHOR CONTRIBUTIONS

Pedro J. Tomaselli: drafted the manuscript and acquired, analyzed, and interpreted data. Alexander M. Rossor: drafted the manuscript and acquired, analyzed, and interpreted data. Alejandro Horga: acquired, analyzed, and interpreted data and revised the manuscript. Zane Jaunmuktane: acquired data and revised the manuscript. Aisling Carr: acquired data and revised the manuscript. Paola Saveiri: acquired and analyzed data and revised the manuscript. Piscoasquito Giuseppe: acquired data and revised the manuscript. David Pareyson: acquired data and revised the manuscript. Matilde Laura: acquired data and revised the manuscript. Julian Blake: acquired data and analyzed the electrophysiologic studies. Roy Poh: acquired, analyzed, and interpreted data and revised the manuscript. James Polke: acquired, analyzed, and interpreted data and revised the manuscript. Henry Houlden: acquired, analyzed, and interpreted data and revised the manuscript. Mary M. Reilly: design and conceptualization of the study, interpreted data, revised the manuscript, and supervised the study.

#### STUDY FUNDING

M.M.R. is supported by the Medical Research Council (MRC, G0601943) and M.M.R., ML, A.M.R. and P.T. are supported by the National Institutes of Neurological Diseases and Stroke and office of Rare Diseases (U54NS065712). The INC (U54NS065712) is a part of the NCATS Rare Diseases Clinical Research Network (RDCRN). This research was also supported by the National Institute for Health Research University College London Hospitals Biomedical Research Centre and by the National Institute for Health Research Biomedical Research Centre at Great Ormond Street Hospital for Children NHS Foundation Trust and

University College London. A.M.R. is funded by a Welcome trust postdoctoral research training fellowship for clinicians. P.T. was supported by the Brazilian National Council for Scientific and Technological Development (CNPq).

#### DISCLOSURE

The authors report no disclosures relevant to the manuscript. Go to Neurology.org for full disclosures.

Received October 15, 2016. Accepted in final form January 18, 2017.

#### REFERENCES

- Murphy SM, Laura M, Fawcett K, et al. Charcot-Marie-Tooth disease: frequency of genetic subtypes and guidelines for genetic testing. J Neurol Neurosurg Psychiatry 2012;83:706–710.
- Scherer SS, Deschenes SM, Xu YT, Grinspan JB, Fischbeck KH, Paul DL. Connexin32 is a myelin-related protein in the PNS and CNS. J Neurosci 1995;15: 8281–8294.
- Balice-Gordon RJ, Bone LJ, Scherer SS. Functional gap junctions in the schwann cell myelin sheath. J Cell Biol 1998;142:1095–1104.
- Piechocki MP, Toti RM, Fernstrom MJ, Burk RD, Ruch RJ. Liver cell-specific transcriptional regulation of connexin32. Biochim Biophys Acta 2000;1491: 107–122.
- Suter U, Snipes GJ, Schoener-Scott R, et al. Regulation of tissue-specific expression of alternative peripheral myelin protein-22 (PMP22) gene transcripts by two promoters. J Biol Chem 1994;269:25795–25808.
- Neuhaus IM, Bone L, Wang S, Ionasescu V, Werner R. The human connexin32 gene is transcribed from two tissue-specific promoters. Biosci Rep 1996;16: 239–248.
- Houlden H, Girard M, Cockerell C, et al. Connexin 32 promoter P2 mutations: a mechanism of peripheral nerve dysfunction. Ann Neurol 2004;56:730–734.
- Chatterjee S, Pal JK. Role of 5'- and 3'-untranslated regions of mRNAs in human diseases. Biol Cell 2009;101: 251–262.
- Murphy SM, Polke J, Manji H, et al. A novel mutation in the nerve-specific 5'UTR of the GJB1 gene causes Xlinked Charcot-Marie-Tooth disease. J Peripher Nerv Syst 2011;16:65–70.
- Rossor AM, Polke JM, Houlden H, Reilly MM. Clinical implications of genetic advances in Charcot-Marie-Tooth disease. Nat Rev Neurol 2013;9:562–571.
- 11. Human Genome Variation Society Database (HGVS). Available at: hgvs.org. Accessed May 15, 2016.
- Ionasescu VV, Searby C, Ionasescu R, Neuhaus IM, Werner R. Mutations of the noncoding region of the connexin32 gene in X-linked dominant Charcot-Marie-Tooth neuropathy. Neurology 1996;47:541–544.
- Li M, Cheng TS, Ho PW, et al. -459C>T point mutation in 5' non-coding region of human GJB1 gene is linked to X-linked Charcot-Marie-Tooth neuropathy. J Peripher Nerv Syst 2009;14:14–21.
- Tsai PC, Chen CH, Liu AB, et al. Mutational analysis of the 5' non-coding region of GJB1 in a Taiwanese cohort with Charcot-Marie-Tooth neuropathy. J Neurol Sci 2013;332:51–55.
- Kabzinska D, Kotruchow K, Ryniewicz B, Kochanski A. Two pathogenic mutations located within the 5'-regulatory sequence of the GJB1 gene affecting initiation of

transcription and translation. Acta Biochim Pol 2011;58: 359–363.

- 16. UCSC Genome Browser. Available at: genome.ucsc.edu. Accessed May 15, 2016.
- NHLBI Exome Sequencing Project Database (EVS). Available at: evs.gs.washington.edu/EVS/. Accessed May 15, 2016.
- Exome Aggregation Consortium Database (ExAC). Available at: exac.broadinstitute.org. Accessed May 15, 2016.
- National Center for Biotechnology Information. Available at: ncbi.nlm.nih.gov/SNP. Accessed May 15, 2016.
- 20. 1000 Genomes Project. Available at: 1000genomes.org. Accessed May 15, 2016.
- Scherer SS, Kleopa KA. X-linked Charcot-Marie-Tooth disease. J Peripher Nerv Syst 2012;17(suppl 3):9–13.
- Birouk N, LeGuern E, Maisonobe T, et al. X-linked Charcot-Marie-Tooth disease with connexin 32 mutations: clinical and electrophysiologic study. Neurology 1998;50:1074–1082.
- Shy ME, Siskind C, Swan ER, et al. CMT1X phenotypes represent loss of GJB1 gene function. Neurology 2007;68: 849–855.
- Bondurand N, Girard M, Pingault V, Lemort N, Dubourg O, Goossens M. Human Connexin 32, a gap junction protein altered in the X-linked form of Charcot-Marie-

Tooth disease, is directly regulated by the transcription factor SOX10. Hum Mol Genet 2001;10:2783–2795.

- Beauvais K, Furby A, Latour P. Clinical, electrophysiological and molecular genetic studies in a family with X-linked dominant Charcot-Marie-Tooth neuropathy presenting a novel mutation in GJB1 Promoter and a rare polymorphism in LITAF/SIMPLE. Neuromuscul Disord 2006;16: 14–18.
- Sivera R, Sevilla T, Vilchez JJ, et al. Charcot-Marie-Tooth disease: genetic and clinical spectrum in a Spanish clinical series. Neurology 2013;81:1617–1625.
- Flagiello L, Cirigliano V, Strazzullo M, et al. Mutation in the nerve-specific 5'non-coding region of Cx32 gene and absence of specific mRNA in a CMTX1 Italian family: mutations in brief no. 195 [online]. Hum Mutat 1998; 12:361.
- Hudder A, Werner R. Analysis of a Charcot-Marie-Tooth disease mutation reveals an essential internal ribosome entry site element in the connexin-32 gene. J Biol Chem 2000;275:34586–34591.
- Bergmann C, Zerres K, Rudnik-Schoneborn S, Eggermann T, Schroder JM, Senderek J. Allelic variants in the 5' non-coding region of the connexin32 gene: possible pitfalls in the diagnosis of X linked Charcot-Marie-Tooth neuropathy (CMTX). J Med Genet 2002;39:e58.

## Seeking Established Neurologists! 2017 Transforming Leaders Program

Are you an experienced US AAN member neurologist 10 or more years out of residency looking for a one-of-a-kind leadership experience that will help you lead at the AAN, in daily life, and in the world of neurology? The Transforming Leaders Program may be for you! Apply by June 1, 2017, at *AAN.com/view/TransformingLeaders*.

### Registration Now Open for 2017 AAN Annual Meeting in Boston

Discover the Annual Meeting everyone's talking about. We changed the entire format and feel of the AAN Annual Meeting for 2016—and people took notice! We're doing it again in 2017, and adding even more changes to the docket. Register today to experience the value, choice, customization, and innovation that only an AAN Annual Meeting can offer. We look forward to seeing you April 22-28 in Boston. Register today at *AAN.com/view/AM17*.

# Neurology®

Mutations in noncoding regions of *GJB1* are a major cause of X-linked CMT Pedro J. Tomaselli, Alexander M. Rossor, Alejandro Horga, et al. *Neurology* 2017;88;1445-1453 Published Online before print March 10, 2017 DOI 10.1212/WNL.00000000003819

Updated Information & Services	including high resolution figures, can be found at: http://www.neurology.org/content/88/15/1445.full.html
Supplementary Material	Supplementary material can be found at: http://www.neurology.org/content/suppl/2017/03/10/WNL.0000000000 003819.DC1
References	This article cites 23 articles, 11 of which you can access for free at: http://www.neurology.org/content/88/15/1445.full.html##ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): <b>All Genetics</b> http://www.neurology.org//cgi/collection/all_genetics <b>EMG</b> http://www.neurology.org//cgi/collection/emg <b>Peripheral neuropathy</b> http://www.neurology.org//cgi/collection/peripheral_neuropathy
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://www.neurology.org/misc/addir.xhtml#reprintsus

#### This information is current as of March 10, 2017

*Neurology* <sup>®</sup> is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright Copyright © 2017 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

