

1 **Supplementary Material**

2 **Supplementary Text: Detailed Description of Underlying Models**

3 *The Statistical Model*

4
5
6
7 The statistical model was based on the database of the PREDICTS (Projecting
8 Responses of Ecological Diversity in Changing Terrestrial Systems) Project (Hudson
9 et al. 2017). These data describe the abundance (or for 17% of records only
10 occurrence) of species sampled at different levels of human pressure (generally
11 different land uses or land-use intensities), collated from numerous published
12 sources (or unpublished sources with a published methodology). This model did not
13 consider the effects of climate change. If sampling effort varied among the sites
14 sampled within a source dataset, abundance values were corrected by assuming
15 that recorded abundance increased linearly with increasing sampling effort (Newbold
16 et al. 2014). Land use at each of the sites in the PREDICTS database was classified
17 based on the description of the habitat, given in the source paper or provided by its
18 authors (see Supplementary Table 1 for criteria). Land use was classified as primary
19 vegetation, secondary vegetation, plantation forest, cropland, pasture or urban
20 (Hudson et al. 2014). This land-use classification is coarse, but was selected so that
21 the models could be generalized over large areas, and for correspondence with
22 available land-use projections. Importantly for this study, pasture describes sites
23 regularly or permanently grazed; sites with some grazing, but not sufficient to
24 substantially alter the habitat architecture, were classified as primary or secondary
25 vegetation (depending whether the natural habitat was destroyed historically). Sites
26 where fire occurs at natural frequency were classified as primary vegetation. In order
27 to understand the effects of small human disturbances – such as grazing or altered
28 fire regimes – within natural (primary and secondary) habitat, we distinguished
29 between minimally and substantially used natural vegetation. The latter incorporated
30 the ‘light’ and ‘intensive’ use-intensity classifications adopted in the PREDICTS
31 database (Hudson et al. 2014; Supplementary Table 1). The baseline for the models
32 and projections – minimally used primary vegetation – does not preclude some
33 human disturbances, of small extent or magnitude, and does not necessarily imply
34 the potential climax vegetation (for example where grassland is maintained naturally
35 through fire). This coarse abstraction of land use, and a degree of subjectivity in the
36 classification, will mean that some potentially important details are lost (land-use
37 classification is particularly challenging for grasslands); but this is necessary in order
38 to develop broad-scale models.

39 We developed a model of sampled species richness and sampled total
40 abundance, as a function of land use, using data from tropical grasslands and
41 savannas in Africa. The initial filtering of data was done by overlaying the global data
42 – extracted from the PREDICTS database on 29th September 2015 – onto a map of
43 biomes (<http://maps.tnc.org/>). We then manually checked each dataset and excluded
44 studies from biomes other than grassland and savanna. The resulting data
45 comprised 170,878 records, for 1,830 uniquely named taxa, from 922 sites. The sites
46 were distributed patchily, but showed a reasonable representation of land uses
47 (Supplementary Figure 2). We fitted generalized linear mixed-effects models with
48 land use as a single categorical fixed effect, and random effects representing the
49 identity of the source study – to capture the wide heterogeneity in sampled taxa,
50 sampling methods and sampling effort among studies – and the spatial blocking

51 structure of sampled sites within each study. Within-sample species richness was
52 modelled using a generalised linear mixed-effects model with a Poisson error
53 distribution. In this model we included an additional observation-level (i.e. site-level)
54 random intercept to control for overdispersion (Rigby et al. 2008). Within-sample
55 total abundance was log transformed and modelled using a linear mixed-effects
56 model (many abundance measurements were non-integers). To assess whether
57 biodiversity in tropical grassland biomes is responding to land use differently to
58 biodiversity globally, we also developed models using the global data across all
59 biomes (2.8 million records, for over 45,000 uniquely named taxa, from 17,064 sites;
60 Supplementary Figure 2). The global models of sampled species richness and total
61 abundance had exactly the same structure as the models for the tropical grassy
62 biomes.

63 To project the models of species richness and total abundance onto estimates
64 of current and future land-use patterns, we followed the methods in Newbold et al.
65 (2015). Mapped estimates of land use were taken from the harmonized land-use
66 data associated with the Representative Concentration Pathways scenarios (Hurt et
67 al. 2011). These data describe the proportion of each half-degree grid cell occupied
68 by the six above-named land-use classes. The model-estimated intactness of
69 biodiversity in each land use was applied, and then values averaged across the land
70 uses in each grid cell, weighted by proportional area. We used land-use estimates
71 for 2005 from the HYDE land-use reconstruction (see 'Land-use Scenarios', in the
72 main text, for more details), and for 2095 under the MESSAGE and MINICAM
73 Representative Concentration Pathways scenarios of future change in human
74 populations and socio-economic systems (Hurt et al. 2011). For more details, see
75 'Land-use Scenarios' in the main text.

76 77 *The Mechanistic Ecosystem Model*

78
79 In the Madingley general ecosystem model (Harfoot et al. 2014), organisms are
80 divided into functional groups: on land, divisions are made between trophic levels
81 (autotrophs, herbivores, omnivores and carnivores), between endotherms and
82 ectotherms, and between semelparous and iteroparous reproductive strategies.
83 Organisms are also characterized by their body mass (juvenile, adult and current
84 body masses). The model includes all photoautotrophs and all heterotrophs of body
85 mass $> 10^{-5}$ g. Ecological processes that involve animals act upon individual
86 organisms (although organisms in the same functional group and with similar
87 masses are grouped together for computational convenience; Purves et al. 2013).

88 The dynamics of plants are modelled using a terrestrial carbon model (Smith et
89 al. 2013), where plant biomass depends upon primary productivity, the division of
90 productivity between evergreen and deciduous plants, the allocation of productivity
91 to leaves, structural tissues or roots, and mortality, all of which depend upon climatic
92 variables (Smith et al. 2013). The plant model was chosen because future
93 projections of the driving climate variables are readily available. Non-climatic factors
94 important in shaping grassland/savanna plant dynamics, such as fire, are captured
95 implicitly to the extent that they correlate with climatic variables (Smith et al. 2013).
96 However, explicit representation of processes such as fire and phenology in future
97 might allow better predictions for grasslands and savannas (e.g. Scheiter and
98 Higgins 2009). The dynamics of heterotrophic animals are based on five ecological
99 processes: 1) predator-prey relationships (including herbivory), which are based on a
100 Holling's Type III functional response (Denno and Lewis 2012), and for predation on


101 a size-based model of predator-prey feeding preferences (Williams et al. 2010); 2)
102 metabolism, which is based on empirical relationships with temperature (Brown et al.
103 2004); 3) reproduction, which is based on a simple allocation of surplus mass to
104 reproductive potential followed by reproductive events once a threshold ratio of
105 reproductive potential to adult body mass is reached ; 4) mortality (in addition to
106 predation mortality), which is of three sources (a constant background rate,
107 starvation if insufficient food is obtained, and senescence, which follows the
108 Gompertz Model (e.g. Pletcher 1999) in assuming an exponentially increasing rate of
109 mortality with time after reproductive maturity is reached); and 5) dispersal, which in
110 the terrestrial realm is either random diffusive dispersal of juvenile organisms or
111 directed dispersal of organisms in response to starvation (Harfoot et al. 2014).

112 For simulating undisturbed ecosystems, the model takes as environmental
113 input the following variables, which determine the rates of ecological processes:
114 monthly near-surface air temperature, diurnal air temperature range, monthly
115 precipitation, soil water capacity, monthly number of frost days, and satellite-derived
116 net primary productivity (used only to determine seasonal patterns). We used a
117 monthly time step.

118 The model has been shown to capture observed properties of individual
119 organisms and the coarse structure of ecosystems reasonably well under
120 environmental conditions without human impact, especially in grassland ecosystems
121 (Harfoot et al. 2014). To simulate land-use impacts in the model, we removed from
122 the model plant biomass calculated as a certain proportion of net primary production,
123 following the Human Appropriation of Net Primary Production (HANPP) paradigm
124 (Haberl et al. 2007). We used published data on HANPP for the year 2000, compiled
125 based on statistics on permanent agricultural and forestry (excluding wood-fuel
126 harvesting), and estimates of global patterns in land use and soil degradation,
127 excluding the effects of shifting cultivation and vegetation loss from fire (important
128 limitations in its application to grassland/savanna systems; Haberl et al. 2007).
129 HANPP estimates are separated into two components: primary production lost as a
130 result of land-use conversion itself, and primary production lost directly to human
131 harvesting (Haberl et al. 2007). To project these estimates, we developed simple
132 spatial models of estimates of each of the two components of HANPP within each
133 half-degree grid cell within African tropical grassland biomes. Estimates of land-use
134 loss and harvest loss were modelled, using linear models, as a function of the total
135 areas of cropland, pasture and urban land use within each cell, and UN subregion (to
136 control for some of the socio-economic factors that might drive spatial differences in
137 human use of the land). Land-use data at half-degree resolution were taken from the
138 HYDE estimates in the harmonized land-use data of (Hurtt et al. 2011). UN
139 subregion was taken from (Sandvik 2009). Land-use areas were fitted with quadratic
140 polynomials and UN subregion as a factor. Land-use HANPP losses are right-
141 skewed but contain negative values (agricultural areas can be more productive than
142 the natural vegetation; Haberl et al. 2007); therefore, we employed a cube-root
143 transformation, preserving the sign of the values. Directly harvested losses of
144 HANPP are also right-skewed, but without negative values, and we therefore
145 employed a simple log transformation. These models explained a substantial
146 proportion of the estimated spatial variation in HANPP (R^2 values were 0.42 for land-
147 use losses and 0.53 for harvest losses). The spatial models of HANPP were then
148 applied to HYDE estimates of land use in 2005, and to estimates of land use in 2095
149 under two of the Representative Concentration Pathways scenarios (MINICAM and
150 MESSAGE; see 'Land-use Scenarios' in the main text).

151
152
153


Supplementary Tables and Figures




154
155
156

Supplementary Figure 1. The structure of this study.

(a)




(b)




157
158
159
160
161
162
163
164
165

Supplementary Figure 2. The location of sites (black points) used in the statistical model, from African grasslands and savannas (a) and worldwide (b). Grey shading in a shows the approximate extent of African grasslands and savannas. The 922 African grassland/savanna sites showed a reasonable representation of land uses: 231 in minimally used primary vegetation, 50 in substantially used primary vegetation, 36 in minimally used secondary vegetation, 33 in substantially used secondary vegetation, 378 in cropland, and 194 in pasture.




166
 167
 168
 169
 170
 171
 172

Supplementary Figure 3. Estimates of land-use change in African tropical grasslands. a-c: natural (primary and secondary) vegetation; d-f: human land uses (plantation forest, cropland, pasture and urban); g-i: human appropriation of net primary production (HANPP; sensu Haberl et al. 2007). Estimates are for 2005 (a, d, g), for 2100 under the MINICAM scenario (b, e, h), and for 2100 under the MESSAGE scenario (c, f, i).



173
 174
 175
 176
 177
 178
 179
 180

Supplementary Figure 4. Predicted temporal changes in the average total abundance of organisms and species richness in local ecological communities in African tropical grasslands and savannas. Shaded areas show $\pm 95\%$ confidence intervals. Land-use change estimates until 2005 were from the HYDE model (Klein Goldewijk et al. 2011). Future estimates were from the harmonized land-use estimates (Hurtt et al. 2011) for two of the Representative Concentration Pathways scenarios: MINICAM (blue) and MESSAGE (red).



181
 182 **Supplementary Figure 5.** Relationship between the area of pasture (a, b), cropland
 183 (b, c) and urban habitat, and the removal of net primary production from ecosystems
 184 by humans. Removal could either be caused directly by the conversion of land from
 185 natural vegetation to human use (a, c, e) or by the harvesting of vegetation by
 186 humans (b, d). The relationships (quadratic polynomials) were modelled using linear
 187 models (one model for land-use losses and one model for harvesting losses). UN
 188 sub-region (Sandvik 2009) was also included in the models to account for some of
 189 the spatial variation in the absolute magnitude of vegetation removal, caused by
 190 socio-economic differences among regions.
 191

192 **Supplementary Table 1.** The land-use and land-use-intensity classification criteria
193 (Hudson et al. 2014).

Level 1 Land Use	Predominant Land Use	Minimal use	Light use	Intense use
No evidence of prior destruction of the vegetation	Primary forest	Any human disturbances identified are very minor (e.g., a trail or path) or very limited in the scope of their effect (e.g., hunting of a particular species of limited ecological importance).	One or more human disturbances of moderate intensity (e.g., selective logging) or breadth of impact (e.g., bushmeat extraction), which are not severe enough to markedly change the nature of the ecosystem. Primary sites in suburban settings are at least Light use.	One or more human disturbances that is severe enough to markedly change the nature of the ecosystem; this includes clear-felling of part of the site too recently for much recovery to have occurred. Primary sites in fully urban settings should be classed as Intense use.
	Primary Non-Forest	As above	As above	As above
Recovering after destruction of the vegetation	Mature Secondary Vegetation	As for Primary Vegetation-Minimal use	As for Primary Vegetation-Light use	As for Primary Vegetation-Intense use
	Intermediate Secondary Vegetation	As for Primary Vegetation-Minimal use	As for Primary Vegetation-Light use	As for Primary Vegetation-Intense use
	Young Secondary Vegetation	As for Primary Vegetation-Minimal use	As for Primary Vegetation-Light use	As for Primary Vegetation-Intense use
	Secondary Vegetation (indeterminate age)	As for Primary Vegetation-Minimal use	As for Primary Vegetation-Light use	As for Primary Vegetation-Intense use
Human use (agricultural)	Plantation forest	Extensively managed or mixed timber, fruit/coffee, oil-palm or rubber plantations in which native understorey and/or other native tree species are tolerated, which are not treated with pesticide or fertiliser, and which have not been recently (< 20 years) clear-felled.	Monoculture fruit/coffee/rubber plantations with limited pesticide input, or mixed species plantations with significant inputs. Monoculture timber plantations of mixed age with no recent (< 20 years) clear-felling. Monoculture oil-palm plantations with no recent (< 20 years) clear-felling.	Monoculture fruit/coffee/rubber plantations with significant pesticide input. Monoculture timber plantations with similarly aged trees or timber/oil-palm plantations with extensive recent (< 20 years) clear-felling.
Human use (agricultural)	Cropland	Low-intensity farms, typically with small fields, mixed crops, crop rotation, little or no inorganic fertiliser use, little or no pesticide use, little or no ploughing, little or no irrigation, little or no mechanisation.	Medium intensity farming, typically showing some but not many of the following: large fields, annual ploughing, inorganic fertiliser application, pesticide application, irrigation, no crop rotation, mechanisation, monoculture crop. Organic farms in developed countries often fall within this category, as may high-intensity farming in developing countries.	High-intensity monoculture farming, typically showing many of the following features: large fields, annual ploughing, inorganic fertiliser application, pesticide application, irrigation, mechanisation, no crop rotation.
	Pasture	Pasture with minimal input of fertiliser and pesticide, and with low stock density (<i>not</i> high enough to cause significant disturbance or to stop regeneration of vegetation).	Pasture either with significant input of fertiliser or pesticide, or with high stock density (high enough to cause significant disturbance or to stop regeneration of vegetation).	Pasture with significant input of fertiliser or pesticide, <i>and</i> with high stock density (high enough to cause significant disturbance or to stop regeneration of vegetation).
Human use (urban)	Urban	Extensive managed green spaces; villages.	Suburban (e.g. gardens), or small managed or unmanaged green spaces in cities.	Fully urban with no significant green spaces.

195 **Supplementary References**

196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239

Brown, J. H. et al. 2004. Toward a metabolic theory of ecology. - Ecology 85: 1771–1789.

Denno, R. F. and Lewis, D. 2012. Predator-prey interactions. - In: Levin, S. A. et al. (eds), The Princeton Guide to Ecology. Princeton University Press, pp. 202–212.

Haberl, H. et al. 2007. Quantifying and mapping the human appropriation of net primary production in earth's terrestrial ecosystems. - Proc. Natl. Acad. Sci. U. S. A. 104: 12942–12947.

Harfoot, M. B. J. et al. 2014. Emergent global patterns of ecosystem structure and function from a mechanistic general ecosystem model. - PLoS Biol. 12: e1001841.

Hudson, L. N. et al. 2014. The PREDICTS database: a global database of how local terrestrial biodiversity responds to human impacts. - Ecol. Evol. 4: 4701–4735.

Hudson, L. N. et al. 2017. The database of the PREDICTS (Projecting Responses of Ecological Diversity In Changing Terrestrial Systems) project. - Ecol. Evol. 7: 145–188.

Hurttt, G. C. et al. 2011. Harmonization of land-use scenarios for the period 1500–2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands. - Clim. Change 109: 117–161.

Klein Goldewijk, K. et al. 2011. The HYDE 3.1 spatially explicit database of human-induced global land-use change over the past 12,000 years. - Glob. Ecol. Biogeogr. 20: 73–86.

Newbold, T. et al. 2014. A global model of the response of tropical and sub-tropical forest biodiversity to anthropogenic pressures. - Proc. R. Soc. London Ser. B, Biol. Sci. 281: 20141371.

Newbold, T. et al. 2015. Global effects of land use on local terrestrial biodiversity. - Nature 520: 45–50.

Pletcher, S. D. 1999. Model fitting and hypothesis testing for age-specific mortality data. - J. Evol. Biol. 12: 430–439.

Purves, D. W. et al. 2013. Time to model all life on Earth. - Nature 493: 295–297.

Rigby, R. A. et al. 2008. A framework for modelling overdispersed count data, including the Poisson-shifted generalized inverse Gaussian distribution. - Comput. Stat. Data Anal. 53: 381–393.

Sandvik, B. 2009. World Borders Dataset.
http://thematicmapping.org/downloads/world_borders.php.

Scheiter, S. and Higgins, S. I. 2009. Impacts of climate change on the vegetation of Africa: an adaptive dynamic vegetation modelling approach. - Glob. Chang. Biol. 15: 2224–2246.

Smith, M. J. et al. 2013. The climate dependence of the terrestrial carbon cycle, including parameter and structural uncertainties. - Biogeosciences 10: 583–606.

Williams, R. J. et al. 2010. The probabilistic niche model reveals the niche structure and role of body size in a complex food web. - PLoS One 5: e12092.