

Antimicrobial Agent	Isolates		
	Serotype a (n=29)	Serotype c (n=17)	All serotypes (n=56)
	Number (%)		
<b>Penicillin</b>	29 (100)	17 (100)	56 (100)
<b>Amoxicillin</b>	22 (75.9)	17 (100)	47 (83.9)
<b>Co-amoxiclav</b>	5 (17.2)	3 (17.6)	8 (14.2)
<b>Metronidazole</b>	29 (100)	17 (100)	56 (100)
<b>Clindamycin</b>	28 (96.6)	14 (82.4)	49 (87.5)
<b>Tetracycline</b>	2 (6.9)	1 (5.9)	5 (8.9)
<b>Ceftazidime</b>	20 (69.0)	15 (88.2)	43 (76.8)
<b>Ciprofloxacin</b>	0 (0)	0 (0)	0 (0)

Table 2

**Resistance phenotype of *A. actinomycetemcomitans* isolates from UK periodontitis patients.** Interpretive criteria for the HACEK group were applied for amoxicillin, amoxicillin/clavulanic acid, ceftazidime, ciprofloxacin and tetracycline; *Pasteurella multocida* criteria were used for penicillin; *Bacteroides fragilis* interpretive criteria were applied for clindamycin and metronidazole [18]