

Anti - inflammatory nutrition and successful ageing in elderly individuals: the multi-national MEDIS study

Stefanos Tyrovolas^{1,2}, Josep Maria Haro¹, Alexandra Foscolou², Dimitra Tyrovola², Anargiros Mariolis³, Vassiliki Bountziouka², Suzanne Piscopo⁴, Giuseppe Valacchi⁵, Foteini Anastasiou⁶, Efthimios Gotsis², George Metallinos², Natassa Papairakleous², Evangelos Polychronopoulos², Antonia – Leda Matalas², Christos Lionis⁶, Akis Zeimbekis⁷, Josep-Antoni Tur⁸, Labros S. Sidossis⁹, Demosthenes Panagiotakos²; for the MEDIS Study Group.

¹Parc Sanitari Sant Joan de Déu, Fundació Sant Joan de Déu, CIBERSAM, Universitat de Barcelona, Barcelona, Spain; ²Department of Nutrition and Dietetics, School of Health Science and Education, Harokopio University, Athens, Greece; ³Health Center of Aeropolis, General Hospital of Sparta, Aeropolis, Greece; ⁴University of Malta, Nutrition, Family and Consumer Studies Office, Msida, Republic of Malta; ⁵Department of Life Sciences and Biotechnology, University of Ferrara, Ferrara, Italy; ⁶Clinic of Social and Family Medicine, School of Medicine, University of Crete, Heraklion, Greece; ⁷Health Center of Kalloni, General Hospital of Mitilini, Mitilini, Greece; ⁸Research Group on Community Nutrition and Oxidative Stress, Universitat de les Illes Balears & CIBERObn, Guillem Colom Bldg, Campus, E-07122 Palma de Mallorca, Spain; ⁹Department of Exercise Science and Sport Studies, School of Arts and Sciences, Rutgers University, NJ, USA

Address for Correspondence

Prof. Demosthenes B. Panagiotakos
46 Paleon Polemiston St., Glyfada, 166 74, Attica, Greece
Tel. +30210-9603116 & Fax. +30210-9600719
E-mail: dbpanag@hua.gr

Abstract

Background: The role of diet and inflammation in successful ageing is not transparent, and as such, is still being investigated. The aim of the present work was to evaluate dietary habit's inflammatory potential in the successful ageing level of a random sample of older adults living in the Mediterranean basin and who participated in the MEDIS (MEDiterranean ISlands) study. **Methods:** During 2005-2016, 3,128 older adults (aged 65-100 years) from 24 Mediterranean islands and the rural Mani region (Peloponnesus) of Greece, were voluntarily enrolled in the study. A multidimensional successful ageing index consisting of 10 components was employed. A validated and reproducible Food Frequency Questionnaire (FFQ) was used to evaluate the dietary habits of the older adults. A nutrition anti - inflammatory (NAI) score based on the participants' specific dietary habits was assessed. **Results:** Participants with high NAI scores (pro-inflammatory nutrition) had a higher prevalence of hypercholesterolemia and lower levels of successful ageing. After adjusting for several confounders, the NAI score was associated with successful ageing [(95% CI): -0.03 (-0.5 to -0.006)]. Stratified analysis by gender and advanced age revealed heterogeneity in the NAI score, predicting successful ageing. **Conclusions:** Nutrition's inflammatory potential was reported as an important factor for successful ageing, suggesting that further research is needed on the role of anti- and pro- inflammatory dietary habits in healthy and successful ageing.

Keywords: successful ageing; dietary habits; dietary inflammation index; older adults, Mediterranean basin

Introduction

Ageing remains among the major public health challenges globally[1]. It was estimated that in 2015, the population over 65 years of age approached 1.5 billion[2]. In order to reduce the negative impact of ageing, great efforts are being put in to achieving healthy ageing. Healthy and successful ageing is considered as a low probability for disease and disability, high cognitive and physical capacity, as well as active participation in various social activities[3,4]. The aforementioned state of "successful ageing" is related to lower mortality rates[5] and better health outcomes[6]. The process of "healthy", "active" and "successful" ageing itself is quite complex and is associated with a variety of factors. Based on the aforementioned, the MEDIS investigators have created a successful ageing index (SAI) using 10 specific attributes. The MEDIS SAI reported to be multi-dimensional with 3 the following components: psycho-social-economic, the bio-clinical and the lifestyle [6] .

Determinants of successful ageing and specifically, the role of the diet's inflammatory potential on ageing remain poorly understood. Previously, well documented studies have reported a strong association between nutrition and inflammation marker's regulation [7-9]. The Mediterranean diet as a healthy dietary pattern[9], as well as flavonoids, polyphenols, n-3 fatty acids and other food components (i.e., coffee, tea, alcohol, etc) are well known for offering anti-inflammatory protection[10]. Moreover, it has been reported that the process of inflammation, oxidative stress and advancing age is strongly related to biological pathways that interacts with the dietary habits of older adults[11]. Recently Tyrovolas and

colleagues[12] reported that increased energy intake and a positive energy balance is associated with lower successful ageing levels. Although it is widely accepted that inflammation throughout atherosclerosis and endothelial dysfunction pathways[13] is associated with cardiovascular health and healthy ageing[14], there is a lack of evidence on the inflammatory role of the diet or specific food items in the ageing process.

Given the complexity of the successful ageing pathway and its interrelation with inflammation and oxidative stress, together with the lack of data among Mediterranean older populations, the aim of the present work was to evaluate the association between nutrition's inflammatory potential in the successful ageing level, of a random sample of older adults living in the Mediterranean basin and who participated in the MEDIS (MEDiterranean ISlands) study.

Methods

The MEDIS study sample

Between 2005-2016, a population-based, multi-national, convenience sampling was performed to voluntarily enroll older people from 24 Mediterranean islands from Greece, Republic of Cyprus, Malta, Italy and Spain [6]. Study's male gender % was 49.9%. According to the study's protocol, individuals were not eligible for inclusion if they resided in assisted-living centers, had a clinical history of cardiovascular disease (CVD) or cancer, or had lived away from the island for a considerable period of time during their lives (i.e., >5 years). These exclusion criteria were applied because the study aimed to assess lifestyle habits that were not subject to modifications due to existing chronic health conditions or by environmental factors, other than living milieu. A group of health scientists (physicians, dietitians and nurses) with experience in field investigation collected all the required information using a quantitative questionnaire and standard procedures.

The study followed the ethical considerations provided by the World Medical Association (52nd WMA General Assembly, Edinburgh, Scotland, October 2000). The Institutional Ethics Board of Harokopio University approved the design and procedures of the study (reference No. 16/19-12-2006). Participants were informed about the aims and procedures of the study and gave their consent prior to being interviewed.

Evaluation of clinical characteristics

All the measurements taken in the different study centres were standardized and the questionnaires were translated in all the cohorts' languages following the World Health Organization (WHO) translation guidelines for tools assessment[15]. Weight, height and waist circumference were measured using a standard protocol; body mass index (BMI) was calculated as the ratio of weight by height squared (kg/m^2). Overweight was defined as BMI between 25 and 29.9 Kg/m^2 and obesity was defined as BMI > 29.9 Kg/m^2 . Diabetes mellitus (type 2) was determined by fasting plasma glucose tests and was analyzed in accordance with the American Diabetes Association diagnostic criteria (glycated haemoglobin $\text{A1C} \geq 6.5$ or fasting blood glucose levels greater than 125 mg/dl or 2-h plasma glucose > 200 mg/dl during an oral glucose tolerance test-OGTT- or a random plasma glucose > 200 mg/dl, or by a prior diagnosis of diabetes). Participants who had blood pressure levels $\geq 140/90$ mmHg or used antihypertensive medications were classified as hypertensive. Fasting blood lipid levels (HDL-, LDL-cholesterol and triglycerides) were also recorded and hypercholesterolemia was defined as total serum cholesterol levels >200 mg/dL or the use of lipid-lowering agents according to the NCEP ATP III guidelines[16]. Symptoms of depression during the previous month were assessed using the validated Greek version (also translated in all the cohort's languages) of the shortened, self-report Geriatric Depression Scale (GDS) (range 0-20)[17]. We classified participants with GDS score >11 in the group of severe depressive symptoms, while those with GDS <10 in the group of mild and low depressive symptoms.

Evaluation of dietary habits, socio-demographic and other lifestyle characteristics of the participants

Dietary habits were assessed through a semi-quantitative, validated and reproducible food-frequency questionnaire[18]. The frequency of consumption of various food groups and beverages (i.e., meat and products, fish and seafood, milk and other dairy, fruits, vegetables, greens and salads, legumes, cereals, coffee and tea and soft-drinks) on daily, weekly or monthly basis, was assessed. Furthermore, intake of various alcoholic beverages (i.e., wine, beer, etc.) was measured in terms of wineglasses adjusted for ethanol intake (e.g., one 100 ml glass of wine was considered to have 12% ethanol). Also, the consumption of various types of coffee drinks (i.e., boiled, filtered, espresso and instant), as well as tea consumption (i.e., green, black) was measured in terms of general consumption per day and as well as cups per day. Energy and macronutrients intake was evaluated through the quantification of the portions of foods and beverages consumed, using food composition tables[19,20].

The calculation of a nutrition anti-inflammatory (NAI) score on the MEDIS study sample was based on previous literature[21,22]. Based on the validated MEDIS-FFQ, a total of 7 food components

were used for the development of a nutrition-related inflammatory score. These were: energy balance, protein, carbohydrates, total fat as percent of energy intake (EI), tea, coffee and alcohol consumption. The creation of a nutrition related inflammatory score was decided in order to avoid the co-linearity phenomenon between the aforementioned components that is essential for correctly performing a multivariable statistical analysis[23]. Individual ratings (-1 and 1) in each of the 7 components were assigned, according to their high anti-inflammatory or low anti-inflammatory effect respectively[24,25]. Higher NAI scores imply lower anti-inflammatory nutrition, following a similar coding methodology that has been reported in the literature [21]. If energy balance was >0 then the coding 1 was given while if it was <0 was given the score -1. If protein, carbohydrates and total fat consumption were within the recommended intakes[26] the score -1 was coded, while if it was over or lower the recommended intake the score 1 was coded. Tea, coffee and alcohol consumption was coded as -1 while no consumption was coded as 1. Moreover, the tertiles of the NAI score were computed, for its classification as high, medium and low anti-inflammatory nutrition.

Socio-demographic characteristics, such as age, gender, years of school, financial status and lifestyle characteristics, such as smoking habits and physical activity status, were recorded. Regarding financial status, the participants were asked to report their mean income during the previous three years using a four-point scale (low, inadequate to cover daily expenses = 1, medium, trying hard to cover daily expenses = 2, good, adequate to cover daily expenses = 3, very good, very adequate to cover daily expenses = 4); this scale was decided upon because of the variety of the populations studied, as well as the common difficulty of accessing exact financial data. The participants that were in the upper category were classified as participants with high financial status while all the others were classified as low and medium financial status (high vs. low-medium financial status). Current smokers were defined as smokers at the time of the interview. Former smokers were defined as those who had previously smoked, but had not done so for a year or more. The remaining participants were defined as occasional or non-current smokers. Physical activity was evaluated in MET-minutes per week, using the shortened, translated in all the cohort's languages and validated in Greek version of the self-reported International Physical Activity Questionnaire (IPAQ)[27]. Minimally active or "health-enhancing physical activity (HEPA) active" were classified as individuals who reported at least 3 MET-minutes per week. Furthermore, the weekly frequency of physical activity was recorded. Additionally, in order to evaluate social participation, the weekly frequency of their social activities with their family, their friends as well as their yearly frequency of excursions were recorded.

Further details about the MEDIS study protocol may be found elsewhere[28].

Evaluated outcomes

Following the multi-dimensional approach to successful ageing that has already been reported by the MEDIS study group[6], 10 components (i.e., education, financial status, physical activity, BMI, GDS score, participation in social activities with friends, with family, yearly excursions, CVD risk factors score and MedDietScore) were incorporated for the measurement of successful ageing. The successful ageing index composed was represented as the cumulative score of the 10 components (theoretical range 0-10); specifically, individual ratings (from 0 to 1) in each of the 10 components were assigned, according to their positive or negative (i.e., reverse scoring) influence on successful ageing.

Statistical analysis

Continuous variables are presented as mean \pm standard deviation (SD) and categorical variables as frequencies. Comparisons of continuous variables between groups were performed using the independent samples t-test (for normal distribution) and the Mann-Whitney U-test (for skewed distribution). Associations between categorical variables were tested using the chi-square test. Linear regression models were applied in order to evaluate the association between various socio-demographic, bio-clinical and nutritional factors (independent variables) and the level of successful ageing (dependent outcome). Colinearity was tested using the Variance Inflation Factor criterion (VIF; values >4 suggested colinearity between independent variables and one of them was excluded from the model). The assumption of homoscedasticity was tested by plotting the scatter plot of standardised residuals over the predicted score values. Results from linear regression models are presented as *b*-coefficients and their 95% Confidence Intervals. All reported *p*-values were based on two-sided tests. SPSS software (version 20) was used for all calculations (IBM Statistics, Greece).

Results

The total range of the NAI score of the MEDIS sample is -7.0 to +5.0. In the entire sample, the mean NAI score was -1.2 ± 2.3 . Demographic, behavioral, clinical and lifestyle characteristics of the total sample, categorized by NAI score tertiles, are summarized in **Table 1**. Compared to the 1st tertile (high anti-inflammatory nutrition), the participants in the 2nd and the 3rd tertile (medium and low anti-inflammatory nutrition) were of female gender ($p=0.001$), less urban residents ($p<0.001$), less physically active

($p < 0.001$), less smokers ($p = 0.03$) and they had higher prevalence of hypercholesterolemia ($p = 0.007$) and lower level of successful ageing ($p < 0.001$). Moreover, no differences were observed between groups of NAI score with regards to obesity, education level, financial status, diabetes mellitus and hypertension.

Table 1. Demographic, behavioral, clinical and lifestyle characteristics of the Multi-national MEDIS sample, nutrition anti-inflammatory (NAI) score categorization.

	NAI score	1 st tertile (1056)	2 st tertile (770)	3 st tertile (713)	P*
N					
Age (years)		74.4±7.3	73.5.1±7.5	73.7±8.1	0.03
Sex (males)		55	53	38	0.001
Urban residence (%)		64	63	53	<0.001
Obesity (%)		32	31	33	0.86
Education (years of school)		6.8±3.9	6.9±3.9	6.4±4.1	0.11
High financial status (%)		20	19	18	0.79
Living alone (%)		23	29	27	0.02
Current smoking (%)		17	17	12	0.03
Physical activity (%)		52	38	34	<0.001
Hypertension (%)		63	62	64	0.75
Diabetes Mellitus (%)		20	23	25	0.09
Hypercholesterolemia (%)		45	52	51	0.007
Successful ageing (0-10)		2.7±1.3	2.6±1.2	2.3±1.3	<0.001

* Comparisons of continuous variables between groups were performed using the independent samples t-test (for normal distribution) and the Mann-Whitney U-test (for skewed distribution).

After adjusting for age, gender, urban residence, waist circumference and smoking habits, it was found a consistent relation between NAI score and the level of successful ageing (*model 1 to model 3*). Specifically it was found that an increase in NAI score (low anti-inflammatory nutrition) was associated with lower levels of successful ageing [b-coefficient (95% CI): -0.03 (-0.5 to -0.006)], ($p = 0.014$). Then, the interactions between NAI and gender, age-group were tested. Although no significant interaction with gender and NAI was observed (p for interaction = 0.28), the analysis was splitted by sex, considering the fact of gender paradox in exploring healthy ageing [...]. The analysis revealed that in females, a low anti-inflammatory nutrition was inversely associated with the successful ageing score [-0.03 (-0.64 to -0.003)], ($p = 0.003$); however in males, there was no association between the NAI score and the level of successful ageing ($p = 0.27$).

A significant interaction was observed with age and NAI (p for interaction = 0.02). Thus, stratified by age-group analysis showed that the lower the anti-inflammatory dietary habits of an older individual, the lower the level of successful ageing score. This association was significant in the group of octogenarians [-0.05 (-0.09 to -0.001), $p = 0.04$], but not in the rest of the participants [-0.05 (-0.09 to -0.001), $p = 0.04$].

Finally the association between anti-inflammatory nutrition and health care services use was also tested. After adjusting for various confounders such age, sex, etc, it was found no relation between NAI score and annual health care services ($p = 0.76$)

Discussion

The present work revealed an inverse association between low anti-inflammatory nutrition, and successful ageing; in particular, the multi-adjusted analysis revealed that the consumption of a low anti-inflammatory diet was associated with lower levels of successful ageing, irrespective of the age, gender, residence area, smoking habits and waist circumference. In addition, when the analysis was stratified by gender and age group, this association remained significant only for females and in those over 80 years old. These relationships have rarely being studied globally; moreover, it is the first time that an association of this kind has been reported among older adults of the Mediterranean basin.

Despite the lack of previous findings regarding successful ageing and nutrition-related inflammation among Mediterranean populations, a number of studies have previously reported that inflammation process is highly related with cardiovascular disease (CVDs)[13,29]. Well documented studies have reported that inflammatory markers such as interleukin-1, CRP and others are related with CVD health[6,30]. In supporting of this, our unadjusted analysis reported that the more pro-inflammatory dietary habits of an individual, the higher was the prevalence for hypercholesterolemia while it seem to be a trend for the presence of diabetes mellitus. Previous results from the MEDIS study reported that the SAI

levels of the ageing insular Mediterranean population is low, with high cardiovascular morbidity. Moreover female gender and urban place of residence were inversely related with the MEDIS SAI [6]. In addition, the use of the SAI based only on the state of well being has been tested in a previous paper of MEDIS group (i.e., modified SAI was inversely associated with BMI, hypertriglyceridemia and was positively associated with tea consumption ($p \leq 0.001$)); however the external validity of this modified type of successful ageing score focused only on the well being was poor {couldn't predict the annual use of health care services [b-coefficient (p value): 1.10, ($p=0.155$)]} [6].

It has been suggested that living longer is a process that reflects complex interrelations among multi-dimensional factors [31]. A wider picture of this concept is described throughout the successful ageing theory which has been associated with better health outcomes[3-5]. Data analysis revealed that the lower the anti-inflammatory nutrition the lower the successful ageing level among the older individuals. The role of healthy diet in ageing and longevity has been well documented in the past [11,14,32] and well conducted studies have shown that healthy nutrition is associated with reduced overall mortality[32]. Nutrition's inflammatory potential have been shown to be associated with various health risk factors such as CVD risk factors burden [33], glucose intolerance, dyslipidemia, components of the metabolic syndrome as well as with anthropometric measurements and established cardiovascular disease [34], [35], [36].

Table 2. Results from linear regression (b-coefficient, 95%CI) performed to evaluate the association of various socio, bio-clinical, lifestyle characteristics and nutrition anti-inflammatory (NAI) score in association to the level of successful ageing (n=1701).

<i>Overall</i>	<i>Model 1</i>		<i>Model 2</i>		<i>Model 3</i>			
	<i>b</i>	95%CI	<i>b</i>	95%CI	<i>b</i>	95%CI		
NAI score (per 1)	-0.05	-0.7, -0.2	-0.03	-0.5, -0.004	-0.03	-0.5, -0.006		
Age (per 1 year)	-	-	-0.02	-0.025, -0.01	-0.01	-0.02, -0.009		
Sex (males vs. females)	-	-	0.58	0.48, 0.68	0.45	0.32, 0.57		
Urban residence (yes vs. no)	-	-	-0.02	-0.12, 0.08	-0.10	-0.22, 0.13		
Waist circumference (per 1 cm)	-	-	-	-	-0.29	-0.03, -0.02		
Current smoking habits (yes vs. no)	-	-	-	-	0.18	0.001, 0.37		
<i>Sub-group analysis</i>	Females		Males		65-80 yrs		80 yrs+	
	<i>b</i>	95%CI	<i>b</i>	95%CI	<i>b</i>	95%CI	<i>b</i>	95%CI
NAI score (per 1)	-0.03	-0.06,- 0.003	-0.02	-0.06, 0.017	-0.02	-0.05, 0.006	-0.05	-0.09, -0.001
Age (per 1 year)	-0.01	-0.02, -0.003	-0.02	-0.03, -0.009	-	-	-	-
Sex (males vs. females)	-	-	-	-	0.48	0.34,0.63	0.27	0.03, 0.51
Urban residence (yes vs. no)	-0.005	-0.16, 0.15	-0.23	-0.12, -0.06	-0.07	-0.20, 0.07	-0.16	-0.39, 0.08
Waist circumference (per 1 cm)	-0.03	-0.03, -0.02	-0.03	-0.04,-0.02	-0.03	-0.4, -0.02	-0.024	-0.03, -0.016
Current smoking habits (yes vs. no)	0.6	0.23, 0.96	0.04	-0.17, 0.26	0.27	0.07, 0.47	-0.19	-0.72, 0.33

In addition to this, other studies have reported the relationship between dietary habits and low grade inflammation[8,9]. It has been illustrated that a nutrition centered around meat consumption or away from the Mediterranean diet is highly related with inflammation[9,10,37]. On the contrary, a diet centered around vegetable or fruit consumption is inversely associated with inflammatory markers[38,39]. Moreover it is well known that high adherence to the traditional Mediterranean dietary pattern has been associated with lower inflammation levels in the human body[40]. Previous studies on dietary habits inflammatory potential and various inflammatory surrogates (i.e., CRP, TNF- α , IL-6, homocysteine, etc) have reported significant associations[41-43]. Inflammation, oxidative stress and healthy ageing are strongly related pathways that are interrelated by the older individual's nutrition[11,12]. However the exact relationship of ageing, healthy eating and the human body's low grade inflammation status still remains unclear[12]. Antioxidant nutrition components (such as, flavonoids, polyphenols, vitamins) as well as specific fatty acids, have been proposed among others as potential nutrition mechanisms that could beneficially contribute to the favorable level of inflammation markers[25,44].

To further explore the association between successful ageing and nutrition's-inflammatory potential, an additional multivariate analysis stratified by sex and age group was applied. This association between the anti-inflammatory nutrition and successful ageing levels remained significant only for the

female individuals and for the octogenarians group. Various well-documented global studies have reported that females live longer with higher morbidity rates, compared to males. Since the role of different trajectories in healthy ageing has also been described recently, the reported result of an anti-inflammatory nutrition, could also possibly explain the different gender morbidity levels[6]. Advanced age has been related with more plasma and/or serum levels of inflammatory mediators (i.e., cytokines, acute phase proteins, etc) as well as with other mediating inflammation factors, such as ageing body composition changes, sex hormones, asymptomatic infections, Alzheimer's disease[44], and the applied analysis reported a significant association between holistic nutrition's inflammatory potential and successful ageing, in the octogenarians. All of the above studies support the proposed relationships between successful ageing, and the NAI score in older individuals, with specific heterogeneity among different population groups. Due to the complexity of ageing, identifying possible anti-inflammatory nutrition-related determinants is of major importance to inform public health authorities and enhance public health nutrition planning.

Strengths and Limitations

The present study has several strengths. To the best of our knowledge, it is the first study that evaluated the effect of the diet's inflammatory potential in the successful ageing of a large sample of "healthy", free-living older people in the Mediterranean basin. The limitation of the study is mainly caused by its cross-sectional design and causal relationships cannot be inferred. Moreover, people with history of cancer and cardiovascular disease have been excluded from the sampling; therefore, potential selection bias may exist. Also, the cumulative successful ageing index that was previously developed by simply adding the presence of the common determinants of the individuals may not accurately estimate the successful ageing status. This methodology however, was based on a standard procedure described in the literature and has previously been used in other ageing-associated definitions (i.e., frailty, healthy ageing)[3,6]. For the development of NAI, 7 semi-quantitative nutrition components were used, and thus, the application of discrete and not continuous coding was obligatory; a fact that may limit the presented results.

Conclusion

The present work revealed the anti-inflammatory role of nutrition on the level of successful ageing among older Mediterranean adults. It is of major interest, to understand the role of nutrition and its inflammatory dynamic on the transforming nature of ageing. The gender and older age-related heterogeneity regarding successful ageing and nutrition's inflammatory potential should be taken into account by public health authorities, in order to develop nutritional education programs and health promotion actions targeted to the needs of specific population subgroups. However, further exploration is needed in order to understand how the anti-inflammatory potential of nutrition is interrelated with the ageing pathway.

Acknowledgement

We are particularly grateful to the men and women from the islands of Malta, Sardinia, Sicily, Mallorca, Menorca, Cyprus, Gökçeada, Lesvos, Samothraki, Crete, Corfu, Lemnos, Zakynthos, Cephalonia, Naxos, Syros, Ikaria, Salamina, Kassos, Rhodes, Karpathos, Tinos, Ai Stratis, Spetses and the rural area of Mani, who participated in this research. We also wish to express our gratitude to the MEDIS Study Group: M. Tornaritis, A. Polystipioti, M. Economou, (field investigators from Cyprus), K. Gelastopoulou, I. Vlachou (field investigator from Lesvos), I. Tsiligianni, M. Antonopoulou, N. Tsakountakis, K. Makri (field investigators from Crete), E. Niforatu, V. Alpentzou, M. Voutsadaki, M. Galiatsatos (field investigators from Cephalonia), K. Voutsas, E. Lioliou, M. Miheli (field investigator from Corfu), Tyrovolas S, G. Pounis, A. Katsarou, E. Papavenetiou, E. Apostolidou, G. Papavassiliou, P. Stravopodis (field investigators from Zakynthos), E. Tourloukis, V. Bountziouka, A. Aggelopoulou, K. Kaldaridou, E. Qira, (field investigators from Syros and Naxos), A. Foscolou (field investigator from Tinos), D. Tyrovolas (field investigators from Kassos), I. Protopappa (field investigator from Ikaria), C. Prekas, O. Blaserou, K.D. Balafouti (field investigators from Salamina), S. Ioakeimidi (field investigator from Rhodes and Karpathos), A. Foscolou, E. Paka, P. Drepanidis (field investigators from Gökçeada), A. Foscolou, P. Drepanidis, K. Katsana, S. Iossifidis (field investigators from Spetses), A. Mariolis (field investigator from Mani), S. Piscopo (field investigator from Malta), J.A. Tur (field investigator from Mallorca and Menorca), G. Valacchi, B. Nanou (field investigators from Sardinia and Sicily) for their substantial assistance in the enrolment of the participants.

Conflict of Interest: All authors declare that there is no conflict of interest.

Funding: The Study was funded by Research grants from the Hellenic Heart Foundation, the Graduate Program of the Department of Nutrition & Dietetics, Harokopio University and the Rutgers University, NJ, USA (GA #5884). Stefanos Tyrovolas's work was supported by the Foundation for Education and European Culture (IPEP), the Sara Borrell postdoctoral program (reference no. CD15/00019 from the Instituto de Salud Carlos III (ISCIII - Spain) and the Fondos Europeo de Desarrollo Regional (FEDER). Josep A. Tur was funded by grants PI11/01791, CIBERobn CB12/03/30038, and CAIB/EU 35/2001.

References

- 1 European Commission: The demographic future of Europe – from challenge to opportunity. Brussels:EC, 2006,
- 2 United Nations. World Population Ageing 1950–2050. 2010 R, DESA, United Nations:
- 3 Bowling A, Dieppe P: What is successful ageing and who should define it? *BMJ (Clinical research ed)* 2005;331:1548-1551.
- 4 Rowe JW, Kahn RL: Successful aging. *The Gerontologist* 1997;37:433-440.
- 5 Schoenfeld DE, Malmrose LC, Blazer DG, Gold DT, Seeman TE: Self-rated health and mortality in the high-functioning elderly--a closer look at healthy individuals: MacArthur field study of successful aging. *Journal of gerontology* 1994;49:M109-115.
- 6 Tyrovolas S, Haro JM, Mariolis A, Piscopo S, Valacchi G, Tsakountakis N, Zeimbekis A, Tyrovola D, Bountziouka V, Gotsis E, Metallinos G, Tur JA, Matalas AL, Lionis C, Polychronopoulos E, Panagiotakos D: Successful aging, dietary habits and health status of elderly individuals: a k-dimensional approach within the multi-national MEDIS study. *Experimental gerontology* 2014;60:57-63.
- 7 Akbaraly TN, Shipley MJ, Ferrie JE, Virtanen M, Lowe G, Hamer M, Kivimaki M: Long-term adherence to healthy dietary guidelines and chronic inflammation in the prospective Whitehall II study. *The American journal of medicine* 2015;128:152-160.e154.
- 8 Dias JA, Wirfalt E, Drake I, Gullberg B, Hedblad B, Persson M, Engstrom G, Nilsson J, Schiopu A, Fredrikson GN, Bjorkbacka H: A high quality diet is associated with reduced systemic inflammation in middle-aged individuals. *Atherosclerosis* 2015;238:38-44.
- 9 Koloverou E, Panagiotakos DB, Pitsavos C, Chrysohoou C, Georgousopoulou EN, Grekas A, Christou A, Chatzigeorgiou M, Skoumas I, Tousoulis D, Stefanadis C: Adherence to Mediterranean diet and 10-year incidence (2002-2012) of diabetes: correlations with inflammatory and oxidative stress biomarkers in the ATTICA cohort study. *Diabetes/metabolism research and reviews* 2016;32:73-81.
- 10 Esposito K, Marfella R, Ciotola M, Di Palo C, Giugliano F, Giugliano G, D'Armiento M, D'Andrea F, Giugliano D: Effect of a mediterranean-style diet on endothelial dysfunction and markers of vascular inflammation in the metabolic syndrome: a randomized trial. *Jama* 2004;292:1440-1446.
- 11 Willcox DC, Willcox BJ, Todoriki H, Suzuki M: The Okinawan diet: health implications of a low-calorie, nutrient-dense, antioxidant-rich dietary pattern low in glycemic load. *Journal of the American College of Nutrition* 2009;28 Suppl:500s-516s.
- 12 Tyrovolas S, Haro JM, Mariolis A, Piscopo S, Valacchi G, Makri K, Zeimbekis A, Tyrovola D, Bountziouka V, Gotsis E, Metallinos G, Tur JA, Matalas A, Lionis C, Polychronopoulos E, Panagiotakos D: The Role of Energy Balance in Successful Aging Among Elderly Individuals: The Multinational MEDIS Study. *Journal of aging and health* 2015;27:1375-1391.
- 13 Hansson GK: Inflammation, atherosclerosis, and coronary artery disease. *The New England journal of medicine* 2005;352:1685-1695.
- 14 Mathers JC: Nutrition and ageing: knowledge, gaps and research priorities. *The Proceedings of the Nutrition Society* 2013;72:246-250.
- 15 WHO: World Health Organization Translation Guidelines., 2016,
- 16 Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, And Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel III). *Jama* 2001;285:2486-2497.
- 17 Fountoulakis KN, Tsolaki M, Iacovides A, Yesavage J, O'Hara R, Kazis A, Ierodiakonou C: The validation of the short form of the Geriatric Depression Scale (GDS) in Greece. *Aging (Milan, Italy)* 1999;11:367-372.
- 18 Tyrovolas S, Pounis G, Bountziouka V, Polychronopoulos E, Panagiotakos DB: Repeatability and validation of a short, semi-quantitative food frequency questionnaire designed for older adults living in Mediterranean areas: the MEDIS-FFQ. *Journal of nutrition for the elderly* 2010;29:311-324.
- 19 U.S. Department of Agriculture ARS UNDLUNNDfSR, 2009, Release 22, 2016,
- 20 tables. Gfc, 2001, 2015,
- 21 Shivappa N, Steck SE, Hurley TG, Hussey JR, Hebert JR: Designing and developing a literature-derived, population-based dietary inflammatory index. *Public health nutrition* 2014;17:1689-1696.

- 22 Georgousopoulou EN, Kouli GM, Panagiotakos DB, Kalogeropoulou A, Zana A, Chrysohoou C, Tsigos C, Tousoulis D, Stefanadis C, Pitsavos C: Anti-inflammatory diet and 10-year (2002-2012) cardiovascular disease incidence: The ATTICA study. *International journal of cardiology* 2016;222:473-478.
- 23 Tu YK, Clerehugh V, Gilthorpe MS: Collinearity in linear regression is a serious problem in oral health research. *European journal of oral sciences* 2004;112:389-397.
- 24 Zaveri NT: Green tea and its polyphenolic catechins: medicinal uses in cancer and noncancer applications. *Life sciences* 2006;78:2073-2080.
- 25 Zampelas A, Panagiotakos DB, Pitsavos C, Chrysohoou C, Stefanadis C: Associations between coffee consumption and inflammatory markers in healthy persons: the ATTICA study. *The American journal of clinical nutrition* 2004;80:862-867.
- 26 Academies IoMotN: Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids (Macronutrients) Washington, DC, National Academies Press, 2005.
- 27 Papathanasiou G, Georgoudis G, Papandreou M, Spyropoulos P, Georgakopoulos D, Kalfakakou V, Evangelou A: Reliability measures of the short International Physical Activity Questionnaire (IPAQ) in Greek young adults. *Hellenic journal of cardiology : HJC = Hellenike kardiologike epitheorese* 2009;50:283-294.
- 28 Tyrovolas S, Polychronopoulos E, Bountziouka V, Zeimbekis A, Tsiligiani I, Papoutsou S, Gotsis E, Metallinos G, Lionis C, Panagiotakos DB: Level of adherence to the Mediterranean diet among elderly individuals living in Mediterranean islands: nutritional report from the Medis Study. *Ecology of food and nutrition* 2009;48:76-87.
- 29 Ross R: Atherosclerosis--an inflammatory disease. *The New England journal of medicine* 1999;340:115-126.
- 30 Ridker PM, Hennekens CH, Buring JE, Rifai N: C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *The New England journal of medicine* 2000;342:836-843.
- 31 Dupre ME, Liu G, Gu D: Predictors of longevity: evidence from the oldest old in China. *American journal of public health* 2008;98:1203-1208.
- 32 Keys A, Menotti A, Aravanis C, Blackburn H, Djordevic BS, Buzina R, Dontas AS, Fidanza F, Karvonen MJ, Kimura N, et al.: The seven countries study: 2,289 deaths in 15 years. *Preventive medicine* 1984;13:141-154.
- 33 Tyrovolas S, Koyanagi A, Kotsakis GA, Panagiotakos D, Shivappa N, Wirth MD, Hebert JR, Haro JM: Dietary inflammatory potential is linked to cardiovascular disease risk burden in the US adult population. *International journal of cardiology* 2017
- 34 Neufcourt L, Assmann KE, Fezeu LK, Touvier M, Graffouillere L, Shivappa N, Hebert JR, Wirth MD, Hercberg S, Galan P, Julia C, Kesse-Guyot E: Prospective Association Between the Dietary Inflammatory Index and Cardiovascular Diseases in the SUPplementation en Vitamines et Mineraux Antioxydants (SU.VI.MAX) Cohort. *Journal of the American Heart Association* 2016;5:e002735.
- 35 Ruiz-Canela M, Zazpe I, Shivappa N, Hebert JR, Sanchez-Tainta A, Corella D, Salas-Salvado J, Fito M, Lamuela-Raventos RM, Rekondo J, Fernandez-Crehuet J, Fiol M, Santos-Lozano JM, Serra-Majem L, Pinto X, Martinez JA, Ros E, Estruch R, Martinez-Gonzalez MA: Dietary inflammatory index and anthropometric measures of obesity in a population sample at high cardiovascular risk from the PREDIMED (PREvencion con DIeta MEDiterranea) trial. *The British journal of nutrition* 2015;113:984-995.
- 36 Garcia-Arellano A, Ramallal R, Ruiz-Canela M, Salas-Salvado J, Corella D, Shivappa N, Schroder H, Hebert JR, Ros E, Gomez-Garcia E, Estruch R, Lapetra J, Aros F, Fiol M, Serra-Majem L, Pinto X, Babio N, Gonzalez JI, Fito M, Martinez JA, Martinez-Gonzalez MA: Dietary Inflammatory Index and Incidence of Cardiovascular Disease in the PREDIMED Study. *Nutrients* 2015;7:4124-4138.
- 37 Lopez-Garcia E, Schulze MB, Fung TT, Meigs JB, Rifai N, Manson JE, Hu FB: Major dietary patterns are related to plasma concentrations of markers of inflammation and endothelial dysfunction. *The American journal of clinical nutrition* 2004;80:1029-1035.
- 38 Esmailzadeh A, Kimiagar M, Mehrabi Y, Azadbakht L, Hu FB, Willett WC: Fruit and vegetable intakes, C-reactive protein, and the metabolic syndrome. *The American journal of clinical nutrition* 2006;84:1489-1497.
- 39 Root MM, McGinn MC, Nieman DC, Henson DA, Heinz SA, Shanely RA, Knab AM, Jin F: Combined fruit and vegetable intake is correlated with improved inflammatory and oxidant status from a cross-sectional study in a community setting. *Nutrients* 2012;4:29-41.
- 40 Salas-Salvado J, Garcia-Arellano A, Estruch R, Marquez-Sandoval F, Corella D, Fiol M, Gomez-Gracia E, Vinales E, Aros F, Herrera C, Lahoz C, Lapetra J, Perona JS, Munoz-Aguado D,

Martinez-Gonzalez MA, Ros E: Components of the Mediterranean-type food pattern and serum inflammatory markers among patients at high risk for cardiovascular disease. *European journal of clinical nutrition* 2008;62:651-659.

41 Cavicchia PP, Steck SE, Hurley TG, Hussey JR, Ma Y, Ockene IS, Hebert JR: A new dietary inflammatory index predicts interval changes in serum high-sensitivity C-reactive protein. *The Journal of nutrition* 2009;139:2365-2372.

42 Shivappa N, Hebert JR, Marcos A, Diaz LE, Gomez S, Nova E, Michels N, Arouca A, Gonzalez-Gil E, Frederic G, Gonzalez-Gross M, Castillo MJ, Manios Y, Kersting M, Gunter MJ, De Henauw S, Antonios K, Widhalm K, Molnar D, Moreno L, Huybrechts I: Association between dietary inflammatory index and inflammatory markers in the HELENA study. *Molecular nutrition & food research* 2016

43 Shivappa N, Hebert JR, Rietzschel ER, De Buyzere ML, Langlois M, Debruyne E, Marcos A, Huybrechts I: Associations between dietary inflammatory index and inflammatory markers in the Asklepios Study. *The British journal of nutrition* 2015;113:665-671.

44 Krabbe KS, Pedersen M, Bruunsgaard H: Inflammatory mediators in the elderly. *Experimental gerontology* 2004;39:687-699.