



Who pays for sex? An analysis of the increasing prevalence of female commercial sex contacts among men in Britain

H Ward, C H Mercer, K Wellings, K Fenton, B Erens, A Copas and A M Johnson

Sex. Transm. Inf. 2005;81:467-471
doi:10.1136/sti.2005.014985

Updated information and services can be found at:
<http://sti.bmjournals.com/cgi/content/full/81/6/467>

These include:

References

This article cites 17 articles, 7 of which can be accessed free at:
<http://sti.bmjournals.com/cgi/content/full/81/6/467#BIBL>

Rapid responses

You can respond to this article at:
<http://sti.bmjournals.com/cgi/eletter-submit/81/6/467>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Topic collections

Articles on similar topics can be found in the following collections

[HIV Infection/AIDS](#) (1153 articles)
[Sexually Transmitted Infections](#) (1179 articles)

Notes

To order reprints of this article go to:
<http://www.bmjournals.com/cgi/reprintform>

To subscribe to *Sexually Transmitted Infections* go to:
<http://www.bmjournals.com/subscriptions/>

SEX WORK

Who pays for sex? An analysis of the increasing prevalence of female commercial sex contacts among men in Britain

H Ward, C H Mercer, K Wellings, K Fenton, B Erens, A Copas, A M Johnson

Sex Transm Infect 2005;**81**:467–471. doi: 10.1136/sti.2005.014985

See end of article for authors' affiliations

Correspondence to: Helen Ward, DIDE, Imperial College London, Norfolk Place, London W2 1PG, UK; h.ward@imperial.ac.uk

Accepted for publication 21 March 2005

Background: In the United Kingdom the incidence of sexually transmitted infections (STI) and risky sexual behaviours is increasing. The role of commercial sex in this trend is poorly understood. Little is known about the men who pay for sex. We examined the epidemiology of female commercial sex contacts reported by men in 1990 and 2000.

Methods: National probability sample surveys of sexual attitudes and lifestyles (Natsal) of men aged 16–44 resident in Britain in 1990 (n = 6000) and 2000 (n = 4762).

Results: The proportion of men who reported paying women for sex in the previous 5 years increased from 2.0% (95% CI 1.6 to 2.5) in 1990 to 4.2% in 2000 (95% CI 3.6 to 4.9). In both surveys, paying for sex was more frequent in men aged between 25 years and 34 years, who were never or previously married, and who lived in London. There was no association with ethnicity, social class, homosexual contact, or injecting drug use. Men who paid for sex were more likely to report 10 or more sexual partners in the previous 5 years; only a minority of their lifetime sexual partners (19.3%) were commercial. They were more likely to meet partners abroad and to report previous STI. Only 15% reported having had an HIV test.

Conclusion: The proportion of men who reported paying for heterosexual sex has increased, and these men have multiple commercial and non-commercial partners. Their higher rates of STI and low level of HIV testing suggest the need for prevention interventions for clients as well as sex workers.

The transmission dynamics of HIV and other sexually transmitted diseases (STI) in the population are fundamentally dependent on rates of partner change, levels of condom use and sexual mixing patterns, particularly between groups with different levels of risk.¹ In some epidemics women who sell sex have been highly vulnerable to HIV infection at an early phase, as have their sexual partners, and many interventions have targeted sex workers in an attempt to reduce transmission.²

Over the past two decades a large body of research on sex workers has built up across the world, showing the variations in background, organisation of sex work, risks of infection, and levels of safety.³ Far less is known about the men who pay for sex. There are some population estimates, mainly from developing countries, where men who pay for sex are seen as a potential “bridge” between higher and lower risk populations. They may have a key role in early stages of an epidemic, particularly where paying for sex is frequent.^{4–6} Interventions targeting this behaviour as well as promoting condom use are said, based on broad trends, to have contributed to control.^{7–9} There are fewer data from industrialised countries and the role of commercial sex in transmission of infection is far from clear. Men who pay for sex may be at increased risk of acquiring and transmitting infections, but as a group are hard to define and therefore difficult to target with health promotion initiatives.

Currently in the United Kingdom there is a resurgence of bacterial STI, with a 103% increase in chlamydia and a 97% increase in gonorrhoea diagnoses from 1997 to 2002, along with an increase in new cases of HIV infection.¹⁰ The underlying causes are not fully understood, but are thought to relate to an increase in high risk sexual behaviours.¹¹ The role of commercial sexual contact in these epidemics is unknown, but data from women who sell sex in the United Kingdom suggest that they are unlikely to be central to transmitting infection since they have a declining incidence of STI and high levels of condom use.¹²

The aim of this paper is to examine the epidemiology of female commercial sex contacts reported by men in Britain in 1990 and 2000. Specifically, we estimated the proportion of men who report paying women for sex at these two points in time, and explore the relation between paying for sex and a number of sociodemographic, sexual, and health behaviour factors.

METHODS

Natsal 2000 is a stratified probability sample survey of 11 161 men and women aged 16–44 years, resident in private households in Britain, who were interviewed between May 1999 and February 2001 using a combination of face to face and computer assisted self interview (CASI). The response rate was 65.4%. Details of the methodology are published elsewhere.^{11–13}

Results from Natsal 2000 were compared with those from respondents aged 16–44 years in Natsal 1990. Natsal 1990 used face to face interview and pen and paper self completion. Comparisons are based on questions with identical wording in the two surveys. All behavioural variables were collected by self completion (pen and paper in Natsal 1990, CASI, in Natsal 2000). Natsal 1990 data were weighted for differential selection probabilities and post-stratified to the 1991 census estimates, following the procedures outlined for Natsal 2000.^{14–15}

In both surveys respondents were asked questions about their sexual lifestyles and attitudes including questions for men about their experience of paying women for sex. Men were asked: “Have you ever paid money for sex with a woman?” If respondents reported “yes,” then they were asked “When was the last time you paid money for sex with a woman?” Response options were: in the last 7 days, between 7 days and 4 weeks ago, between 4 weeks and 1 year ago,

Abbreviations: CASI, computer assisted self interview; STD, sexually transmitted diseases; STI, sexually transmitted infections

Table 1 Prevalence of reporting paying for heterosexual sex

Time frame	Natsal 1990	Base (weighted, unweighted)	Natsal 2000	Base (weighted, unweighted)	OR adjusted for age for 1990–2000 change in prevalence (95% CI)*
Ever	% (95% CI)		% (95% CI)		
	5.6 (4.9 to 6.3)	6678, 5766	8.8 (7.9 to 9.7)	5613, 4698	1.54 (1.30 to 1.83)
	No of different women paid†	329, 302	No of different women paid†	490, 475	
	Mean (SD)		Mean (SD)		–
	3.4 (4.8)		3.6 (7.9)		
Past 5 years	Median		Median		
	2		2		
	Range		Range		
	39 (max 40)		149 (max 150)		–
	% (95% CI)		% (95% CI)		
Past 5 years	2.0 (1.6 to 2.5)	6675, 5764	4.2 (3.6 to 4.9)	5614, 4699	2.09 (1.61 to 2.73)
Past year	% (95% CI)		% (95% CI)		
	0.5 (0.3 to 0.7)	6675, 5764	1.3 (1.0 to 1.7)	5613, 4698	2.70 (1.62 to 4.49)

Base: All men aged 16–44 years.

* $p < 0.0001$ for all age adjusted odds ratios (OR).

†Of men reporting to have ever paid for heterosexual sex.

between 1 year and 5 years ago, longer than 5 years ago. Men were then asked “In your lifetime, to about how many different women, altogether, have you paid money for sex?” In Natsal 2000 these questions were asked as part of a CASI component; in the previous survey there were asked in a self completed pen and paper questionnaire. The denominator population for this study is all men in Natsal 2000 (4672), and all men aged 16–44 years from Natsal 1990 (6000).

All analyses were performed using Stata version 7.0 to account for stratification, clustering, and weighting of the sample.¹⁶ The data were weighted to correct for unequal selection probabilities and to match the age/sex population profile as previously described.^{11–13}

We used logistic regression to examine the association between paying for sex and a number of sociodemographic, sexual, and health behaviour factors. The crude and adjusted odds ratios (OR) are given with the corresponding 95% confidence interval (CI). The adjusted OR considers the association after controlling for age, marital status, area of residence, and partner numbers. Statistical significance is considered at $p < 0.05$ for all analyses. We obtained ethical approval from University College Hospital, North Thames Multicentre, and all local research ethics committees in Britain. The funders of this study did not have any role in the design of this study, the collection, analysis or interpretation of data, or in the drafting of this paper.

RESULTS

In Natsal 2000, 1.3% of men reported paying for sex in the previous 12 months, 4.2% in the previous 5 years, and 8.8% in their lifetime (table 1). These proportions had all increased significantly between 1990 and 2000. There was also a small increase in the mean number of different women who had ever been paid for sex.

In Natsal 2000, men who reported paying women for sex in the previous 5 years were more likely to be aged 25–34, to be previously or never married, and to be resident in London (table 2). There was no association with ethnicity, social class, or education.

Paying women for sex in the previous 5 years was associated with increased numbers of sexual partners: 36.5% of men who had paid for sex in the past 5 years reported 10 or more sexual partners in the past 5 years, and 54% had new sexual partners while abroad in that time (table 3). There was no association with having male sexual partners. Of 70 954 heterosexual partnerships (ever) reported by 5567 men in Natsal 2000, 2.5% (1744) were with

commercial sex workers. Among men paying for sex, 19.3% of all reported heterosexual partnerships were with commercial sex workers.

Table 3 also shows that less than one in five of the men who had paid for sex in the past 5 years had attended a sexually transmitted disease (STD) clinic in that time. Only 15.1% had been screened for HIV, although 9.1% reported an STI in the past 5 years.

We did not obtain specific information on the most recent commercial sexual contact, and therefore have limited data on condom use. However, taking only those men who had paid for sex in the previous 12 months, 35% had never used a condom in that time.

Men who paid for sex in the past 5 years were more likely to meet new sexual partners while abroad and from abroad, as shown in table 3. Table 4 shows the regions of origin of new sex partners reported by men who had paid for sex, although not all of these would have been commercial partners. This table also shows the potential for geographically widespread sexual mixing, including regions with much higher rates of HIV and STI than the United Kingdom.

DISCUSSION

This population based study shows that a significant proportion of men in Britain pay women for sex, and that this proportion is increasing. This behaviour is more common in men who live in London, who are in their late twenties and early thirties, and who are currently single, with the highest proportion in men who have been previously married. We have shown that men who pay for sex have other risks for STI, including higher numbers of partners and being more likely to meet new partners abroad. It is possible that some of the change between 1990 and 2000 is the result of increased reporting, with a greater acceptability of commercial sexual contact, but we think this is unlikely to account for the whole increase.¹⁷

The rate of divorce has increased, as has the proportion of men who are never or previously married, and this may explain some of the increased “demand” for commercial sex.⁸ Further evidence for the growth in commercial sex can be found from studies of sex workers. While there are no precise estimates on the numbers of sex workers in the United Kingdom or elsewhere, all reports suggest an increasingly large and diverse sex industry, with more opportunities for the sale and purchase of sex via sex clubs, escort agencies, the internet, and sex tourism.¹⁸

Table 2 Variations of the prevalence of reporting paying for heterosexual sex in the past 5 years by sociodemographic characteristics

	Prevalence of reporting paying for heterosexual sex, past 5 years			Base (weighted, unweighted)
	Row % (95% CI)	Crude ORs* (95% CI)	Adjusted ORs† (95% CI)	
All men	4.2 (3.6 to 4.9)	–	–	5614, 4699
Age (years)	p=0.014	p=0.013	p=0.0054	
16–24	3.4 (2.4 to 4.8)	1.00	1.00	1496, 1215
25–34	5.4 (4.4 to 6.6)	1.63 (1.07 to 2.46)	2.23 (1.33 to 3.72)	2107, 1771
35–44	3.6 (2.8 to 4.6)	1.05 (0.68 to 1.64)	1.69 (0.95 to 3.00)	2011, 1713
Marital status	p=0.0001	p=0.0001	p=0.0002	
Married	2.8 (2.1 to 3.8)	1.00	1.00	2237, 1546
Cohabiting	3.6 (2.4 to 5.6)	1.30 (0.76 to 2.22)	1.28 (0.76 to 2.17)	922, 652
Previously married	8.3 (5.6 to 12.2)	3.11 (1.83 to 5.29)	2.97 (1.72 to 5.12)	245, 320
Never married	5.4 (4.5 to 6.6)	1.97 (1.36 to 2.87)	2.27 (1.41 to 3.67)	2203, 2174
Self reported ethnicity	p=0.278	p=0.504	p=0.790	
White	4.1 (3.5 to 4.8)	1.00	1.00	5122, 4193
Black Caribbean	6.8 (2.1 to 19.8)	1.69 (0.49 to 5.81)	1.32 (0.37 to 4.80)	84, 101
Black African	3.4 (1.3 to 8.7)	0.82 (0.30 to 2.25)	0.38 (0.12 to 1.14)	53, 75
Indian	3.7 (1.6 to 8.3)	0.89 (0.37 to 2.13)	0.89 (0.37 to 2.13)	97, 85
Pakistani	0.7 (0.1 to 4.7)	0.15 (0.02 to 1.15)	0.18 (0.03 to 1.33)	48, 33
Other	6.6 (4.0 to 11.6)	1.65 (0.88 to 3.12)	1.34 (0.70 to 2.55)	200, 202
Social class	p=0.525	p=0.554	p=0.835	
I/II	4.3 (3.4 to 5.5)	1.00	1.00	1863, 1605
III non-manual/III manual	4.2 (3.3 to 5.3)	0.97 (0.69 to 1.37)	0.98 (0.69 to 1.39)	2201, 1790
IV/V	5.2 (3.8 to 7.3)	1.22 (0.79 to 1.88)	1.13 (0.71 to 1.78)	979, 823
Area of residence	p<0.0001	p<0.0001	p<0.0001	
London	8.6 (6.9 to 10.6)	1.00	1.00	809, 1156
Urban/city centre, not London	3.7 (2.3 to 5.8)	0.41 (0.24 to 0.70)	0.44 (0.25 to 0.77)	847, 608
Suburban residential	3.0 (2.3 to 3.9)	0.33 (0.23 to 0.48)	0.38 (0.26 to 0.56)	2473, 1814
Country town/village	4.1 (3.1 to 5.6)	0.46 (0.31 to 0.68)	0.54 (0.35 to 0.81)	1486, 1121
Education	p=0.147	p=0.170	p=0.368	
Degree	5.2 (4.0 to 6.8)	1.00	1.00	1205, 1090
A level/equivalent	4.7 (3.7 to 6.1)	0.91 (0.62 to 1.34)	1.10 (0.74 to 1.64)	1733, 1430
GCSE/O levels/equivalent	3.4 (2.6 to 4.4)	0.64 (0.43 to 0.95)	0.80 (0.53 to 1.19)	1990, 1591
Foreign/other	3.5 (0.7 to 15.2)	0.65 (0.13 to 3.34)	0.47 (0.09 to 2.35)	36, 39
None	3.5 (2.1 to 5.6)	0.66 (0.37 to 1.18)	0.72 (0.38 to 1.36)	639, 538

Base: All men aged 16–44 years in Natsal 2000.

*Crude odds ratio of reporting paying for heterosexual sex in the past 5 years.

†Odds ratio of reporting paying for heterosexual sex in the past 5 years adjusted for age, marital status, area of residence, and partner numbers in the past 5 years.

There are few other studies of the prevalence of commercial sexual contact by men, particularly from developed countries. A recent telephone population survey in Australia found that one in six men (15.6%) had ever paid for sex, with 1.9% in the

past year, an estimate similar to our own. The majority (97%) of these contacts were with women, 3% with men.¹⁹ Mikl²⁰ reported that 6% of men attending an STD clinic in Prague were clients of sex workers, but no time frame was given.

Table 3 Variations in the prevalence of reported sexual behaviours and sexual health indicators in the past 5 years by whether or not men reported paying for heterosexual sex in the past 5 years

In past 5 years	Reported paying for heterosexual sex in past 5 years		Odds ratio* (95% CI)	p Value
	Yes % (95% CI)	No % (95% CI)		
Sexual behaviour				
Partner numbers				<0.0001
0	0	7.7 (6.9 to 8.6)	1.00	
1	4.4 (2.4 to 8.0)	44.2 (42.6 to 45.7)	1.13 (0.24 to 5.24)	
2–4	26.9 (20.8 to 34.1)	27.6 (26.1 to 29.1)	11.17 (2.66 to 46.9)	
5–9	31.5 (25.1 to 38.8)	12.5 (11.4 to 13.6)	28.9 (6.91 to 121.1)	
10+	36.5 (29.7 to 44.0)	8.1 (7.3 to 9.0)	51.6 (12.4 to 215.9)	
Male sexual partner(s)	2.9 (1.5 to 5.8)	2.8 (2.3 to 3.3)	1.07 (0.51 to 2.23)	0.867
New sexual partner(s) while abroad	54.0 (46.5 to 61.2)	11.8 (10.8 to 12.9)	8.73 (6.38 to 12.0)	<0.0001
New sexual partner(s) while abroad from outside UK	50.7 (43.3 to 58.0)	7.1 (6.3 to 8.0)	13.4 (9.84 to 18.4)	<0.0001
Sexual health indicator				
Attended an STD clinic	18.9 (13.6 to 25.6)	7.1 (6.3 to 8.0)	3.06 (2.04 to 4.58)	<0.0001
STI diagnosis/es	9.1 (5.2 to 15.3)	2.7 (2.2 to 3.3)	3.61 (1.92 to 6.78)	<0.0001
HIV test	15.1 (10.6 to 21.0)	8.8 (7.9 to 9.8)	1.84 (1.21 to 2.81)	0.005
Base (weighted, unweighted)	236, 247	5378, 4452	–	–

Base: All men aged 16–44 years in Natsal 2000.

*Crude odds ratio of reporting paying for heterosexual sex in the past 5 years if reporting sexual behaviour or sexual health indicator relative to not reporting sexual behaviour or sexual health indicator except for partner numbers.

Table 4 Country/region of origin of new sexual partners while abroad (commercial and non-commercial) in the past 5 years

Country/region	% (95% CI) reporting new sexual partner(s) from country/region*†‡
United Kingdom	13.1 (9.0 to 18.7)
Other European countries	32.8 (26.2 to 40.3)
Australia/New Zealand	5.9 (3.3 to 10.3)
North America	7.4 (4.1 to 13.0)
South America/Central America	3.2 (1.5 to 6.8)
Caribbean countries	3.9 (1.6 to 9.1)
Asian countries	12.2 (8.2 to 17.7)
Middle East/North Africa	3.4 (1.1 to 9.5)
African countries (other than North Africa)	4.9 (2.3 to 10.0)
Other countries	2.4 (1.1 to 5.6)
Base (weighted, unweighted)	236, 269

Base: Men who reported paying for heterosexual sex in the past 5 years in Natsal 2000.

*Percentages do not sum to 100% since men could report new sexual Natsal 2000 partner(s) from 1+ countries/regions

†“Sexual partner” defined as “people who have had sex together—whether just once, or a few times, or as regular partners, or as married partners.”

‡Respondents were asked “Where did this/these new partner(s) normally live?” The CASI emphasised that “we are interested in which country the person comes from, not the country where you met”. Response options as listed in table 4.

There are more data from developing countries where health surveys include a relevant indicator—namely, the percentage of male respondents reporting sex with a sex worker in the past 12 months. Estimates range from less than 2% in Nepal (2001), Kazakhstan (1999), Uganda (2000/1), Rwanda (2000), and Namibia (2000), to 10% or more in the Central African Republic (1994–5), Dominican Republic (1996), Mozambique (2003), Benin (1996), Cameroon

(1998), and Zambia (1996). With the exception of Cameroon (20%) and Mozambique (13%), those with the higher figures used a broader definition (the number of male respondents reporting that they had given or received money, gifts, or favours in return for sex in the last 12 months).²¹

It is of concern that only a minority of the men who paid for sex had been to an STD clinic or been screened for HIV, suggesting that they do not see themselves as being at risk. These men may be assuming that the women will take responsibility for safety and screening, which may be valid in some areas since sex workers have been shown to be very good consumers of safer sex advice.

We have demonstrated an increase in the proportion of men paying for female sex contacts. We have also demonstrated that men who pay for sex have a higher incidence of reported STI than men who do not. It is not clear however whether this increased risk is related to commercial sexual contacts per se or related to the much higher rate of sex partner change (commercial and non-commercial) in this group. For men who paid for sex, less than one in five (19%) of all partners were commercial, and we know from other studies that condom use is relatively high in commercial sex, probably higher than in non-commercial casual partnerships.

It is clear that some of these men could be linking otherwise separate sexual networks. They are meeting new partners abroad, some of whom are likely to be sex workers. The distribution of regions where new partners were met is consistent with sex tourism: western Europe (where the Netherlands and Germany have large, visible sex industries) and Asia (where, for example, Thailand and India have large numbers of sex workers). Cheap air travel has led to an increase in the number of people travelling to these destinations, and many will have sex with sex workers and with other people they meet.²²

In conclusion, men who pay for sex are a substantial group who are at increased risk of acquiring and potentially transmitting STI and HIV. However, their increased risk of STI is likely to be linked as much to their high rates of sexual partner change with non-commercial as with commercial contacts. To understand the risks better we should be improving surveillance, through routinely asking men attending clinics whether they have paid for sex, and monitoring how many STIs may have been acquired or transmitted during commercial sex.

Men who pay for sex should be the target of health promotion campaigns and screening initiatives as a result of their increased risk and high rate of partner change. However, as this study shows, it is very difficult to identify clients as a distinct group—the behaviour occurs across ages, social classes and ethnic groups, and is increasing. Health promotion therefore needs to address men in general, starting with school based sex education and mass media campaigns. Groups that warrant special attention are single and previously married men and tourists. While men generally do not discuss commercial sexual contacts with each other, there are situations where this does occur including holidays and “stag party” trips where commercial sex may be a collective experience. Targeting these groups could lead to peer interventions to improve safety. Any campaigns need to take into account the stigma of sex work; women who sell sex are vulnerable to violence and other abuse, and any campaign targeting them as the “source” of infections would be both misleading, since they have low levels of infection in the United Kingdom, and could rebound through increased violence. Greater understanding of the changing nature of commercial sex, and the engagement of men who pay for sex in both descriptive research and the design of interventions, is an essential next step.

What this paper adds

- A significant and growing proportion of men in the United Kingdom have paid women for sex
- Paying women for sex is most likely to be reported by single men, men living in London, and men aged between 25 years and 34 years, but there was no association with ethnicity or social class
- Men who pay for sex also have higher rates of sexual partner change with non-commercial contacts, and report higher rates of STI than other men. Few have been tested for HIV

Policy implications

- We need health promotion campaigns to outline the need for safer sex in commercial contacts, and the need for regular screening for clients as well as sex workers. These will have to be aimed at a broad audience given the wide range of men who pay for sex. Policies towards prostitution should be based on an understanding of the widespread nature of commercial sexual contact, and focus on harm minimisation

ACKNOWLEDGEMENTS

We thank the study participants, the team of interviewers and operations, and computing staff from the National Centre for Social Research who carried out the interviews. The study was supported by a grant from the Medical Research Council with funds from the Department of Health, the Scottish Executive, and the National Assembly for Wales. The late Jane Wadsworth made important contributions to survey design.

CONTRIBUTORS

HW participated in developing the concept, analysing and interpreting the data, and drafted the paper; AJ participated in developing the concept, analysing and interpreting the data, and preparation of the paper; CM did the statistical analysis and participated in the preparation of the paper; AJ, KW, KF, BE, and AC participated in the design and management of the main study and preparation of this paper.

Authors' affiliations

H Ward, Department of Infectious Disease Epidemiology, Imperial College London, UK

C H Mercer, K Fenton, A Copas, A M Johnson, Department of Primary Care and Population Sciences, University College London, UK

K Wellings, Centre for Reproductive and Sexual Health, London School of Hygiene and Tropical Medicine, UK

B Erens, National Centre for Social Research, UK

Conflicts of interest: none.

REFERENCES

- Garnett GP**. The geographical and temporal evolution of sexually transmitted disease epidemics. *Sex Transm Infect* 2002;**78**:14i-19.
- Aral SO, Blanchard JF**. Phase specific approaches to the epidemiology and prevention of sexually transmitted diseases. *Sex Transm Infect* 2002;**78**:1i-2.
- Vanwesenbeeck I**. Another decade of social scientific work on sex work: a review of research 1990-2000. *Annu Rev Sex Res* 2001;**12**:242-89.
- Morris M, Podhisita C, Wawer M, et al**. Bridge populations in the spread of HIV/AIDS in Thailand. *AIDS* 1996;**10**:1265-71.
- Ramjee G, Gouws aE**. Prevalence of HIV among truck drivers visiting sex workers in KwaZulu-Natal, South Africa. *Sex Transm Dis* 2002;**29**:44-9.
- Lau JT, Thomas J**. Risk behaviours of Hong Kong male residents travelling to mainland China: a potential bridge population for HIV infection. *AIDS Care* 2001;**13**:71-81.
- Ghys PD, Saitel T, Vu HT, et al**. Growing in silence: selected regions and countries with expanding HIV/AIDS epidemics. *AIDS* 2003;**17**(Suppl 4):S45-50.
- Creese A, Floyd K, Alban A, et al**. Cost-effectiveness of HIV/AIDS interventions in Africa: a systematic review of the evidence. *Lancet* 2002;**359**:1635-42.
- Gorbach PM, Sopheab H, Phalla T, et al**. Sexual bridging by Cambodian men: potential importance for general population spread of STD and HIV epidemics. *Sex Transm Dis* 2000;**27**:320-6.
- Brown AE, Sadler KE, Tomkins SE, et al**. Recent trends in HIV and other STIs in the United Kingdom: data to the end of 2002. *Sex Transm Infect* 2004;**80**:159-66.
- Johnson AM, Mercer CH, Erens B, et al**. Sexual behaviour in Britain: partnerships, practices, and HIV risk behaviours. *Lancet* 2001;**358**:1835-42.
- Ward H, Day S, Green A, et al**. Declining prevalence of STI in the London sex industry, 1985 to 2002. *Sex Transm Infect* 2004;**80**:374-9.
- Erens B, McManus S, Field J, et al**. *National survey of sexual attitudes and lifestyles. II: Technical report*. London: National Centre for Social Research, 2001.
- Johnson AM, Wadsworth J, Wellings K, et al**. Sexual lifestyles and HIV risk. *Nature* 1992;**360**:410-12.
- Johnson AM, Wadsworth J, Wellings K, et al**. *Sexual attitudes and lifestyles*. Oxford: Blackwell Scientific Press, 1994.
- Stata Corp**. *Stata statistical software: Release 7.0*. Texas: Stata Corporation, 2001.
- Copas AJ, Wellings K, Erens B, et al**. The accuracy of reported sensitive sexual behaviour in Britain: exploring the extent of change 1990-2000. *Sex Transm Infect* 2002;**78**:26-30.
- Ward H, Day S**. Sex work in context. In: Day S, Ward H, eds. *Sex work, mobility and health in Europe*. London: Kegan Paul, 2004.
- Rissel CE, Richters J, Grulich AE, et al**. Sex in Australia: experiences of commercial sex in a representative sample of adults. *Aust N Z J Public Health* 2003;**27**:191-7.
- Mikl J, Sudar Z, Smith PF, et al**. HIV infection and high risk behavior among patients attending an STD referral clinic in Prague, Czech Republic. *Sex Transm Infect* 1998;**74**:128-30.
- HIV/AIDS Survey Indicators Database (www.measuredhs.com/hivdata/about_db.cfm) accessed August, 2004.
- Bellis MA, Hughes K, Thomson R, et al**. Sexual behaviour of young people in international tourist resorts. *Sex Transm Infect* 2004;**80**:43-7.

ECHO

Late diagnosis still delays HIV treatment for many



Please visit the Sexually Transmitted Infections website [www.stijournal.com] for a link to the full text of this article.

Missed opportunities for diagnosing HIV are denying patients the benefit of early treatment, say researchers. More patients at risk should be encouraged to have an HIV test, and health professionals must get better at recognising symptoms of the infection.

This follows a national case review of new diagnoses of HIV infection between January and March 2003 by a survey to all adult HIV care providers in the United Kingdom and Ireland. The survey covered clinical and immune state and symptoms or episodes in the previous 12 months.

A third of the 977 patients identified had presented late, as indicated by a CD4 lymphocyte count <200 cells/ μ l. They were commonly older patients and black Africans. Diagnosis as part of a routine screen and testing at genitourinary, sexual health, and HIV clinics were both independently associated with a lower chance of late diagnosis, after adjustment for demographic confounders. Seventeen per cent (168) of all patients had had an episode in the previous year suggesting HIV infection, including 58 hospital admissions. In 160 of these patients CD4 counts were below the threshold for starting treatment advocated by the British HIV Association, possibly indicating delayed treatment. The response rate to the survey was 76% (113/148).

The results reflect a national trend for late diagnosis of HIV infection reported by the Health Protection Agency. Estimates in 2001 disclosed that 59% of patients with HIV in the United Kingdom starting treatment had low CD4 counts (<200 cells/ μ l), mainly owing to late diagnosis.

▲ Sullivan AK, et al. *BMJ* 2005;**330**:1301-1302.