

## Search for Maximal Flavor Violating Scalars in Same-Charge Lepton Pairs in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV

T. Aaltonen,<sup>24</sup> J. Adelman,<sup>14</sup> T. Akimoto,<sup>55</sup> M. G. Albrow,<sup>18</sup> B. Álvarez González,<sup>12</sup> S. Amerio,<sup>43</sup> D. Amidei,<sup>35</sup> A. Anastassov,<sup>52</sup> A. Annovi,<sup>20</sup> J. Antos,<sup>15</sup> M. Aoki,<sup>25</sup> G. Apollinari,<sup>18</sup> A. Apresyan,<sup>48</sup> T. Arisawa,<sup>57</sup> A. Artikov,<sup>16</sup> W. Ashmanskas,<sup>18</sup> A. Attal,<sup>4</sup> A. Aurisano,<sup>53</sup> F. Azfar,<sup>42</sup> P. Azzi-Bacchetta,<sup>43</sup> P. Azzurri,<sup>46</sup> N. Bacchetta,<sup>43</sup> W. Badgett,<sup>18</sup> A. Barbaro-Galtieri,<sup>29</sup> V. E. Barnes,<sup>48</sup> B. A. Barnett,<sup>26</sup> S. Baroiant,<sup>8</sup> S. Bar-Shalom,<sup>61</sup> V. Bartsch,<sup>31</sup> G. Bauer,<sup>33</sup> P.-H. Beauchemin,<sup>34</sup> F. Bedeschi,<sup>46</sup> P. Bednar,<sup>15</sup> S. Behari,<sup>26</sup> G. Belletini,<sup>46</sup> J. Bellinger,<sup>59</sup> A. Belloni,<sup>23</sup> D. Benjamin,<sup>17</sup> A. Beretvas,<sup>18</sup> J. Beringer,<sup>29</sup> T. Berry,<sup>30</sup> A. Bhatti,<sup>50</sup> M. Binkley,<sup>18</sup> D. Bisello,<sup>43</sup> I. Bizjak,<sup>31</sup> R. E. Blair,<sup>3</sup> C. Blocker,<sup>7</sup> B. Blumenfeld,<sup>26</sup> A. Bocci,<sup>17</sup> A. Bodek,<sup>49</sup> V. Boisvert,<sup>49</sup> G. Bolla,<sup>48</sup> A. Bolshov,<sup>33</sup> D. Bortoletto,<sup>48</sup> J. Boudreau,<sup>47</sup> A. Boveia,<sup>11</sup> B. Brau,<sup>11</sup> A. Bridgeman,<sup>25</sup> L. Brigliadori,<sup>6</sup> C. Bromberg,<sup>36</sup> E. Brubaker,<sup>14</sup> J. Budagov,<sup>16</sup> H. S. Budd,<sup>49</sup> S. Budd,<sup>25</sup> K. Burkett,<sup>18</sup> G. Busetto,<sup>43</sup> P. Bussey,<sup>22</sup> A. Buzatu,<sup>34</sup> K. L. Byrum,<sup>3</sup> S. Cabrera,<sup>17,r</sup> M. Campanelli,<sup>36</sup> M. Campbell,<sup>35</sup> F. Canelli,<sup>18</sup> A. Canepa,<sup>45</sup> D. Carlsmith,<sup>59</sup> R. Carosi,<sup>46</sup> S. Carrillo,<sup>19,l</sup> S. Carron,<sup>34</sup> B. Casal,<sup>12</sup> M. Casarsa,<sup>18</sup> A. Castro,<sup>6</sup> P. Catastini,<sup>46</sup> D. Cauz,<sup>54</sup> M. Cavalli-Sforza,<sup>4</sup> A. Cerri,<sup>29</sup> L. Cerrito,<sup>31,p</sup> S. H. Chang,<sup>28</sup> Y. C. Chen,<sup>1</sup> M. Chertok,<sup>8</sup> G. Chiarelli,<sup>46</sup> G. Chlachidze,<sup>18</sup> F. Chlebana,<sup>18</sup> K. Cho,<sup>28</sup> D. Chokheli,<sup>16</sup> J. P. Chou,<sup>23</sup> G. Choudalakis,<sup>33</sup> S. H. Chuang,<sup>52</sup> K. Chung,<sup>13</sup> W. H. Chung,<sup>59</sup> Y. S. Chung,<sup>49</sup> C. I. Ciobanu,<sup>25</sup> M. A. Ciocci,<sup>46</sup> A. Clark,<sup>21</sup> D. Clark,<sup>7</sup> G. Compostella,<sup>43</sup> M. E. Convery,<sup>18</sup> J. Conway,<sup>8</sup> B. Cooper,<sup>31</sup> K. Copic,<sup>35</sup> M. Cordelli,<sup>20</sup> G. Cortiana,<sup>43</sup> F. Crescioli,<sup>46</sup> C. Cuenca Almenar,<sup>8,r</sup> J. Cuevas,<sup>12,o</sup> R. Culbertson,<sup>18</sup> J. C. Cully,<sup>35</sup> D. Dagenhart,<sup>18</sup> M. Datta,<sup>18</sup> T. Davies,<sup>22</sup> P. de Barbaro,<sup>49</sup> S. De Cecco,<sup>51</sup> A. Deisher,<sup>29</sup> G. De Lentdecker,<sup>49,d</sup> G. De Lorenzo,<sup>4</sup> M. Dell'Orso,<sup>46</sup> L. Demortier,<sup>50</sup> J. Deng,<sup>17</sup> M. Deninno,<sup>6</sup> D. De Pedis,<sup>51</sup> P. F. Derwent,<sup>18</sup> G. P. Di Giovanni,<sup>44</sup> C. Dionisi,<sup>51</sup> B. Di Ruzza,<sup>54</sup> J. R. Dittmann,<sup>5</sup> M. D'Onofrio,<sup>4</sup> S. Donati,<sup>46</sup> P. Dong,<sup>9</sup> J. Donini,<sup>43</sup> T. Dorigo,<sup>43</sup> S. Dube,<sup>52</sup> J. Efron,<sup>39</sup> R. Erbacher,<sup>8</sup> D. Errede,<sup>25</sup> S. Errede,<sup>25</sup> R. Eusebi,<sup>18</sup> H. C. Fang,<sup>29</sup> S. Farrington,<sup>30</sup> W. T. Fedorko,<sup>14</sup> R. G. Feild,<sup>60</sup> M. Feindt,<sup>27</sup> J. P. Fernandez,<sup>32</sup> C. Ferrazza,<sup>46</sup> R. Field,<sup>19</sup> G. Flanagan,<sup>48</sup> R. Forrest,<sup>8</sup> S. Forrester,<sup>8</sup> M. Franklin,<sup>23</sup> J. C. Freeman,<sup>29</sup> I. Furic,<sup>19</sup> M. Gallinaro,<sup>50</sup> J. Galyardt,<sup>13</sup> F. Garbersson,<sup>11</sup> J. E. Garcia,<sup>46</sup> A. F. Garfinkel,<sup>48</sup> K. Genser,<sup>18</sup> H. Gerberich,<sup>25</sup> D. Gerdes,<sup>35</sup> S. Giagu,<sup>51</sup> V. Giakoumopolou,<sup>46,a</sup> P. Giannetti,<sup>46</sup> K. Gibson,<sup>47</sup> J. L. Gimmell,<sup>49</sup> C. M. Ginsburg,<sup>18</sup> N. Giokaris,<sup>16,a</sup> M. Giordani,<sup>54</sup> P. Giromini,<sup>20</sup> M. Giunta,<sup>46</sup> V. Glagolev,<sup>16</sup> D. Glenzinski,<sup>18</sup> M. Gold,<sup>37</sup> N. Goldschmidt,<sup>19</sup> A. Golossanov,<sup>18</sup> G. Gomez,<sup>12</sup> G. Gomez-Ceballos,<sup>33</sup> M. Goncharov,<sup>53</sup> O. González,<sup>32</sup> I. Gorelov,<sup>37</sup> A. T. Goshaw,<sup>17</sup> K. Goulianos,<sup>50</sup> A. Gresele,<sup>43</sup> S. Grinstein,<sup>23</sup> C. Grosso-Pilcher,<sup>14</sup> R. C. Group,<sup>18</sup> U. Grundler,<sup>25</sup> J. Guimaraes da Costa,<sup>23</sup> Z. Gunay-Unalan,<sup>36</sup> C. Haber,<sup>29</sup> K. Hahn,<sup>33</sup> S. R. Hahn,<sup>18</sup> E. Halkiadakis,<sup>52</sup> A. Hamilton,<sup>21</sup> B.-Y. Han,<sup>49</sup> J. Y. Han,<sup>49</sup> R. Handler,<sup>59</sup> F. Happacher,<sup>20</sup> K. Hara,<sup>55</sup> D. Hare,<sup>52</sup> M. Hare,<sup>56</sup> S. Harper,<sup>42</sup> R. F. Harr,<sup>58</sup> R. M. Harris,<sup>18</sup> M. Hartz,<sup>47</sup> K. Hatakeyama,<sup>50</sup> J. Hauser,<sup>9</sup> C. Hays,<sup>42</sup> M. Heck,<sup>27</sup> A. Heijboer,<sup>45</sup> B. Heinemann,<sup>29</sup> J. Heinrich,<sup>45</sup> C. Henderson,<sup>33</sup> M. Herndon,<sup>59</sup> J. Heuser,<sup>27</sup> S. Hewamanage,<sup>5</sup> D. Hidas,<sup>17</sup> C. S. Hill,<sup>11,c</sup> D. Hirschbuehl,<sup>27</sup> A. Hocker,<sup>18</sup> S. Hou,<sup>1</sup> M. Houlden,<sup>30</sup> S.-C. Hsu,<sup>10</sup> B. T. Huffman,<sup>42</sup> R. E. Hughes,<sup>39</sup> U. Husemann,<sup>60</sup> J. Huston,<sup>36</sup> J. Incandela,<sup>11</sup> G. Introzzi,<sup>46</sup> M. Iori,<sup>51</sup> A. Ivanov,<sup>8</sup> B. Iyutin,<sup>33</sup> E. James,<sup>18</sup> B. Jayatilaka,<sup>17</sup> D. Jeans,<sup>51</sup> E. J. Jeon,<sup>28</sup> S. Jindariani,<sup>19</sup> W. Johnson,<sup>8</sup> M. Jones,<sup>48</sup> K. K. Joo,<sup>28</sup> S. Y. Jun,<sup>13</sup> J. E. Jung,<sup>28</sup> T. R. Junk,<sup>25</sup> T. Kamon,<sup>53</sup> D. Kar,<sup>19</sup> P. E. Karchin,<sup>58</sup> Y. Kato,<sup>41</sup> R. Kephart,<sup>18</sup> U. Kerzel,<sup>27</sup> V. Khotilovich,<sup>53</sup> B. Kilminster,<sup>39</sup> D. H. Kim,<sup>28</sup> H. S. Kim,<sup>28</sup> J. E. Kim,<sup>28</sup> M. J. Kim,<sup>18</sup> S. B. Kim,<sup>28</sup> S. H. Kim,<sup>55</sup> Y. K. Kim,<sup>14</sup> N. Kimura,<sup>55</sup> L. Kirsch,<sup>7</sup> S. Klimentenko,<sup>19</sup> M. Klute,<sup>33</sup> B. Knuteson,<sup>33</sup> B. R. Ko,<sup>17</sup> S. A. Koay,<sup>11</sup> K. Kondo,<sup>57</sup> D. J. Kong,<sup>28</sup> J. Konigsberg,<sup>19</sup> A. Korytov,<sup>19</sup> A. V. Kotwal,<sup>17</sup> J. Kraus,<sup>25</sup> M. Kreps,<sup>27</sup> J. Kroll,<sup>45</sup> N. Krumnack,<sup>5</sup> M. Kruse,<sup>17</sup> V. Krutelyov,<sup>11</sup> T. Kubo,<sup>55</sup> S. E. Kuhlmann,<sup>3</sup> T. Kuhr,<sup>27</sup> N. P. Kulkarni,<sup>58</sup> Y. Kusakabe,<sup>57</sup> S. Kwang,<sup>14</sup> A. T. Laasanen,<sup>48</sup> S. Lai,<sup>34</sup> S. Lami,<sup>46</sup> S. Lammel,<sup>18</sup> M. Lancaster,<sup>31</sup> R. L. Lander,<sup>8</sup> K. Lannon,<sup>39</sup> A. Lath,<sup>52</sup> G. Latino,<sup>46</sup> I. Lazzizzera,<sup>43</sup> T. LeCompte,<sup>3</sup> J. Lee,<sup>49</sup> J. Lee,<sup>28</sup> Y. J. Lee,<sup>28</sup> S. W. Lee,<sup>53,q</sup> R. Lefèvre,<sup>21</sup> N. Leonardo,<sup>33</sup> S. Leone,<sup>46</sup> S. Levy,<sup>14</sup> J. D. Lewis,<sup>18</sup> C. Lin,<sup>60</sup> C. S. Lin,<sup>29</sup> J. Linacre,<sup>42</sup> M. Lindgren,<sup>18</sup> E. Lipeles,<sup>10</sup> A. Lister,<sup>8</sup> D. O. Litvintsev,<sup>18</sup> T. Liu,<sup>18</sup> N. S. Lockyer,<sup>45</sup> A. Loginov,<sup>60</sup> M. Loretì,<sup>43</sup> L. Lovas,<sup>15</sup> R.-S. Lu,<sup>1</sup> D. Lucchesi,<sup>43</sup> J. Lueck,<sup>27</sup> C. Luci,<sup>51</sup> P. Lujan,<sup>29</sup> P. Lukens,<sup>18</sup> G. Lungu,<sup>19</sup> L. Lyons,<sup>42</sup> J. Lys,<sup>29</sup> R. Lysak,<sup>15</sup> E. Lytken,<sup>48</sup> P. Mack,<sup>27</sup> D. MacQueen,<sup>34</sup> R. Madrak,<sup>18</sup> K. Maeshima,<sup>18</sup> K. Makhoul,<sup>33</sup> T. Maki,<sup>24</sup> P. Maksimovic,<sup>26</sup> S. Malde,<sup>42</sup> S. Malik,<sup>31</sup> G. Manca,<sup>30</sup> A. Manousakis,<sup>16,a</sup> F. Margaroli,<sup>48</sup> C. Marino,<sup>27</sup> C. P. Marino,<sup>25</sup> A. Martin,<sup>60</sup> M. Martin,<sup>26</sup> V. Martin,<sup>22,j</sup> M. Martínez,<sup>4</sup> R. Martínez-Ballarín,<sup>32</sup> T. Maruyama,<sup>55</sup> P. Mastrandrea,<sup>51</sup> T. Masubuchi,<sup>55</sup> M. E. Mattson,<sup>58</sup> P. Mazzanti,<sup>6</sup> K. S. McFarland,<sup>49</sup> P. McIntyre,<sup>53</sup> R. McNulty,<sup>30,i</sup> A. Mehta,<sup>30</sup> P. Mehtala,<sup>24</sup> S. Menzemer,<sup>12,k</sup> A. Menzione,<sup>46</sup> P. Merkel,<sup>48</sup> C. Mesropian,<sup>50</sup> A. Messina,<sup>36</sup> T. Miao,<sup>18</sup> N. Miladinovic,<sup>7</sup> J. Miles,<sup>33</sup> R. Miller,<sup>36</sup> C. Mills,<sup>23</sup> M. Milnik,<sup>27</sup> A. Mitra,<sup>1</sup> G. Mitselmakher,<sup>19</sup> H. Miyake,<sup>55</sup> S. Moed,<sup>23</sup> N. Moggi,<sup>6</sup> C. S. Moon,<sup>28</sup> R. Moore,<sup>18</sup> M. Morello,<sup>46</sup> P. Movilla Fernandez,<sup>29</sup>

J. Mülmenstädt,<sup>29</sup> A. Mukherjee,<sup>18</sup> Th. Muller,<sup>27</sup> R. Mumford,<sup>26</sup> P. Murat,<sup>18</sup> M. Mussini,<sup>6</sup> J. Nachtman,<sup>18</sup> Y. Nagai,<sup>55</sup> A. Nagano,<sup>55</sup> J. Naganoma,<sup>57</sup> K. Nakamura,<sup>55</sup> I. Nakano,<sup>40</sup> A. Napier,<sup>56</sup> V. Necula,<sup>17</sup> C. Neu,<sup>45</sup> M. S. Neubauer,<sup>25</sup> J. Nielsen,<sup>29,f</sup> L. Nodulman,<sup>3</sup> M. Norman,<sup>10</sup> O. Norriella,<sup>25</sup> E. Nurse,<sup>31</sup> S. H. Oh,<sup>17</sup> Y. D. Oh,<sup>28</sup> I. Oksuzian,<sup>19</sup> T. Okusawa,<sup>41</sup> R. Oldeman,<sup>30</sup> R. Orava,<sup>24</sup> K. Osterberg,<sup>24</sup> S. Pagan Griso,<sup>43</sup> C. Pagliarone,<sup>46</sup> E. Palencia,<sup>18</sup> V. Papadimitriou,<sup>18</sup> A. Papaikonomou,<sup>27</sup> A. A. Paramonov,<sup>14</sup> B. Parks,<sup>39</sup> S. Pashapour,<sup>34</sup> J. Patrick,<sup>18</sup> G. Pauletta,<sup>54</sup> M. Paulini,<sup>13</sup> C. Paus,<sup>33</sup> D. E. Pellett,<sup>8</sup> A. Penzo,<sup>54</sup> T. J. Phillips,<sup>17</sup> G. Piacentino,<sup>46</sup> J. Piedra,<sup>44</sup> L. Pinera,<sup>19</sup> K. Pitts,<sup>25</sup> C. Plager,<sup>9</sup> L. Pondrom,<sup>59</sup> X. Portell,<sup>4</sup> O. Poukhov,<sup>16</sup> N. Pounder,<sup>42</sup> F. Prakoshyn,<sup>16</sup> A. Pronko,<sup>18</sup> J. Proudfoot,<sup>3</sup> F. Ptohos,<sup>18,h</sup> G. Punzi,<sup>46</sup> J. Pursley,<sup>59</sup> J. Rademacker,<sup>42,c</sup> A. Rahaman,<sup>47</sup> A. Rajaraman,<sup>61</sup> V. Ramakrishnan,<sup>59</sup> N. Ranjan,<sup>48</sup> I. Redondo,<sup>32</sup> B. Reisert,<sup>18</sup> V. Rekovic,<sup>37</sup> P. Renton,<sup>42</sup> M. Rescigno,<sup>51</sup> S. Richter,<sup>27</sup> F. Rimondi,<sup>6</sup> L. Ristori,<sup>46</sup> A. Robson,<sup>22</sup> T. Rodrigo,<sup>12</sup> E. Rogers,<sup>25</sup> S. Rolli,<sup>56</sup> R. Roser,<sup>18</sup> M. Rossi,<sup>54</sup> R. Rossin,<sup>11</sup> P. Roy,<sup>34</sup> A. Ruiz,<sup>12</sup> J. Russ,<sup>13</sup> V. Rusu,<sup>18</sup> H. Saarikko,<sup>24</sup> A. Safonov,<sup>53</sup> W. K. Sakumoto,<sup>49</sup> G. Salamanna,<sup>51</sup> O. Saltó,<sup>4</sup> L. Santi,<sup>54</sup> S. Sarkar,<sup>51</sup> L. Sartori,<sup>46</sup> K. Sato,<sup>18</sup> A. Savoy-Navarro,<sup>44</sup> T. Scheidle,<sup>27</sup> P. Schlabach,<sup>18</sup> E. E. Schmidt,<sup>18</sup> M. A. Schmidt,<sup>14</sup> M. P. Schmidt,<sup>60</sup> M. Schmitt,<sup>38</sup> T. Schwarz,<sup>8</sup> L. Scodellaro,<sup>12</sup> A. L. Scott,<sup>11</sup> A. Scribano,<sup>46</sup> F. Scuri,<sup>46</sup> A. Sedov,<sup>48</sup> S. Seidel,<sup>37</sup> Y. Seiya,<sup>41</sup> A. Semenov,<sup>16</sup> L. Sexton-Kennedy,<sup>18</sup> A. Sfyria,<sup>21</sup> S. Z. Shalhout,<sup>58</sup> M. D. Shapiro,<sup>29</sup> T. Shears,<sup>30</sup> P. F. Shepard,<sup>47</sup> D. Sherman,<sup>23</sup> M. Shimojima,<sup>55,n</sup> M. Shochet,<sup>14</sup> Y. Shon,<sup>59</sup> I. Shreyber,<sup>21</sup> A. Sidoti,<sup>46</sup> P. Sinervo,<sup>34</sup> A. Sisakyan,<sup>16</sup> A. J. Slaughter,<sup>18</sup> J. Slaunwhite,<sup>39</sup> K. Sliwa,<sup>56</sup> J. R. Smith,<sup>8</sup> F. D. Snider,<sup>18</sup> R. Snihur,<sup>34</sup> M. Soderberg,<sup>35</sup> A. Soha,<sup>8</sup> S. Somalwar,<sup>52</sup> V. Sorin,<sup>36</sup> J. Spalding,<sup>18</sup> F. Spinella,<sup>46</sup> T. Spreitzer,<sup>34</sup> P. Squillacioti,<sup>46</sup> M. Stanitzki,<sup>60</sup> R. St. Denis,<sup>22</sup> B. Stelzer,<sup>9</sup> O. Stelzer-Chilton,<sup>42</sup> D. Stentz,<sup>38</sup> J. Strologas,<sup>37</sup> D. Stuart,<sup>11</sup> J. S. Suh,<sup>28</sup> A. Sukhanov,<sup>19</sup> H. Sun,<sup>56</sup> I. Suslov,<sup>16</sup> T. Suzuki,<sup>55</sup> A. Taffard,<sup>25,e</sup> R. Takashima,<sup>40</sup> Y. Takeuchi,<sup>55</sup> R. Tanaka,<sup>40</sup> M. Tecchio,<sup>35</sup> P. K. Teng,<sup>1</sup> K. Terashi,<sup>50</sup> J. Thom,<sup>18,g</sup> A. S. Thompson,<sup>22</sup> G. A. Thompson,<sup>25</sup> E. Thomson,<sup>45</sup> P. Tipton,<sup>60</sup> V. Tiwari,<sup>13</sup> S. Tkaczyk,<sup>18</sup> D. Toback,<sup>53</sup> S. Tokar,<sup>15</sup> K. Tollefson,<sup>36</sup> T. Tomura,<sup>55</sup> D. Tonelli,<sup>18</sup> S. Torre,<sup>20</sup> D. Torretta,<sup>18</sup> S. Tourneur,<sup>44</sup> W. Trischuk,<sup>34</sup> Y. Tu,<sup>45</sup> N. Turini,<sup>46</sup> F. Ukegawa,<sup>55</sup> S. Uozumi,<sup>55</sup> S. Vallecorsa,<sup>21</sup> N. van Remortel,<sup>24</sup> A. Varganov,<sup>35</sup> E. Vataga,<sup>37</sup> F. Vázquez,<sup>19,1</sup> G. Velev,<sup>18</sup> C. Vellidis,<sup>46,a</sup> V. Veszpremi,<sup>48</sup> M. Vidal,<sup>32</sup> R. Vidal,<sup>18</sup> I. Vila,<sup>12</sup> R. Vilar,<sup>12</sup> T. Vine,<sup>31</sup> M. Vogel,<sup>37</sup> I. Volobouev,<sup>29,q</sup> G. Volpi,<sup>46</sup> F. Würthwein,<sup>10</sup> P. Wagner,<sup>45</sup> R. G. Wagner,<sup>3</sup> R. L. Wagner,<sup>18</sup> J. Wagner-Kuhr,<sup>27</sup> W. Wagner,<sup>27</sup> T. Wakisaka,<sup>41</sup> R. Wallny,<sup>9</sup> S. M. Wang,<sup>1</sup> A. Warburton,<sup>34</sup> D. Waters,<sup>31</sup> M. Weinberger,<sup>53</sup> W. C. Wester III,<sup>18</sup> B. Whitehouse,<sup>56</sup> D. Whiteson,<sup>45,e</sup> A. B. Wicklund,<sup>3</sup> E. Wicklund,<sup>18</sup> G. Williams,<sup>34</sup> H. H. Williams,<sup>45</sup> P. Wilson,<sup>18</sup> B. L. Winer,<sup>39</sup> P. Wittich,<sup>18,g</sup> S. Wolbers,<sup>18</sup> C. Wolfe,<sup>14</sup> T. Wright,<sup>35</sup> X. Wu,<sup>21</sup> S. M. Wynne,<sup>30</sup> A. Yagil,<sup>10</sup> K. Yamamoto,<sup>41</sup> J. Yamaoka,<sup>52</sup> T. Yamashita,<sup>40</sup> C. Yang,<sup>60</sup> U. K. Yang,<sup>14,m</sup> Y. C. Yang,<sup>28</sup> W. M. Yao,<sup>29</sup> G. P. Yeh,<sup>18</sup> J. Yoh,<sup>18</sup> K. Yorita,<sup>14</sup> T. Yoshida,<sup>41</sup> G. B. Yu,<sup>49</sup> F. Yu,<sup>61</sup> I. Yu,<sup>28</sup> S. S. Yu,<sup>18</sup> J. C. Yun,<sup>18</sup> L. Zanello,<sup>51</sup> A. Zanetti,<sup>54</sup> I. Zaw,<sup>23</sup> X. Zhang,<sup>25</sup> Y. Zheng,<sup>9,b</sup> and S. Zucchelli<sup>6</sup>

(CDF Collaboration)<sup>s</sup><sup>1</sup>*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*<sup>2</sup>*Informatics Institute, University of Amsterdam, Amsterdam, The Netherlands*<sup>3</sup>*Argonne National Laboratory, Argonne, Illinois 60439, USA*<sup>4</sup>*Institut de Física d'Altes Energies, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*<sup>5</sup>*Baylor University, Waco, Texas 76798, USA*<sup>6</sup>*Istituto Nazionale di Fisica Nucleare, University of Bologna, I-40127 Bologna, Italy*<sup>7</sup>*Brandeis University, Waltham, Massachusetts 02254, USA*<sup>8</sup>*University of California, Davis, Davis, California 95616, USA*<sup>9</sup>*University of California, Los Angeles, Los Angeles, California 90024, USA*<sup>10</sup>*University of California, San Diego, La Jolla, California 92093, USA*<sup>11</sup>*University of California, Santa Barbara, Santa Barbara, California 93106, USA*<sup>12</sup>*Instituto de Física de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*<sup>13</sup>*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*<sup>14</sup>*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*<sup>15</sup>*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*<sup>16</sup>*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*<sup>17</sup>*Duke University, Durham, North Carolina 27708, USA*<sup>18</sup>*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*<sup>19</sup>*University of Florida, Gainesville, Florida 32611, USA*<sup>20</sup>*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*<sup>21</sup>*University of Geneva, CH-1211 Geneva 4, Switzerland*<sup>22</sup>*Glasgow University, Glasgow G12 8QQ, United Kingdom*<sup>23</sup>*Harvard University, Cambridge, Massachusetts 02138, USA*

- <sup>24</sup>*Division of High Energy Physics, Department of Physics, University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland*
- <sup>25</sup>*University of Illinois, Urbana, Illinois 61801, USA*
- <sup>26</sup>*The Johns Hopkins University, Baltimore, Maryland 21218, USA*
- <sup>27</sup>*Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany*
- <sup>28</sup>*Center for High Energy Physics: Kyungpook National University, Daegu 702-701, Korea; Seoul National University, Seoul 151-742, Korea; Sungkyunkwan University, Suwon 440-746, Korea; Korea Institute of Science and Technology Information, Daejeon, 305-806, Korea; Chonnam National University, Gwangju, 500-757, Korea*
- <sup>29</sup>*Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*
- <sup>30</sup>*University of Liverpool, Liverpool L69 7ZE, United Kingdom*
- <sup>31</sup>*University College London, London WC1E 6BT, United Kingdom*
- <sup>32</sup>*Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain*
- <sup>33</sup>*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*
- <sup>34</sup>*Institute of Particle Physics: McGill University, Montréal, Canada H3A 2T8; and University of Toronto, Toronto, Canada M5S 1A7*
- <sup>35</sup>*University of Michigan, Ann Arbor, Michigan 48109, USA*
- <sup>36</sup>*Michigan State University, East Lansing, Michigan 48824, USA*
- <sup>37</sup>*University of New Mexico, Albuquerque, New Mexico 87131, USA*
- <sup>38</sup>*Northwestern University, Evanston, Illinois 60208, USA*
- <sup>39</sup>*The Ohio State University, Columbus, Ohio 43210, USA*
- <sup>40</sup>*Okayama University, Okayama 700-8530, Japan*
- <sup>41</sup>*Osaka City University, Osaka 588, Japan*
- <sup>42</sup>*University of Oxford, Oxford OX1 3RH, United Kingdom*
- <sup>43</sup>*University of Padova, Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, I-35131 Padova, Italy*
- <sup>44</sup>*LPNHE, Universite Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France*
- <sup>45</sup>*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*
- <sup>46</sup>*Istituto Nazionale di Fisica Nucleare Pisa, Universities of Pisa, Siena and Scuola Normale Superiore, I-56127 Pisa, Italy*
- <sup>47</sup>*University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA*
- <sup>48</sup>*Purdue University, West Lafayette, Indiana 47907, USA*
- <sup>49</sup>*University of Rochester, Rochester, New York 14627, USA*
- <sup>50</sup>*The Rockefeller University, New York, New York 10021, USA*
- <sup>51</sup>*Istituto Nazionale di Fisica Nucleare, Sezione di Roma I, University of Rome “La Sapienza,” I-00185 Roma, Italy*
- <sup>52</sup>*Rutgers University, Piscataway, New Jersey 08855, USA*
- <sup>53</sup>*Texas A&M University, College Station, Texas 77843, USA*
- <sup>54</sup>*Istituto Nazionale di Fisica Nucleare, University of Trieste/Udine, Italy*
- <sup>55</sup>*University of Tsukuba, Tsukuba, Ibaraki 305, Japan*
- <sup>56</sup>*Tufts University, Medford, Massachusetts 02155, USA*
- <sup>57</sup>*Waseda University, Tokyo 169, Japan*
- <sup>58</sup>*Wayne State University, Detroit, Michigan 48201, USA*
- <sup>59</sup>*University of Wisconsin, Madison, Wisconsin 53706, USA*
- <sup>60</sup>*Yale University, New Haven, Connecticut 06520, USA*
- <sup>61</sup>*University of California, Irvine, California 92617, USA*
- (Received 29 September 2008; published 27 January 2009)

Models of maximal flavor violation (MxFV) in elementary particle physics may contain at least one new scalar SU(2) doublet field  $\Phi_{\text{FV}} = (\eta^0, \eta^+)$  that couples the first and third generation quarks ( $q_1, q_3$ ) via a Lagrangian term  $\mathcal{L}_{\text{FV}} = \xi_{13} \Phi_{\text{FV}} q_1 q_3$ . These models have a distinctive signature of same-charge top-quark pairs and evade flavor-changing limits from meson mixing measurements. Data corresponding to  $2 \text{ fb}^{-1}$  collected by the Collider Dectector at Fermilab II detector in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96 \text{ TeV}$  are analyzed for evidence of the MxFV signature. For a neutral scalar  $\eta^0$  with  $m_{\eta^0} = 200 \text{ GeV}/c^2$  and coupling  $\xi_{13} = 1$ ,  $\sim 11$  signal events are expected over a background of  $2.1 \pm 1.8$  events. Three events are observed in the data, consistent with background expectations, and limits are set on the coupling  $\xi_{13}$  for  $m_{\eta^0} = 180\text{--}300 \text{ GeV}/c^2$ .

DOI: 10.1103/PhysRevLett.102.041801

PACS numbers: 13.85.Ni, 12.15.Ff, 12.60.Fr, 13.85.Qk

Measurements of low energy flavor-changing (FC) transitions, such as neutral meson mixing and rare  $B$  and  $K$  decays [1], largely confirm the minimal flavor violation

(MFV) ansatz of the standard model's quark-mixing matrix. This suggests that any new physics that couples quark flavors must either be well aligned with the standard model

couplings or mediated by particles that are too heavy to give observable deviations in current data. If the proposed new physics can be written in terms of a coupling matrix  $\xi_{ij}$  between quark flavors  $i$  and  $j$ , MFV imposes strict constraints on models that couple the top quark to lighter quarks, namely  $\xi_{31}, \xi_{13}, \xi_{32}, \xi_{23} \sim 0$ .

However, there is no strong theoretical motivation for an alignment between the flavor structure of the standard model and new physics. On the contrary, theories beyond the standard model typically predict large flavor-changing transitions if no additional symmetries are imposed. Moreover, new results on  $CP$ -violating asymmetries from  $B_s \rightarrow J/\psi\phi$  [2,3] suggest that new flavor structure beyond that of the CKM may be required [4].

A new class of scalar-mediated models has recently been proposed with a Lagrangian describing a scalar  $SU(2)$  doublet field  $\Phi_{FV} = (\eta^0, \eta^+)$  that couple left-handed quark fields of flavor  $i$  ( $Q_{iL}$ ) to right-handed up-type quark fields of flavor  $j$  ( $u_{jR}$ ) with a strength  $\xi_{ij}$  [5]

$$\mathcal{L}_{FV} = \xi_{ij} \bar{Q}_{iL} \tilde{\Phi}_{FV} u_{jR} + \text{H.c.} \quad (1)$$

This departs maximally from the MFV ansatz by allowing real  $\xi_{31}, \xi_{13} \sim 1$  or  $\xi_{32}, \xi_{23} \sim 1$  with all other terms zero. Contrary to previous understanding, these models are not excluded by current measurements, which constrain the products of terms in the coupling matrix, e.g.,  $\xi_{32} \cdot \xi_{31}$  [5], even with a light  $\eta^0$  mass of  $\mathcal{O}(200)$  GeV/ $c^2$ .

In the model investigated here ( $\xi_{31}, \xi_{13} \sim 1$ , called  $M_{X_{FV}1}$  in Ref. [5]), the  $\eta^0$  decays with equal probability to quark-antiquark pairs  $t + \bar{u}$  and  $\bar{t} + u$ . This leads to production of same-charge top-quark pairs in association with light-quark jets through the processes  $ug \rightarrow t\eta^0 \rightarrow tt\bar{u}$ ,  $ug \rightarrow tt\bar{u}$  ( $\eta^0$  exchange),  $u\bar{u} \rightarrow \eta^0\eta^0 \rightarrow tt\bar{u}\bar{u}$ , and  $uu \rightarrow tt$  ( $\eta^0$  exchange) and their Hermitian conjugates [6]. The predicted cross section is  $\mathcal{O}(1)$  pb over a range of light  $\eta^0$  masses, 180–300 GeV/ $c^2$ .

As suggested in Ref. [6], the case in which both  $W$  bosons decay leptonically ( $t \rightarrow Wb \rightarrow l\nu b$ ) presents an

experimentally attractive signature of two same-charge leptons, missing transverse energy ( $\cancel{E}_T$ ) [7,8], and at least one  $b$  jet ( $\ell^\pm \ell^\pm \cancel{E}_T b$ ) accompanied by additional jets. Though CDF has examined its inclusive same-charge lepton data set in smaller data subsets [9], and has looked for new physics in exclusive final states [10] there has not been an experimental study of the inclusive  $\ell^\pm \ell^\pm \cancel{E}_T b$  final state, where many of the contributions to a  $\ell^\pm \ell^\pm$  final state are suppressed by the requirement of  $b$  jet identification or large missing transverse energy. Thus, there may be new flavor-violating processes which have large cross sections, low backgrounds, and no direct experimental constraints. In addition, this final state is sensitive to a broad array of new physics topics which give same-charge top quarks, such as heavy down-type quarks [11] or scalar gluons [12].

Reference [6] used a parametrized detector description to describe the signal and background rates. This Letter estimates the dominant backgrounds using data-driven models, validates those estimates in control samples, calculates the signal acceptance with a realistic GEANT-based detector simulation [13], and reports the results of the first experimental search for particles that couple up and top quarks in this generic manner.

We use data collected between 2002 and 2007 with the CDF II detector, corresponding to an integrated luminosity of 2.0 fb $^{-1}$ . CDF II [8,14] is a general purpose detector designed to study  $p\bar{p}$  collisions at the Fermilab Tevatron. The tracking system consists of a cylindrical open-cell drift chamber and silicon microstrip detectors in a 1.4 T magnetic field parallel to the beam axis; the momentum resolution is  $\delta p_T/p_T^2 = 0.1\%/GeV/c$ . The silicon detectors provide tracking information for pseudorapidity  $|\eta| < 2$  and are used to reconstruct collision and decay points. The drift chamber surrounds the silicon detectors and gives full coverage in the central pseudorapidity region  $|\eta| < 1$ . Electromagnetic and hadronic calorimeters surround the tracking system and measure particle energies. Drift chambers and scintillators located outside the calorimeters detect muons in the central pseudorapidity region  $|\eta| < 1$ . The data used in this measurement are collected with

TABLE I. Cross sections  $\sigma(tt)$ ,  $\sigma(tt\bar{u})$ , and  $\sigma(tt\bar{u}\bar{u})$ , acceptance ( $\epsilon$ ) and number ( $N$ ) of selected  $\ell^\pm \ell^\pm b \cancel{E}_T$  events expected in the sample [6].

	$M_{\eta^0}$ [GeV/ $c^2$ ]	180	190	200	225	250	300
$tt$	$\sigma$ [pb]	0.50	0.45	0.41	0.33	0.27	0.19
	$\epsilon$ [%]	0.5	0.5	0.5	0.5	0.5	0.5
	$N$	4.4	4.3	3.8	2.6	2.1	0.9
$tt\bar{u}$	$\sigma$ [pb]	0.54	0.50	0.42	0.28	0.22	0.10
	$\epsilon$ [%]	0.5	0.5	0.5	0.5	0.5	0.5
	$N$	4.7	3.9	3.9	3.0	2.4	1.7
$tt\bar{u}\bar{u}$	$\sigma$ [pb]	0.68	0.45	0.38	0.17	0.06	0.02
	$\epsilon$ [%]	0.5	0.5	0.5	0.5	0.5	0.5
	$N$	5.7	3.5	3.3	1.4	0.5	0.2
Total $N(\ell^\pm \ell^\pm b \cancel{E}_T)$		14.7	11.7	10.9	7.0	5.0	2.7

TABLE II. Number of expected events for each background process, and the observed same-charge lepton events with  $b$ -tag and missing transverse energy, see text.

Source	$ee$	$\mu\mu$	$e\mu$	$\ell\ell$
$Z \rightarrow \ell\ell$	0.01	0.02	0.02	$0.1 \pm 0.1$
$t\bar{t}$	0.09	0.03	0.11	$0.2 \pm 0.1$
MisID	0.6	0.71	0.50	$1.8 \pm 1.8$
Total	0.7	0.8	0.6	$2.1 \pm 1.8$
Data	0	1	2	3

lepton triggers that require an electron or muon with  $p_T > 18 \text{ GeV}/c$ .

To isolate the same-charge top-quarks signal we follow [6] to define the  $\ell^\pm \ell^\pm b \cancel{E}_T$  signature by requiring two same-charge reconstructed leptons (electrons or muons) in the central region of the detector, each with  $p_T > 20 \text{ GeV}/c$ , one of which must be isolated [15]; at least one jet with  $|\eta| < 2.4$  and transverse energy of at least 15 GeV identified as a  $b$  jet by the SECVTX algorithm that searches for a displaced secondary vertex [16]; and at least 20 GeV of missing transverse energy,  $\cancel{E}_T$  [7].

To calculate the expected number of  $t\bar{t}$  and  $t\bar{t}$  events, we generate events for each of the same-charge processes using CALCHEP [17] with CTEQ6M proton parton distribution functions [18] followed by parton fragmentation and hadronization by PYTHIA [19]. Detector resolution and acceptance are modeled using the GEANT-based detector simulation, CDFSIM [13]. Table I shows the number of expected events in our sample.

Backgrounds to same-charge lepton pairs come from two classes of processes. In the first class, a real lepton is paired with a jet, which is misidentified as a same-charge lepton. The second class of processes comes from a pair of real opposite-charge leptons which include an electron; a hard photon emission from an electron converts into an electron-positron pair with strongly asymmetric momenta so that only one leg is reconstructed.

Backgrounds in which the second lepton arises from a misidentified jet or the decay of a heavy quark are largely due to production of  $W$  + jets or semileptonic  $t\bar{t}$  decays and are described using a model from jet data [20] in which the rate of lepton reconstruction in inclusive jets is measured and applied to  $W$  + jet events. The misidentification model is validated for light-quark jets by comparing the predicted and observed rates of same-charge events as a function of the missing transverse energy without a  $b$ -tag requirement. Discrepancies in rates in control regions motivate a 40% uncertainty. The selected sample may have a larger heavy-flavor fraction than the jets from which the lepton misidentification model was derived. Studies in simulated events show that the rate of misidentified leptons in a heavy-flavor enriched sample may be 50%–75% higher, and examination of the equivalent opposite-charge




FIG. 1 (color online). Following the prescriptions in Refs. [6,24], horizontal bands in measured  $\xi$  are shown which include 95% of simulated experiments, for various values of true  $\xi$ , with  $m_{\eta^0} = 180 \text{ GeV}/c^2$ .

sample motivate a 100% total uncertainty on the background prediction from lepton misidentification.

Backgrounds in which the same-charge lepton is due to a hard photon emission come from  $Z/\gamma^* + \text{jets}$  and top-quark pairs with electron-positron decays. Estimates of the backgrounds from  $Z/\gamma^* + \text{jets}$  processes are made with the ALPGEN [21] simulation code matched with PYTHIA in the MLM scheme [21] for the hadronization and fragmentation and normalized to data in opposite-sign events. To validate the modeling of the rate of hard emission, we compare our prediction for the contribution of  $Z \rightarrow e^+e^-$  to the observed sample of same-charge electrons or positron without a  $b$ -tag or missing transverse




FIG. 2. Jet multiplicity for background, observed data, and  $MxFV_1$  signal with best-fit value of  $\xi = 0.41$  for  $M_{\eta^0} = 180 \text{ GeV}/c^2$ .

TABLE III. 95% confidence level upper limits on the coupling  $\xi$  as a function of the mass of  $\eta^0$ .

Mass [GeV/ $c^2$ ]	180	190	200	225	250	300
(95% C.L.) $\xi <$	0.78	0.81	0.87	1.03	1.11	1.39

energy requirement. The shape and yield of the observed signal at the  $Z$  mass agrees well with the prediction. The  $t\bar{t}$  backgrounds are estimated using events generated in PYTHIA at  $m_t = 172$  GeV/ $c^2$ . Modeling of the  $t\bar{t}$  contribution is validated by comparing predicted and observed rates of opposite-sign leptons with  $\cancel{E}_T$  and a  $b$  tag, where  $t\bar{t}$  is expected to dominate. The detector response for both  $Z +$  jets and  $t\bar{t}$  processes is evaluated using CDFSIM, where, to avoid double-counting, the same-charge leptons are required to originate from the  $W$  or  $Z$  decays rather than from misidentified jets.


Backgrounds from charge mismeasurement are insignificant, as the charge of a particle with momentum of 100 GeV/ $c$  is typically determined with a significance greater than  $5\sigma$  [22]. The largest potential source comes from top-quark pair events, in which the lepton momenta are typically softer. Charge mismeasurement is very rare in this range, confirmed by the absence of a significant signal near the  $Z$  mass in observed same-charge muon events.

Backgrounds from diboson production  $WW$ ,  $WZ$ ,  $ZZ$ ,  $W\gamma$ , and  $Z\gamma$  in association with  $b$  jets are modeled with PYTHIA and BAUR [23] generators. These have non-negligible contributions to the inclusive same-charge lepton pair sample, but in the final selection they are insignificant due to the requirement of a  $b$  tag.

The final background estimate, shown in Table II, is 2.1 events with an uncertainty of 1.8 events.

From the observed number of events, one could directly measure the value of the  $Mx\text{FV}_1$  coupling  $\xi = \xi_{31} = \xi_{13}$ . As suggested in [6], the precision of a coupling measurement can be enhanced by simultaneously fitting for the number of signal and background events in the data. This exploits the different number of jets expected in signal and background events (see Fig. 2). Our likelihood fit for the number of signal events is binned in the number of reconstructed jets and takes into account that  $\sigma(ug \rightarrow t\eta^0 \rightarrow t\bar{t}\bar{u}) \propto \xi^2$  while  $\sigma(uu \rightarrow tt)$  and  $\sigma(u\bar{u} \rightarrow \eta^0\eta^0 \rightarrow t\bar{t}\bar{u}\bar{u}) \propto \xi^4$ . The fitted number of signal events can be transformed into an estimated value for  $\xi$ .

Following the Feldman-Cousins prescription [24], we use simulated experiments to construct bands which contain 95% of the fitted values of  $\xi$  at various true values of  $\xi$  for a mass of  $\eta^0$  (Fig. 1). The simulated experiments include fluctuations in the nuisance parameters, including the uncertainty in the jet energy scale, initial and final state radiation, parton distribution functions and signal and background normalization uncertainties. The confidence band in the space of the true  $\xi$  for an individual experiment is the intersection of a line drawn at the observed  $\xi$ .

FIG. 3 (color online). Observed 95% C.L. regions in the  $\xi - m_{\eta^0}$  plane.

We observe 3 events, in good agreement with the background expectation. The distribution of jets can be seen in Fig. 2 for the data as well as for the signal and background for the best-fit value of  $\xi = 0.41$  for  $m_{\eta^0} = 180$  GeV/ $c^2$ . As shown in Fig. 1, this corresponds to an upper limit  $\xi < 0.78$  at 95% C.L. Table III and Fig. 3 give upper limits on the value of the coupling  $\xi$  for  $m_{\eta^0} = 180\text{--}300$  GeV/ $c^2$ .

In conclusion, we find no evidence of the signature for maximal flavor violation, and set the first limits on the flavor-changing coupling between the top and up quark in such a model.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A. P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korean Science and Engineering Foundation and the Korean Research Foundation; the Science and Technology Facilities Council and the Royal Society, UK; the Institut National de Physique Nucleaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010, Spain; the Slovak R&D Agency; and the Academy of Finland.

<sup>a</sup>Visitor from University of Athens, 15784 Athens, Greece.<sup>b</sup>Visitor from Chinese Academy of Sciences, Beijing 100864, China.

- <sup>c</sup>Visitor from University of Bristol, Bristol BS8 1TL, United Kingdom.
- <sup>d</sup>Visitor from University Libre de Bruxelles, B-1050 Brussels, Belgium.
- <sup>e</sup>Visitor from University of California Irvine, Irvine, CA 92697, USA.
- <sup>f</sup>Visitor from University of California Santa Cruz, Santa Cruz, CA 95064, USA.
- <sup>g</sup>Visitor from Cornell University, Ithaca, NY 14853, USA.
- <sup>h</sup>Visitor from University of Cyprus, Nicosia CY-1678, Cyprus.
- <sup>i</sup>Visitor from University College Dublin, Dublin 4, Ireland.
- <sup>j</sup>Visitor from University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom.
- <sup>k</sup>Visitor from University of Heidelberg, D-69120 Heidelberg, Germany.
- <sup>l</sup>Visitor from Universidad Iberoamericana, Mexico D.F., Mexico.
- <sup>m</sup>Visitor from University of Manchester, Manchester M13 9PL, United Kingdom.
- <sup>n</sup>Visitor from Nagasaki Institute of Applied Science, Nagasaki, Japan.
- <sup>o</sup>Visitor from University de Oviedo, E-33007 Oviedo, Spain.
- <sup>p</sup>Visitor from Queen Mary, University of London, London, E1 4NS, United Kingdom.
- <sup>q</sup>Visitor from Texas Tech University, Lubbock, TX 79409, USA.
- <sup>r</sup>Visitor from IFIC(CSIC-Universitat de Valencia), 46071 Valencia, Spain.
- <sup>s</sup><http://www-cdf.fnal.gov>
- [1] W.-M. Yao *et al.*, J. Phys. G **33**, 1 (2006).
- [2] T. Aaltonen *et al.*, Phys. Rev. Lett. **100**, 161802 (2008).
- [3] V. Abazov *et al.*, Phys. Rev. Lett. **101**, 241801 (2008).
- [4] Z. Ligeti *et al.*, Phys. Rev. Lett. **97**, 101801 (2006).
- [5] S. Bar-Shalom and A. Rajaraman, Phys. Rev. D **77**, 095011 (2008).
- [6] S. Bar-Shalom *et al.*, Phys. Rev. D **78**, 033003 (2008).
- [7] Missing transverse energy,  $\cancel{E}_T$ , is defined as the magnitude of the vector  $-\sum_i E_T^i \vec{n}_i$  where  $E_T^i$  are the magnitudes of transverse energy contained in each calorimeter tower  $i$ , and  $\vec{n}_i$  is the unit vector from the interaction vertex to the tower in the transverse ( $x, y$ ) plane.  $\cancel{E}_T$  is further corrected for the energy of identified muons.
- [8] CDF uses a cylindrical coordinate system with the  $z$  axis along the proton beam axis. Pseudorapidity is  $\eta \equiv -\ln(\tan(\theta/2))$ , where  $\theta$  is the polar angle, and  $\phi$  is the azimuthal angle relative to the proton beam direction, while  $p_T = |p| \sin\theta$ ,  $E_T = E \sin\theta$ .
- [9] A. Abulencia *et al.*, Phys. Rev. Lett. **98**, 221803 (2007).
- [10] A. Abulencia *et al.*, Phys. Rev. D **78**, 012003 (2008).
- [11] R. Contino and G. Servant, J. High Energy Phys. **06** (2008) 026.
- [12] T. Plehn and T. Tait, arXiv:0810.3919.
- [13] T. Affolder *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **447**, 1 (2000).
- [14] D. Acosta *et al.*, Phys. Rev. D **71**, 032001 (2005).
- [15] A lepton is isolated if the total  $E_T$  within a cone  $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} \leq 0.4$ , minus the lepton  $E_T$  is  $< 10\%$  of the lepton  $E_T$ .
- [16] D. Acosta *et al.*, Phys. Rev. D **71**, 052003 (2005).
- [17] A. Pukhov *et al.*, Technical Report No. 98-41/542, INP MSU, 1998.
- [18] J. Pumplin *et al.*, J. High Energy Phys. **07** (2002) 012.
- [19] T. Sjostrand *et al.*, Comput. Phys. Commun. **238**, 135 (2001).
- [20] D. Acosta *et al.*, Phys. Rev. D **72**, 052003 (2005).
- [21] M. L. Mangano *et al.*, J. High Energy Phys. **07** (2003) 001.
- [22] A. Abulencia *et al.*, J. Phys. G **34**, 2457 (2007).
- [23] U. Baur and E. Berger, Phys. Rev. D **41**, 1476 (1990).
- [24] G. Feldman and R. Cousins, Phys. Rev. D **57**, 3873 (1998).