

Measurement of the Strong Coupling Constant from Inclusive Jet Production at the Tevatron $\bar{p}p$ Collider

T. Affolder,¹ H. Akimoto,² A. Akopian,³ M. G. Albrow,⁴ P. Amaral,⁵ D. Amidei,⁶ K. Anikeev,⁷ J. Antos,⁸ G. Apollinari,⁴ T. Arisawa,² A. Artikov,⁹ T. Asakawa,¹⁰ W. Ashmanskas,⁵ F. Azfar,¹¹ P. Azzi-Bacchetta,¹² N. Bacchetta,¹² H. Bachacou,¹ S. Bailey,¹³ P. de Barbaro,¹⁴ A. Barbaro-Galtieri,¹ V. E. Barnes,¹⁵ B. A. Barnett,¹⁶ S. Baroiant,¹⁷ M. Barone,¹⁸ G. Bauer,¹⁹ F. Bedeschi,²⁰ S. Belforte,²¹ W. H. Bell,²² G. Bellettini,²⁰ J. Bellinger,²³ D. Benjamin,²⁴ J. Bensinger,²⁵ A. Beretvas,⁴ J. P. Berge,⁴ J. Berryhill,⁵ A. Bhatti,³ M. Binkley,⁴ D. Bisello,¹² M. Bishai,⁴ R. E. Blair,²⁶ C. Blocker,²⁵ K. Bloom,⁶ B. Blumenfeld,¹⁶ S. R. Blusk,¹⁴ A. Bocci,³ A. Bodek,¹⁴ W. Bokhari,²⁷ G. Bolla,¹⁵ Y. Bonushkin,²⁸ D. Bortoletto,¹⁵ J. Boudreau,²⁹ A. Brandl,³⁰ S. van den Brink,¹⁶ C. Bromberg,³¹ M. Brozovic,²⁴ E. Brubaker,¹ N. Bruner,³⁰ E. Buckley-Geer,⁴ J. Budagov,⁹ H. S. Budd,¹⁴ K. Burkett,¹³ G. Busetto,¹² A. Byon-Wagner,⁴ K. L. Byrum,²⁶ S. Cabrera,²⁴ P. Calafiura,¹ M. Campbell,⁶ W. Carithers,¹ J. Carlson,⁶ D. Carlsmith,²³ W. Caskey,¹⁷ A. Castro,³² D. Cauz,²¹ A. Cerri,²⁰ A. W. Chan,⁸ P. S. Chang,⁸ P. T. Chang,⁸ J. Chapman,⁶ C. Chen,²⁷ Y. C. Chen,⁸ M.-T. Cheng,⁸ M. Chertok,¹⁷ G. Chiarelli,²⁰ I. Chirikov-Zorin,⁹ G. Chlachidze,⁹ F. Chlebana,⁴ L. Christofek,³³ M. L. Chu,⁸ Y. S. Chung,¹⁴ C. I. Ciobanu,³⁴ A. G. Clark,³⁵ A. P. Colijn,⁴ A. Connolly,¹ M. Convery,³ J. Conway,³⁶ M. Cordelli,¹⁸ J. Cranshaw,³⁷ R. Cropp,³⁸ R. Culbertson,⁴ D. Dagenhart,³⁹ S. D'Auria,²² F. DeJongh,⁴ S. Dell'Agnello,¹⁸ M. Dell'Orso,²⁰ S. Demers,³ L. Demortier,³ M. Deninno,³² P. F. Derwent,⁴ T. Devlin,³⁶ J. R. Dittmann,⁴ A. Dominguez,¹ S. Donati,²⁰ J. Done,⁴⁰ M. D'Onofrio,²⁰ T. Dorigo,¹³ N. Eddy,³³ K. Einsweiler,¹ J. E. Elias,⁴ E. Engels, Jr.,²⁹ R. Erbacher,⁴ D. Errede,³³ S. Errede,³³ Q. Fan,¹⁴ H.-C. Fang,¹ R. G. Feild,⁴¹ J. P. Fernandez,⁴ C. Ferretti,²⁰ R. D. Field,⁴² I. Fiori,³² B. Flaughner,⁴ G. W. Foster,⁴ M. Franklin,¹³ J. Freeman,⁴ J. Friedman,¹⁹ Y. Fukui,⁴³ I. Furic,¹⁹ S. Galeotti,²⁰ A. Gallas,^{13,*} M. Gallinaro,³ T. Gao,²⁷ M. Garcia-Sciveres,¹ A. F. Garfinkel,¹⁵ P. Gatti,¹² C. Gay,⁴¹ D. W. Gerdes,⁶ P. Giannetti,²⁰ P. Giromini,¹⁸ V. Glagolev,⁹ D. Glenzinski,⁴ M. Gold,³⁰ J. Goldstein,⁴ I. Gorelov,³⁰ A. T. Goshaw,²⁴ Y. Gotra,²⁹ K. Goulianos,³ C. Green,¹⁵ G. Grim,¹⁷ P. Gris,⁴ L. Groer,³⁶ C. Grosso-Pilcher,⁵ M. Guenther,¹⁵ G. Guillian,⁶ J. Guimaraes da Costa,¹³ R. M. Haas,⁴² C. Haber,¹ S. R. Hahn,⁴ C. Hall,¹³ T. Handa,⁴⁴ R. Handler,²³ W. Hao,³⁷ F. Happacher,¹⁸ K. Hara,¹⁰ A. D. Hardman,¹⁵ R. M. Harris,⁴ F. Hartmann,⁴⁵ K. Hatakeyama,³ J. Hauser,²⁸ J. Heinrich,²⁷ A. Heiss,⁴⁵ M. Herndon,¹⁶ C. Hill,¹⁷ K. D. Hoffman,¹⁵ C. Holck,²⁷ R. Hollebeek,²⁷ L. Holloway,³³ B. T. Huffman,¹¹ R. Hughes,³⁴ J. Huston,³¹ J. Huth,¹³ H. Ikeda,¹⁰ J. Incandela,^{4,†} G. Introzzi,²⁰ A. Ivanov,¹⁴ J. Iwai,² Y. Iwata,⁴⁴ E. James,⁶ M. Jones,²⁷ U. Joshi,⁴ H. Kambara,³⁵ T. Kamon,⁴⁰ T. Kaneko,¹⁰ K. Karr,³⁹ S. Kartal,⁴ H. Kasha,⁴¹ Y. Kato,⁴⁶ T. A. Keaffaber,¹⁵ K. Kelley,¹⁹ M. Kelly,⁶ D. Khazins,²⁴ T. Kikuchi,¹⁰ B. Kilminster,¹⁴ B. J. Kim,⁴⁷ D. H. Kim,⁴⁷ H. S. Kim,³³ M. J. Kim,⁴⁷ S. B. Kim,⁴⁷ S. H. Kim,¹⁰ Y. K. Kim,¹ M. Kirby,²⁴ M. Kirk,²⁵ L. Kirsch,²⁵ S. Klimenko,⁴² P. Koehn,³⁴ K. Kondo,² J. Konigsberg,⁴² A. Korn,¹⁹ A. Korytov,⁴² E. Kovacs,²⁶ J. Kroll,²⁷ M. Kruse,²⁴ S. E. Kuhlmann,²⁶ K. Kurino,⁴⁴ T. Kuwabara,¹⁰ A. T. Laasanen,¹⁵ N. Lai,⁵ S. Lami,³ S. Lammel,⁴ J. Lancaster,²⁴ M. Lancaster,¹ R. Lander,¹⁷ A. Lath,³⁶ G. Latino,²⁰ T. LeCompte,²⁶ A. M. Lee IV,²⁴ K. Lee,³⁷ S. Leone,²⁰ J. D. Lewis,⁴ M. Lindgren,²⁸ T. M. Liss,³³ J. B. Liu,¹⁴ Y. C. Liu,⁸ D. O. Litvintsev,⁴ O. Lobban,³⁷ N. Lockyer,²⁷ J. Loken,¹¹ M. Loreti,¹² D. Lucchesi,¹² P. Lukens,⁴ S. Lusin,²³ L. Lyons,¹¹ J. Lys,¹ R. Madrak,¹³ K. Maeshima,⁴ P. Maksimovic,¹³ L. Malferrari,³² M. Mangano,²⁰ M. Mariotti,¹² G. Martignon,¹² A. Martin,⁴¹ J. A. J. Matthews,³⁰ J. Mayer,³⁸ P. Mazzanti,³² K. S. McFarland,¹⁴ P. McIntyre,⁴⁰ E. McKigney,²⁷ M. Menguzzato,¹² A. Menzione,²⁰ P. Merkel,⁴ C. Mesropian,³ A. Meyer,⁴ T. Miao,⁴ R. Miller,³¹ J. S. Miller,⁶ H. Minato,¹⁰ S. Miscetti,¹⁸ M. Mishina,⁴³ G. Mitselmakher,⁴² Y. Miyazaki,⁴⁶ N. Moggi,³² E. Moore,³⁰ R. Moore,⁶ Y. Morita,⁴³ T. Moulik,¹⁵ M. Mulhearn,¹⁹ A. Mukherjee,⁴ T. Muller,⁴⁵ A. Munar,²⁰ P. Murat,⁴ S. Murgia,³¹ J. Nachtman,²⁸ V. Nagaslaev,³⁷ S. Nahn,⁴¹ H. Nakada,¹⁰ I. Nakano,⁴⁴ C. Nelson,⁴ T. Nelson,⁴ C. Neu,³⁴ D. Neuberger,⁴⁵ C. Newman-Holmes,⁴ C.-Y. P. Ngan,¹⁹ H. Niu,²⁵ L. Nodulman,²⁶ A. Nomerotski,⁴² S. H. Oh,²⁴ Y. D. Oh,⁴⁷ T. Ohmoto,⁴⁴ T. Ohsugi,⁴⁴ R. Oishi,¹⁰ T. Okusawa,⁴⁶ J. Olsen,²³ W. Orejudos,¹ C. Pagliarone,²⁰ F. Palmonari,²⁰ R. Paoletti,²⁰ V. Papadimitriou,³⁷ D. Partos,²⁵ J. Patrick,⁴ G. Pauletta,²¹ M. Paulini,^{1,‡} C. Paus,¹⁹ D. Pellett,¹⁷ L. Pescara,¹² T. J. Phillips,²⁴ G. Piacentino,²⁰ K. T. Pitts,³³ A. Pompos,¹⁵ L. Pondrom,²³ G. Pope,²⁹ M. Popovic,³⁸ F. Prokoshin,⁹ J. Proudfoot,²⁶ F. Ptohos,¹⁸ O. Pukhov,⁹ G. Punzi,²⁰ A. Rakitine,¹⁹ F. Ratnikov,³⁶ D. Reher,¹ A. Reichold,¹¹ P. Renton,¹¹ A. Ribon,¹² W. Riegler,¹³ F. Rimondi,³² L. Ristori,²⁰ M. Rivelino,³⁸ W. J. Robertson,²⁴ A. Robinson,³⁸ T. Rodrigo,⁴⁸ S. Rolli,³⁹ L. Rosenson,¹⁹ R. Roser,⁴ R. Rossin,¹² C. Rott,¹⁵ A. Roy,¹⁵ A. Ruiz,⁴⁸ A. Safonov,¹⁷ R. St. Denis,²² W. K. Sakumoto,¹⁴ D. Saltzberg,²⁸ C. Sanchez,³⁴ A. Sansoni,¹⁸ L. Santi,²¹ H. Sato,¹⁰ P. Savard,³⁸ P. Schlabach,⁴ E. E. Schmidt,⁴ M. P. Schmidt,⁴¹ M. Schmitt,^{13,*} L. Scodellaro,¹² A. Scott,²⁸ A. Scribano,²⁰ S. Segler,⁴ S. Seidel,³⁰ Y. Seiya,¹⁰ A. Semenov,⁹

F. Semeria,³² T. Shah,¹⁹ M.D. Shapiro,¹ P.F. Shepard,²⁹ T. Shibayama,¹⁰ M. Shimojima,¹⁰ M. Shochet,⁵ A. Sidoti,¹² J. Siegrist,¹ A. Sill,³⁷ P. Sinervo,³⁸ P. Singh,³³ A. J. Slaughter,⁴¹ K. Sliwa,³⁹ C. Smith,¹⁶ F.D. Snider,⁴ A. Solodsky,³ J. Spalding,⁴ T. Speer,³⁵ P. Sphicas,¹⁹ F. Spinella,²⁰ M. Spiropulu,⁵ L. Spiegel,⁴ J. Steele,²³ A. Stefanini,²⁰ J. Strologas,³³ F. Strumia,³⁵ D. Stuart,⁴ K. Sumorok,¹⁹ T. Suzuki,¹⁰ T. Takano,⁴⁶ R. Takashima,⁴⁴ K. Takikawa,¹⁰ P. Tamburello,²⁴ M. Tanaka,¹⁰ B. Tannenbaum,²⁸ M. Tecchio,⁶ R. Tesarek,⁴ P.K. Teng,⁸ K. Terashi,³ S. Tether,¹⁹ A. S. Thompson,²² R. Thurman-Keup,²⁶ P. Tipton,¹⁴ S. Tkaczyk,⁴ D. Toback,⁴⁰ K. Tollefson,¹⁴ A. Tollestrup,⁴ D. Tonelli,²⁰ H. Toyoda,⁴⁶ W. Trischuk,³⁸ J.F. de Troconiz,¹³ J. Tseng,¹⁹ N. Turini,²⁰ F. Ukegawa,¹⁰ T. Vaiculis,¹⁴ J. Valls,³⁶ S. Vejcik III,⁴ G. Velev,⁴ G. Veramendi,¹ R. Vidal,⁴ I. Vila,⁴⁸ R. Vilar,⁴⁸ I. Volobouev,¹ M. von der Mey,²⁸ D. Vucinic,¹⁹ R.G. Wagner,²⁶ R.L. Wagner,⁴ N.B. Wallace,³⁶ Z. Wan,³⁶ C. Wang,²⁴ M.J. Wang,⁸ B. Ward,²² S. Waschke,²² T. Watanabe,¹⁰ D. Waters,¹¹ T. Watts,³⁶ R. Webb,⁴⁰ H. Wenzel,⁴⁵ W.C. Wester III,⁴ A.B. Wicklund,²⁶ E. Wicklund,⁴ T. Wilkes,¹⁷ H.H. Williams,²⁷ P. Wilson,⁴ B.L. Winer,³⁴ D. Winn,⁶ S. Wolbers,⁴ D. Wolinski,⁶ J. Wolinski,³¹ S. Wolinski,⁶ S. Worm,³⁰ X. Wu,³⁵ J. Wyss,²⁰ W. Yao,¹ G.P. Yeh,⁴ P. Yeh,⁸ J. Yoh,⁴ C. Yosef,³¹ T. Yoshida,⁴⁶ I. Yu,⁴⁷ S. Yu,²⁷ Z. Yu,⁴¹ A. Zanetti,²¹ F. Zetti,¹ and S. Zucchelli³²

(CDF Collaboration)

¹Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720

²Waseda University, Tokyo 169, Japan

³Rockefeller University, New York, New York 10021

⁴Fermi National Accelerator Laboratory, Batavia, Illinois 60510

⁵Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637

⁶University of Michigan, Ann Arbor, Michigan 48109

⁷Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

⁸Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China

⁹Joint Institute for Nuclear Research, RU-141980 Dubna, Russia

¹⁰University of Tsukuba, Tsukuba, Ibaraki 305, Japan

¹¹University of Oxford, Oxford OX1 3RH, United Kingdom

¹²Universita di Padova, Istituto Nazionale di Fisica Nucleare, Sezione di Padova, I-35131 Padova, Italy

¹³Harvard University, Cambridge, Massachusetts 02138

¹⁴University of Rochester, Rochester, New York 14627

¹⁵Purdue University, West Lafayette, Indiana 47907

¹⁶The Johns Hopkins University, Baltimore, Maryland 21218

¹⁷University of California at Davis, Davis, California 95616

¹⁸Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy

¹⁹Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

²⁰Istituto Nazionale di Fisica Nucleare, University and Scuola Normale Superiore of Pisa, I-56100 Pisa, Italy

²¹Istituto Nazionale di Fisica Nucleare, University of Trieste/Udine, Italy

²²Glasgow University, Glasgow G12 8QQ, United Kingdom

²³University of Wisconsin, Madison, Wisconsin 53706

²⁴Duke University, Durham, North Carolina 27708

²⁵Brandeis University, Waltham, Massachusetts 02254

²⁶Argonne National Laboratory, Argonne, Illinois 60439

²⁷University of Pennsylvania, Philadelphia, Pennsylvania 19104

²⁸University of California at Los Angeles, Los Angeles, California 90024

²⁹University of Pittsburgh, Pittsburgh, Pennsylvania 15260

³⁰University of New Mexico, Albuquerque, New Mexico 87131

³¹Michigan State University, East Lansing, Michigan 48824

³²Istituto Nazionale di Fisica Nucleare, University of Bologna, I-40127 Bologna, Italy

³³University of Illinois, Urbana, Illinois 61801

³⁴The Ohio State University, Columbus, Ohio 43210

³⁵University of Geneva, CH-1211 Geneva 4, Switzerland

³⁶Rutgers University, Piscataway, New Jersey 08855

³⁷Texas Tech University, Lubbock, Texas 79409

³⁸Institute of Particle Physics, University of Toronto, Toronto M5S 1A7, Canada

³⁹Tufts University, Medford, Massachusetts 02155

⁴⁰Texas A&M University, College Station, Texas 77843

⁴¹Yale University, New Haven, Connecticut 06520

⁴²University of Florida, Gainesville, Florida 32611

⁴³High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305, Japan

⁴⁴*Hiroshima University, Higashi-Hiroshima 724, Japan*

⁴⁵*Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany*

⁴⁶*Osaka City University, Osaka 588, Japan*

⁴⁷*Center for High Energy Physics, Kyungpook National University, Taegu 702-701, Korea
and Seoul National University, Seoul 151-742, Korea
and SungKyunKwan University, Suwon 440-746, Korea*

⁴⁸*Instituto de Fisica de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain
(Received 29 August 2001; published 9 January 2002)*

We report a measurement of the strong coupling constant, $\alpha_s(M_Z)$, extracted from inclusive jet production in $p\bar{p}$ collisions at $\sqrt{s} = 1800$ GeV. The QCD prediction for the evolution of α_s with jet transverse energy E_T is tested over the range $40 < E_T < 450$ GeV using E_T for the renormalization scale. The data show good agreement with QCD in the region below 250 GeV. The value of α_s at the mass of the Z^0 boson averaged over the range $40 < E_T < 250$ GeV is found to be $\alpha_s(M_Z) = 0.1178 \pm 0.0001(\text{stat})_{-0.0095}^{+0.0081}(\text{expt. syst.})$. The associated theoretical uncertainties are mainly due to the choice of renormalization scale (${}_{-4\%}^{+6\%}$) and input parton distribution functions (5%).

DOI: 10.1103/PhysRevLett.88.042001

PACS numbers: 12.38.Qk, 13.87.Ce

Jet production at hadron colliders provides an excellent opportunity for testing the theory of strong interactions, quantum chromodynamics (QCD) [1]. QCD has achieved remarkable success in describing hadron interactions at short distances (large momentum transfers), owing to the property of asymptotic freedom [2]. Asymptotic freedom predicts a logarithmic decrease of the coupling strength, $\alpha_s(\mu)$, as the momentum scale μ characterizing a process increases. Processes with large momentum transfer can then be described by an expansion in powers of $\alpha_s(\mu)$. The value of α_s , a free parameter of QCD, is one of the fundamental constants of nature. Its determination is the essential measurement of QCD, and the observation of its evolution, or *running*, with momentum transfer is one of the key tests of the theory. At e^+e^- colliders α_s has been measured from the fragmentation functions [3], event shapes [4], jet production rates [5], and τ lepton decay properties [6]. In lepton-hadron collisions, α_s has been measured from scaling violations [7], jet production rates [8], and momentum sum rules [9]. A precise value for α_s has also been obtained from a global fit to properties of the Z^0 boson measured at the CERN LEP and the SLAC SLC e^+e^- colliders and the W boson and top quark masses [10]. A review of these and other α_s measurements can be found in [11]. In this Letter, we present a measurement of α_s from the inclusive jet cross section in $p\bar{p}$ collisions over the jet transverse energy (E_T) range from 40 to 450 GeV.

This measurement is based on a data sample of integrated luminosity 87 pb^{-1} collected by the Collider Detector at Fermilab (CDF) during the 1994-1995 run (Run 1b) of the Fermilab Tevatron $p\bar{p}$ collider operating at $\sqrt{s} = 1.8$ TeV. The CDF detector is described elsewhere [12]. Details of the measurement of the inclusive jet differential cross section can be found in [13]. Briefly, jets are reconstructed using an iterative fixed cone algorithm with a radius $R = (\Delta\eta^2 + \Delta\phi^2)^{1/2} = 0.7$, where $\eta \equiv -\ln(\tan\frac{\theta}{2})$ is the pseudorapidity, evaluated from the angle θ between the centerline of the jet cone and the proton beam line, and ϕ is the azimuthal angle. The inclusive jet

cross section includes all jets in an event in the pseudo-rapidity range $0.1 < |\eta| < 0.7$. The measured spectrum is corrected for the calorimeter response, resolution, and the underlying event energy using an iterative unsmearing procedure which changes both the energy scale and the normalization simultaneously. The value of α_s is determined by comparing the jet cross section with the next to leading order (NLO) perturbative QCD predictions [14]. In the E_T region studied, the nonperturbative contributions to the inclusive jet cross section are estimated to be negligible [15]. The procedure of extracting α_s can be summarized by the equation

$$\frac{d\sigma}{dE_T} = \alpha_s^2(\mu_R)\hat{X}^{(0)}(\mu_F, E_T) \times [1 + \alpha_s(\mu_R)k_1(\mu_R, \mu_F, E_T)], \quad (1)$$

where $\frac{d\sigma}{dE_T}$ is the transverse energy distribution of the inclusive jets, μ_R and μ_F , related to E_T by a scale factor, are the renormalization and factorization scales, $\alpha_s^2(\mu_R)\hat{X}^{(0)}(\mu_F, E_T)$ is the leading order (LO) prediction for the inclusive jet cross section, and $\alpha_s^3(\mu_R)\hat{X}^{(0)}(\mu_F, E_T)k_1(\mu_R, \mu_F, E_T)$ is the NLO contribution. Both $\hat{X}^{(0)}(\mu_F, E_T)$ and $k_1(\mu_R, \mu_F, E_T)$ are calculated with the JETRAD Monte Carlo program [16] based on the techniques described in [17] and the matrix elements of [18]. NLO QCD predictions for the inclusive jet cross section are also available in [19,20] and agree well with those of JETRAD. All calculations are performed in the modified minimal subtraction, $\overline{\text{MS}}$, scheme [21]. The JETRAD Monte Carlo program generates events with weighting factors, so that the jet clustering algorithm and E_T and η cuts, mimicking the experimental requirements, are directly applied to the final state partons. The jet clustering in JETRAD is governed by a cone radius R and a phenomenological parameter \mathcal{R}_{sep} (default value = 1.3), introduced to match the experimental efficiency of identifying overlapping jets [15]. If two partons are more than

$\mathcal{R}_{\text{sep}} \times R$ apart, they are identified as two distinct jets; otherwise they are merged into a single jet.

The inclusive jet data are divided into 33 E_T bins, from which we obtain statistically independent measurements of α_s for 33 different values of μ_R . The α_s values derived for $\mu_R = \mu_F = E_T$ using CTEQ4M [22] parton distribution functions (PDFs) are presented in Fig. 1. For $E_T < 250$ GeV, there is good agreement with QCD predictions for the running of the coupling constant. The behavior of α_s at higher E_T values is a direct reflection of the excess observed in the inclusive jet cross section [13]. The discrepancy with the NLO QCD predictions in this region, though not well understood, may be accommodated by the flexibility allowed by the world data in determining the high- x gluon component in the parton distributions [22].

The measured values of $\alpha_s(\mu_R)$ are evolved to the mass of the Z^0 boson, M_Z , by using the solution to the two-loop renormalization group equation:

$$\alpha_s(M_Z) = \frac{\alpha_s(\mu_R)}{1 - \alpha_s(\mu_R)[b_0 + b_1\alpha_s(\mu_R)]\ln(\mu_R/M_Z)}, \quad (2)$$

$$b_0 = \frac{33 - 2n_f}{6\pi} \quad b_1 = \frac{306 - 38n_f}{24\pi^2}, \quad (3)$$

where n_f is the number of active flavors, which is equal to five (six) for μ_R smaller (larger) than the top quark mass. The values of $\alpha_s(M_Z)$ for all 33 measurements are




FIG. 1. The strong coupling constant as a function of E_T for $\mu_R = E_T$ measured using CTEQ4M parton distributions. The shaded area shows the experimental systematic uncertainties. The curved line represents the NLO QCD prediction for the evolution of $\alpha_s(E_T)$ using $\alpha_s(M_Z) = 0.1178$, the average value obtained in the region $40 < E_T < 250$ GeV. The $\alpha_s(M_Z)$ extracted from $\alpha_s(E_T)$ is shown in the inset along with the weighted average as the horizontal line.

shown in the inset of Fig. 1. Averaging over the range 40-250 GeV, we obtain

$$\alpha_s(M_Z) = 0.1178 \pm 0.0001(\text{stat}).$$

Inclusion of the data with $E_T > 250$ GeV results in an increase of the average value by 0.0001.

The running of α_s is tested by verifying if $\alpha_s(M_Z)$ is independent of the energy scale E_T at which the jet cross section is measured. The 27 values of $\alpha_s(M_Z)$ obtained from the data in the jet E_T range 40–250 GeV are fit to the linear function $P_0 + P_1 \times E_T$. The fit yields $P_0 = 0.1173 \pm 0.0005$ and $P_1 = (0.3 \pm 0.3) \times 10^{-5} \text{ GeV}^{-1}$ with $\chi^2/\text{d.o.f.} = 1.3$, showing that $\alpha_s(M_Z)$ is independent of E_T within one standard deviation. The estimated experimental systematic uncertainty on the value of P_1 is $\pm 5.0 \times 10^{-5} \text{ GeV}^{-1}$. The nontrivial result of this fit demonstrates the correctness of the QCD prediction for the evolution of α_s over the above range.

The experimental systematic uncertainties on the value of $\alpha_s(M_Z)$ are derived from those on the inclusive jet cross section. For each source of systematic uncertainty described below, except normalization, the inclusive jet cross section was reevaluated by varying the corresponding parameter in the detector response by 1σ . For the normalization uncertainty, it was changed by a scale factor [13]. These uncertainties were propagated to $\alpha_s(M_Z)$ by repeating the procedure described above using the spectra given in Table VI of Ref. [13]. The results are shown in Fig. 2. The deviations of $\alpha_s(M_Z)$ for each spectrum from the central value are given in Table I. The dominant experimental systematic uncertainty in the inclusive jet cross section measurement is due to the calorimeter response to jets.




FIG. 2. Experimental systematic uncertainties for α_s measurement (the lines are 1 standard deviation contours), with CTEQ4M as input PDF and $\mu_R = \mu_F = E_T$.

TABLE I. Experimental systematic uncertainties on $\alpha_s(M_Z)$ extracted using CTEQ4M parton distribution functions.

Source of uncertainty	$\Delta\alpha_s$	$(\frac{\Delta\alpha_s}{\alpha_s})\%$
Calorimeter high P_T pion response	+0.0036	+3.1
	-0.0055	-4.7
Calorimeter low P_T pion response	+0.0027	+2.3
	-0.0033	-2.8
Energy scale stability	+0.0030	+2.6
	-0.0030	-2.6
Neutral pion response	+0.0016	+1.4
	-0.0021	-1.8
Underlying event energy	+0.0025	+2.1
	-0.0027	-2.3
Jet fragmentation functions	+0.0046	+3.9
	-0.0044	-3.7
Jet energy resolution	+0.0015	+1.3
	-0.0017	-1.4
Normalization	+0.0022	+2.0
	-0.0023	-1.9

The detector response and jet energy corrections are derived from a combination of test-beam data and Monte Carlo simulations. The calorimeter response to charged pions was evaluated separately for high and low transverse momentum (P_T) pions from test-beam data and isolated charged tracks from $p\bar{p}$ data with an uncertainty of $\pm 5\%$ for $P_T \leq 5$ GeV, $\pm 3\%$ for $5 \text{ GeV} < P_T < 15$ GeV and $^{+3.6}_{-2.0}\%$ for $P_T \geq 15$ GeV. During the run, the calorimeter response was monitored by using muons, isolated particles, and the measured inclusive jet cross section. The response was found to be stable within $\pm 1\%$. The electromagnetic calorimeter was calibrated using electrons from $p\bar{p}$ interactions. The associated uncertainty, labeled in Fig. 2 as neutral pion response, arises from the modeling of calorimeter response to very low energy electrons. The underlying event energy (nonjet energy contribution to the jet E_T) was measured from minimum bias data, and its mean value was varied by $\pm 30\%$ [13] to evaluate the effect on the jet cross section. The error from the jet fragmentation functions is due to the extrapolation of the track momentum and multiplicity distribution to the high E_T region and from uncertainties in the track reconstruction efficiency. The detector jet energy response has a Gaussian shape with exponential tails on both the high and low sides and a resolution with an uncertainty of $\pm 10\%$. Finally, the overall normalization of the inclusive jet cross section has an uncertainty of $\pm 4.5\%$, dominated by the contribution from the total cross section measurement. Summing in quadrature all the above uncertainties after propagation to $\alpha_s(M_Z)$ yields a total experimental systematic uncertainty of $\pm \begin{matrix} 0.0081 \\ 0.0095 \end{matrix}$.

The theoretical uncertainties are mainly due to the choice of renormalization and factorization scales and parton distribution functions. The scales μ_F and μ_R are expected to be of the same order as the characteristic scale of the process, which in this case is the jet E_T . We have evaluated the changes in $\alpha_s(M_Z)$ resulting

from independently varying μ_F and μ_R from $E_T/2$ to $2E_T$ and found that the largest changes occur for $\mu_R = \mu_F$. For all results presented in this Letter the two scales were set equal. The sensitivity of the measured $\alpha_s(M_Z)$ to changes in these scales is indicated by the shaded band in Fig. 3(a). Over the E_T range from 40 to 250 GeV, the shift in $\alpha_s(M_Z)$ induced by changing the scale from $E_T/2$ [$\alpha_s(M_Z) = 0.1129 \pm 0.0001$] [23] to $2E_T$ [$\alpha_s(M_Z) = 0.1249 \pm 0.0001$] is approximately $^{+6\%}_{-4\%}$, independently of E_T .

The coefficients \hat{X}_0 and k_1 in Eq. (1) depend on the PDFs, which are obtained from global fits to deep inelastic scattering (DIS), Drell-Yan production, and other collider data, including the inclusive jet results from Tevatron. Each PDF set has an associated strong coupling constant, α_s^{PDF} . The gluon PDF [$G(x)$] determined from the fit is highly correlated with α_s^{PDF} because, in equations describing all the processes used, the $G(x)$ is always accompanied by α_s^{PDF} . To calculate the above coefficients, the PDFs are evolved using α_s^{PDF} . For this procedure of measuring α_s to be valid, the extracted value of α_s should be consistent with the input α_s^{PDF} , although not necessarily equal. The variation in parton distributions, especially in the gluon distribution, allowed by the world data was studied by the CTEQ Collaboration by fixing the value of α_s^{PDF} to 0.110, 0.113, 0.116, 0.119, or 0.122, with resulting χ^2 of 1388, 1323, 1323, 1388, or 1543 for 1297 nonjet data points [22]. We use the CTEQ4A series to study the $\alpha_s(M_Z)$ dependence on the PDFs. In addition, we have studied $\alpha_s(M_Z)$ using PDF sets which do not include Tevatron jet results, such as the MRST(g ↑) set [24], the MRSA' series [25], and two MRS-R sets [26]. The χ^2 , calculated by comparing the data with the theoretical prediction in




FIG. 3. Uncertainties in $\alpha_s(M_Z)$ due to the renormalization scale μ , (a), and parton distribution functions, (b).

the restricted range of 40–250 GeV, is used to quantify the agreement. The minimal $\chi^2/\text{d.o.f.} = 1.38$ is obtained for CTEQ4M ($\alpha_s^{\text{PDF}} = 0.116$), therefore we use this PDF in our final fit. Excluding the PDFs which have obvious disagreement ($\chi^2/\text{d.o.f.} \geq 5$), we estimate the uncertainty on the $\alpha_s(M_Z)$ due to the PDF choice to be $\pm 5\%$.

Finally, the variation of \mathcal{R}_{sep} , the jet clustering parameter, from 1.3 to 2.0 results in a 5%–7% normalization change of the inclusive jet cross section. The corresponding variation in the $\alpha_s(M_Z)$ measurement is 2%–3%.

To explore the flexibility in the gluon distribution at high E_T , a special PDF set, CTEQ4HJ, was generated by including CDF jet data in the global fit with higher statistical weight assigned to high E_T points and a new parametrization of the gluon distribution [22]. This PDF yields good agreement between Tevatron data and theoretical predictions. Using this set, we obtain $\alpha_s(M_Z) = 0.1185 \pm 0.0001$, averaged over the entire E_T range.

In conclusion, we have tested the evolution of the strong coupling constant α_s using the inclusive jet cross section data from $\bar{p}p$ collisions at $\sqrt{s} = 1800$ GeV. Our results demonstrate that for E_T in the range of 40–250 GeV with $\mu_R = E_T$ the running of α_s is in good agreement with QCD predictions. The value of α_s expressed at the Z^0 boson mass is found to be

$$\alpha_s(M_Z) = 0.1178 \pm 0.0001(\text{stat})_{-0.0095}^{+0.0081}(\text{expt. syst}).$$

This value is in good agreement with the world average $\alpha_s(M_Z) = 0.1181 \pm 0.0020$ [27]. The theoretical uncertainties associated with the choice of parton distribution functions ($\sim 5\%$) and the choice of the renormalization scale ($_{-4\%}^{+6\%}$) are comparable to the experimental systematic uncertainty.

We thank W. Giele for providing the JETRAD program and for helpful discussions and comments on all stages of this work. We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Science, Sports and Culture of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium fuer Bildung und Forschung, Germany; the Korea Science and Engineering Foundation (KoSEF); the Korea Research Foundation; and the Comision Interministerial de Ciencia y Tecnologia, Spain.

*Now at Northwestern University, Evanston, Illinois 60208.

†Now at University of California, Santa Barbara, California 93106.

‡Now at Carnegie Mellon University, Pittsburgh, PA 15213.

- [1] H. Fritzsch, M. Gell-Mann, and H. Leutwyler, Phys. Lett. **47B**, 365 (1973).
- [2] G. 't Hooft (unpublished); D. Gross and F. Wilczek, Phys. Rev. Lett. **30**, 1343 (1973); H. D. Politzer, Phys. Rev. Lett. **30**, 1346 (1973).
- [3] P. Abreu *et al.*, Phys. Lett. B **398**, 194 (1997).
- [4] P. Abreu *et al.*, Eur. Phys. J. C **14**, 557 (2000).
- [5] P. Pfeifenschneider *et al.*, Eur. Phys. J. C **17**, 19 (2000).
- [6] T. Coan *et al.*, Phys. Lett. B **356**, 580 (1995); R. Barate *et al.*, Eur. Phys. J. C **4**, 409 (1998); K. Ackerstaff *et al.*, Eur. Phys. J. C **7**, 571 (1999).
- [7] J. Santiago and F. J. Yndurain, Nucl. Phys. (Proc. Suppl.) **86**, 69 (2000); C. Adloff *et al.*, Eur. Phys. J. C **21**, 33 (2001).
- [8] J. Breitweg *et al.*, Phys. Lett. B **507**, 70 (2001); C. Adloff *et al.*, Eur. Phys. J. C **19**, 269 (2001).
- [9] J. H. Kim *et al.*, Phys. Rev. Lett. **81**, 3595 (1998).
- [10] D. Abbaneo *et al.*, CERN-EP/2001-021 (2001).
- [11] I. Hinchliffe and A. V. Manohar, Annu. Rev. Nucl. Part. Sci. **50**, 643 (2000).
- [12] F. Abe *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **271**, 387 (1988).
- [13] T. Affolder *et al.*, Phys. Rev. D **64**, 032001 (2001).
- [14] W. Giele, E. W. N. Glover, and J. Yu, Phys. Rev. D **53**, 120 (1996).
- [15] S. D. Ellis, Z. Kunszt, and D. E. Soper, Phys. Rev. Lett. **69**, 3615 (1992).
- [16] W. Giele, E. W. N. Glover, and D. A. Kosower, Phys. Rev. Lett. **73**, 2019 (1994).
- [17] W. Giele, E. W. N. Glover, and D. A. Kosower, Nucl. Phys. **B403**, 633 (1993).
- [18] R. K. Ellis and J. Sexton, Nucl. Phys. **B269**, 445 (1986).
- [19] S. D. Ellis, Z. Kunszt, and D. E. Soper, Phys. Rev. Lett. **64**, 2121 (1990).
- [20] F. Aversa *et al.*, Phys. Lett. B **211**, 465 (1988).
- [21] W. A. Bardeen *et al.*, Phys. Rev. D **18**, 3998 (1978).
- [22] H. L. Lai *et al.*, Phys. Rev. D **55**, 1280 (1997).
- [23] The $\alpha_s(M_Z)$ value quoted in the preprint by Mesropian [FERMILAB-CONF-99-177-E (1999)] was obtained with $\mu_R = E_T/2$; however, for this Letter we repeated the analysis with $\mu_R = E_T$.
- [24] A. D. Martin *et al.*, Eur. Phys. J. C **4**, 463 (1998).
- [25] A. D. Martin, R. Roberts, and W. J. Stirling, Phys. Lett. B **356**, 89 (1995).
- [26] A. D. Martin, R. Roberts, and W. J. Stirling, Phys. Lett. B **387**, 419 (1996).
- [27] Particle Data Group, D. E. Groom *et al.*, Eur. Phys. J. C **15**, 1 (2000).