

## **Work Stress and Coronary Heart Disease- What are the Mechanisms?**

Tarani Chandola<sup>1</sup>, DPhil. t.chandola@ucl.ac.uk      Corresponding author

Annie Britton<sup>1</sup>, PhD a.britton@ucl.ac.uk

Eric Brunner<sup>1</sup>, PhD e.brunner@ucl.ac.uk

Harry Hemingway<sup>1</sup> FRCP h.hemingway@ucl.ac.uk

Marek Malik<sup>2</sup> PhD, MD marek.malik@btinternet.com

Meena Kumari m.kumari@ucl.ac.uk

Ellena Badrick e.badrick@ucl.ac.uk

Mika Kivimaki, PhD m.kivimaki@ucl.ac.uk

Michael Marmot<sup>1</sup>, PhD m.marmot@ucl.ac.uk

<sup>1</sup>Department of Epidemiology and Public Health

University College London

Tel: +44 20 7679 5629

Fax: +44 20 7813 0242

<sup>2</sup>Department of Cardiac and Vascular Sciences, St. George's University of London,

London, UK

Word Count:3910

The Whitehall II study has been supported by grants from the Medical Research Council; Economic and Social Research Council; British Heart Foundation; Health and Safety Executive; Department of Health; National Heart Lung and Blood Institute (HL36310), US, NIH: National Institute on Aging (AG13196), US, NIH; Agency for Health Care Policy Research (HS06516); and the John D and Catherine T MacArthur Foundation Research Networks on Successful Midlife Development and Socio-economic Status and Health. Michael Marmot is supported by an MRC Research Professorship, Harry Hemingway by a public health career scientist award from the Department of Health, and Mika Kivimaki by the Academy of Finland (grant 117 604).

## **Work Stress and Coronary Heart Disease- What are the Mechanisms?**

**Aims-** To determine the biological and behavioural factors linking work stress with CHD.

**Methods and Results—**10,308 London-based male and female civil servants aged 35-55 at phase 1 (1985-1988) of the Whitehall II study. Exposures included work stress (assessed at phases 1 and 2). Outcomes included behavioural risk factors (phase 3), the metabolic syndrome (phase 3), heart rate variability, morning rise in cortisol (phase 7) and incident CHD (phases 2 to 7) based on CHD death, non-fatal myocardial infarction or definite angina. Chronic work stress was associated with CHD and this association was stronger among participants aged under 50 (RR 1.68, 95% CI 1.17-2.42). There were similar associations between work stress and low physical activity, poor diet, the metabolic syndrome, its components, and lower heart rate variability. Cross-sectionally, work stress was associated with a higher morning rise in cortisol. Around 32% of the effect of work stress on CHD was attributable to its effect on health behaviours and the metabolic syndrome.

**Conclusions--**Work stress may be an important determinant of CHD among working age populations which is mediated through indirect effects on health behaviours and direct effects on neuroendocrine stress pathways.

## **Introduction**

Stress at work is associated with an increased risk of coronary heart disease (CHD) but the mechanisms underlying this association remain unclear. (1) Work stress may affect CHD through direct activation of neuroendocrine responses to stressors, or more indirectly through unhealthy behaviours which increase the risk of CHD, such as smoking, lack of exercise or excessive alcohol consumption. One of the main axes of neuroendocrine stress responses is the autonomic nervous system (ANS). Repeated activation of the ANS is characterised by lowered heart rate variability which has been associated with work stress among men in cross sectional studies (2), (3) Furthermore, work stress may affect dysregulation of the hypothalamic-pituitary-adrenal (HPA) axis, (4) which is associated with disturbances in the circadian rhythm of cortisol and the development of the metabolic syndrome. (5),(6)

Accumulation of work stress is associated with higher risks of the metabolic syndrome,(7) and incident obesity.(8) However, there are few longitudinal studies examining the effect of cumulative work stress on other intermediate mechanisms, despite evidence that chronic stress predicts cardiovascular mortality and morbidity.(9) It is important to examine cumulative exposures in order to show "dose-response" relations,(10) which would contribute a causal understanding of the association between work stress and CHD. In addition, there is little longitudinal evidence on the mechanisms by which work stress affects CHD. Stronger associations between work stress and CHD risk among working age populations would also increase the specificity of this association.

This study addresses the following questions: (1) Is the accumulation of work stress associated with higher risks of incident CHD and risk factors? (2) Is this association

stronger among working age populations? (3) Does work stress affect CHD directly through neuroendocrine mechanisms and/or indirectly through behavioural risk factors for CHD?

## **Methods**

### *Study Sample and Design*

The Whitehall II study conducted in 1985-1988 (phase 1) recruited 10,308 participants from 20 civil service departments in London. After initial participation, data collection was carried out in 1989-90 (phase 2), 1991-1993 (phase 3) 1995 (phase 4), 1997-1999 (phase 5), 2001 (phase 6) and 2002-04 (phase 7). Phases 2, 4 and 6 were postal questionnaires, while phases 3, 5 and 7 also included a clinical examination. Full details of the clinical examinations are reported elsewhere. (11) Ethical approval for the Whitehall II study was obtained from the University College London Medical School Committee on the ethics of human research. Informed consent was obtained from the study participants.

### *Assessment of Work Stress*

Self-reported *work stress* was measured by the job-strain questionnaire. (12) Participants report job-strain when their responses to the job demands questions are high and decision latitude (job control) questions are low (defined as being above or below the median score for the measures of job demands and decision latitude). In addition, participants are said to have iso-strain when they report job-strain and are socially isolated at work (i.e., without supportive coworkers or supervisors). (13), (14), (7) A cumulative measure of work stress was created by adding together the number of times the participant reported iso-strain at phases 1 and 2 (range 0-2), giving us a measure on the duration of exposure to work stress, albeit measured on two occasions only. Participants who lacked work stress data at either phase were

assigned a missing value. The prevalence of work stress (iso-strain) was lowest in the highest civil service grade.

#### *Follow-up Measurements*

CHD events included fatal CHD (ICD9 codes 410-414 or ICD10 I20-25) or incident non fatal myocardial infarction (MI) from phases 2-7 (an average of 12 years of follow up), with or without angina. Non-fatal MI was defined following MONICA criteria (15) based on study electrocardiograms, hospital acute ECGs and cardiac enzymes and excluded participants with existing MI at either phases 1 or 2. Incident angina were defined on the basis of clinical records and nitrate medication use, excluding cases based solely on self-reported data without clinical verification and participants with definite angina at either phases 1 or 2.

Biological risk factors for CHD included the ATPIII(16) metabolic syndrome measured at phase 3, its components (waist circumference: men>102cm women>88cm, serum triglycerides:  $\geq 150$ mg/dL, HDL cholesterol: men<40mg/dL, women<50mg/dL, blood pressure:  $\geq 130/\geq 85$  mmHg or on antihypertensive medication, fasting glucose:  $\geq 110$ mg/dL), morning rise in cortisol and low heart rate variability (both measured at phase 7).

For evaluation of heart rate variability, 5-minute of RR interval data were collected and analyzed both in the time domain (standard deviation of all intervals between normal-to-normal sinus rhythm R waves [SDNN]) and in the frequency domains: low frequency (LF), 0.04 to 0.15 Hz (in  $ms^2$ ) and high frequency (HF), 0.15 to 0.4 Hz (in  $ms^2$ ). These measures were log transformed to obtain a more normal distribution for the regression analyses.

For evaluation of cortisol participants were asked to provide samples of saliva collected at waking and 30 minutes after waking. Participants were asked to record time of waking. Samples were posted back and stored at -80°C for subsequent hormone analysis. Cortisol was measured as previously described.(17) Morning rise in cortisol was calculated as the difference between cortisol levels at waking and 30 minutes post waking.

Behavioural risk factors (at phase 3) for CHD included alcohol, smoking, activity and diet. Alcohol consumption in the previous week was categorised into non-drinker, recommended (1-14 units for women/1-21 units for men) and unsafe (14+ units for women/21+ units for men). Cigarette smoking categories were non-smoker, ex-smoker, 1-9 cigs/day, 10-19 cigs/day and 20+ cigs/day. Physical activity was measured by self-reported frequency of moderate activities (3+ times a week, at least once a week, at least once a month, never). Diet was measured by self-reported fruit or vegetable consumption (less than weekly, less than daily and at least daily). For logistic regression analyses, these health behaviours were coded into binary variables of current vs. never/ex smokers, unsafe drinkers vs. non/recommended limit drinkers, less than daily fruit/veg consumption vs. daily, and no physical activity vs. some activity.

#### *Missing Data and Statistical Methods*

There were 10,308 civil servants who participated in the baseline (phase 1) study. By phase 7, of the 9,692 participants still alive, 6,484 attended the clinical examination, 71% on whom we measured heart rate variability. Of those participants who were asked to collect saliva samples, 90.1% (n=4609) returned samples. Some samples were not assayed for technical reasons. Participants taking corticosteroid

medication were excluded from analysis (n=236). Any participants taking the first sample more than 10 minutes after waking were excluded from analysis (n=634), this is the commonly used cut off when investigating daytime cortisol levels as the cortisol awakening response is already substantially under way.

A missing value on the work stress measure could indicate that the data were not available at a particular phase, the participant dropped out, or the participant was not in employment. There were 7721 Participants who were still in employment at phase 2 with work stress data at both phases 1 and 2. Out of these participants, 98% had follow up data on incident CHD, 86-90% had information on health behaviours and the metabolic syndrome at phase 3, 45-49% had information on heart rate variability and cortisol at phase 7.

Cox proportional hazard regression models were used to model the association between the cumulative work stress measures (from phases 1 and 2) and incident CHD events (from phases 2 to 7), adjusted for age, sex and employment grade, smoking history, total cholesterol and hypertension (systolic blood pressure >140 and diastolic blood pressure >90, or on antihypertensive medication). Logistic/linear regression models were then used to model the association between cumulative work stress and binary/continuous CHD risk factors. Finally, Cox proportional hazard regression models were used again to examine the reduction in the hazard ratios of cumulative work stress on CHD, adjusted for potential intermediate pathways (health behaviours and the metabolic syndrome). Heart rate variability and cortisol could not be examined as potential mediators as they were not measured in the first few phases of data collection. All statistical significance testing used a two-sided test at the 0.05 significance level. As the main exposure (work stress) consisted of two pairwise comparisons (no report vs. one report, and no report vs. two reports),

Bonferroni corrected p values (a conservative statistical adjustment to adjust for multiple comparisons) are reported to reduce the risk of Type 1 errors. Some of the analyses were stratified by age-group if there was a significant interaction between age and work stress.

## **Results**

The distribution of all the variables in the analysis is shown in Appendix Table A1.

Table 1 displays the hazard ratios of incident CHD by cumulative measures of work stress from phases 1 and 2. Greater reports of work stress were associated with a higher risk of CHD. This was true for both major CHD events (fatal events and myocardial infarction) and definite angina. Although reporting bias may lead to a spurious association between self-reports of stress and angina pectoris,(18) the estimated risks of myocardial infarction and definite angina were similar and so further analyses combined these two CHD outcomes.

There was a significant interaction between age and two reports of work stress ( $p=0.04$ ), so the analysis is stratified by age group. Among younger participants (aged 37 to 49 at phase 2), there was a clear dose response association between greater reports of work stress and higher risks of incident CHD events. Among older participants (aged 50 to 60), there was little association between work stress and CHD. Stratifying by employment status at phase 5 revealed similar effects (analysis not shown).

Table 2 shows the association of work stress (measured at phases 1 and 2) with the metabolic syndrome, its components and health behaviours (all from phase 3) among younger (aged under 50) respondents in the Whitehall II cohort. Greater reports of work stress were associated with poorer health behaviours in terms of

eating less fruit and vegetables, and less physical activity. In addition, work stress was associated with not drinking any alcohol (which increased the risk of CHD- see appendix table A2). Work stress was also associated with the overall metabolic syndrome and four of its five components. Adjusting for health behaviours only slightly reduced the association between work stress and the overall metabolic syndrome.

Table 3 shows the association between work stress (at phases 1 and 2) and low heart rate variability (at phase 7), and morning rise in cortisol (at phase 7) for participants at all ages (there was no significant interaction between age and work stress). Greater reports of work stress were associated with lower heart rate variability in terms of lowering of the total variance and low and high frequency components. There was little association with morning rise in cortisol. However, additional cross sectional analysis at phase 7 between work stress and cortisol revealed significantly elevated morning rise in cortisol among those reporting work stress ( $p < 0.05$ ). All the analyses in Table 3 were adjusted for age, sex, employment grade, hypertension, total cholesterol, smoking and other health behaviours.

Table 4 displays the hazard ratios of incident CHD for the younger respondents (aged under 50) by work adjusted for behavioural risk factors and the metabolic syndrome. There was a 16% reduction in the hazard ratios when behavioural risk factors were adjusted for, and a similar reduction when adjusting for the overall metabolic syndrome. Adjusting for both health behaviours and the metabolic syndrome reduced the work stress-CHD association by around 32%.

## **Discussion**

Cumulative work stress is a risk factor for CHD and neuroendocrine stress responses, especially among the younger, working age population. Around 32% of the effect of work stress on CHD can be explained by the effect of work stress on health behaviours (low physical activity and poor diet in particular) and the metabolic syndrome.

The association between work stress and CHD was stronger among employees younger than 50 and those still in employment. This is in agreement with previous age group analyses of work stress(19) and is consistent with the fact that more robust work stress-CHD associations have been found in studies employing younger(20),(21) than older cohorts.(22),(23) Among older employees, the impact of work stress might be attenuated due to a healthy worker survivor bias. Retirement during the follow-up removes work stress and this exposure misclassification may also reduce the effect of work stress. Furthermore, an increasing number of other age-related causes of CVD may eclipse the effect of work stress as these other causes figure into both the numerator and the denominator of the ratio.

An important case-control study (INTERHEART) of 11,119 patients with a first myocardial infarction (MI) and 13,648 age and sex-matched controls in 52 countries found that 'permanent' stress at work was associated with over twice the odds of MI compared to those reporting no stress at work (24). However few studies have been able to move from demonstrating associations to causality. This paper builds on the INTERHEART and other studies by advancing a causal understanding of this association in terms of dose-response associations, establishing the plausibility of this association in terms of underlying biological and behavioural mechanisms, and demonstrating the specificity of this association among working age populations.

There are relatively few studies which have found associations between work stress and (un)healthy behaviours. Work stress is associated with smoking and exercise,(25) while fatty food intake increases under stressful conditions.(26) Work stress has also been linked with problem drinking, although in this cohort, non-drinkers had the highest risk of CHD (and were more likely to report work stress).

Previous cross-sectional analysis from the Whitehall II study has shown low control at work is associated with poor autonomic function,(2) and neuroendocrine activation during the working day.(4) Longitudinal analyses from the study has shown that work stress is related to CHD(14), the metabolic syndrome(7), and predicts weight gain and incident obesity.(8) This study adds to the literature by showing a linear association between work stress and CHD events, the components of the metabolic syndrome, and lower heart variability. In addition, around 16% of the effect of work stress on CHD can be explained by the effect of work stress on the metabolic syndrome. As there was little reduction in the association between work stress and the metabolic syndrome after adjusting for health behaviours, work stress may directly affect neuroendocrine stress mechanisms independently of health behaviours, resulting in increased risks of the metabolic syndrome. Direct biological stress-effects are additionally possible through acute work-related stressors triggering myocardial infarction in susceptible individuals,(27) a possibility which is consistent with the relatively small effect attenuation after adjustment for metabolic components and the fact that the association between work stress and CHD diluted in individuals who stopped work during follow-up. Heart rate variability and cortisol were not measured in the early phases of the study so their role as a potential mediator of the work stress- CHD association could not be examined. However, adjusting for health behaviours did not change the association between work stress

and (low) heart rate variability, suggesting a direct effect on the autonomic nervous system and neuroendocrine function, rather than indirect effects through health behaviours. The association between work stress and the heart rate variability components suggests that work stress leads to vagal withdrawal and sympathetic saturation indicating a prevalence of sympathetic mechanisms leading to cardiac electrical instability. (28)

Cumulative work stress did not predict a greater cortisol awakening response. However, there was a cross sectional association between work stress and greater cortisol awakening response. A lag period of around 12 years between exposure (work stress) and disturbances in the circadian rhythm of cortisol may not be optimal for detection of the hypothesised neuroendocrine effect.

The Whitehall II cohort is a sample of primarily office-based white-collar workers. There were few manual workers in the cohort. It is possible that the mechanisms underlying the association of work stress with CHD may differ in manual workers although there is little evidence for this hypothesis.(29) Previous research has suggested that the effect of work stress on cardiovascular is less consistent among women.(30) The Whitehall II cohort is predominantly male (67%), although gender stratified analysis revealed similar estimates of work stress on CHD among younger men and women. Missing data is a common problem all cohort studies face. Non-responders at the later clinical examinations were more likely to report work stress, consume less alcohol, have poor diets and high cholesterol, come from lower employment grades, be smokers, physically inactive and obese, resulting in an underestimation of these effects in the analyses. The results on the heart rate variability and cortisol are less robust compared to the other outcomes due to the greater non-response at phase 7. The metabolic syndrome has been criticized as a

purely artificial construct,(31) not contributing any further information over its component risk factors although recent results suggest otherwise.(32) This paper acknowledges this debate on the metabolic syndrome, and presents results on the syndrome itself as well as its components. There may be unmeasured confounders which may “cause” the association between work stress and CHD, such as other sources of stress and personality type.

This study adds to the evidence that the work stress-CHD association is causal in nature.(10) We demonstrate, within a population of office staff largely unexposed to physical occupational hazards, a prospective dose-response relation between psychosocial stress at work and CHD over 12 years of follow-up. We confirm, during the same exposure period, the plausibility of the proposed pathways involving behavioural mechanisms, neuroendocrine and autonomic activation, and development of risk factor clustering, represented by the metabolic syndrome.(1) (6) (7) (2) Further, those who are older (and are more likely to be retired and less exposed to work stress) are less susceptible to the work psychosocial effect, presenting a coherent pattern in our findings. This study demonstrates that stress at work can lead to coronary heart disease through direct activation of neuroendocrine stress pathways and indirectly through health behaviours.

## **Acknowledgments**

We thank all participating civil service departments and their welfare, personnel, and establishment officers; the Occupational Health and Safety Agency; the Council of Civil Service Unions; all participating civil servants in the Whitehall II study; and all members of the Whitehall II study team.

## Reference List

1. Brunner EJ, Marmot MG. Social organisation, stress and health. In: Marmot MG, Wilkinson RG, eds. *Social determinants of health*. Oxford: Oxford University Press; 1999.
2. Hemingway H, Shipley M, Brunner E, Britton A, Malik M, Marmot M. Does autonomic function link social position to coronary risk? The Whitehall II study. *Circulation* 2005 Jun 14;111(23):3071-3077.
3. Vrijkotte TGM, van Doornen LJP, de Geus EJC. Effects of work stress on ambulatory blood pressure, heart rate, and heart rate variability. *Hypertension* 2000 Apr;35(4):880-886.
4. Kunz-Ebrecht SR, Kirschbaum C, Steptoe A. Work stress, socioeconomic status and neuroendocrine activation over the working day. *Soc Sci Med* 2004 Apr;58(8):1523-1530.
5. Bjorntorp P, Rosmond R. The metabolic syndrome - a neuroendocrine disorder? *Br J Nutr* 2000;83(Suppl. 1):S49-S57.
6. Brunner EJ, Hemingway H, Walker BR, Page M, Clarke P, Juneja M, Shipley MJ, Kumari M, Andrew R, Seckl JR, Papadopoulos A, Checkley S, Rumley A, Lowe GDO, Stansfeld SA, Marmot MG. Adrenocortical, autonomic, and inflammatory causes of the metabolic syndrome: nested case-control study. *Circulation* 2002 Nov 19;106(21):2659-2665.
7. Chandola T, Brunner E, Marmot M. Chronic stress at work and the metabolic syndrome: prospective study. *British Medical Journal* 2006 Mar 4;332(7540):521-524A.
8. Brunner EJ, Chandola T, Marmot MG. Prospective effect of job strain on general and central obesity in the Whitehall II Study. *Am J Epidemiol* 2007 Apr 1;165(7):828-837.
9. Ohlin B, Nilsson PM, Nilsson JA, Berglund G. Chronic psychosocial stress predicts long-term cardiovascular morbidity and mortality in middle-aged men. *Eur Heart J* 2004 May;25(10):867-873.
10. Hill AB. The environment and disease: association or causation? *Proc Royal Soc Med* 1965;58:295-300.
11. Marmot M, Brunner E. Cohort profile: The Whitehall II study. *International Journal of Epidemiology* 2005 Apr;34(2):251-256.

12. Karasek R, Theorell T. *Healthy work: stress, productivity, and the reconstruction of working life*. New York: Basic Books; 1990.
13. Landsbergis PA, Schnall PL, Warren K, Pickering TG, Schwartz JE. Association between ambulatory blood pressure and alternative formulations of job strain. *Scand J Work Environ Health* 1994;20:349-363.
14. Kuper H, Marmot M. Job strain, job demands, decision latitude, and the risk of coronary heart disease within the Whitehall II study. *J Epidemiol Community Health* 2003;57(2):147-153.
15. Tunstall-Pedoe H, Kuulasmaa K, Amouyel P, Arveiler D, Rajakangas AM, Pajak A. Myocardial-Infarction and Coronary Deaths in the World-Health-Organization Monica Project - Registration Procedures, Event Rates, and Case-Fatality Rates in 38 Populations from 21 Countries in 4 Continents. *Circulation* 1994 Jul;90(1):583-612.
16. Expert Panel on Detection. Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, And Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel III). *JAMA* 2001;285:2486-2497.
17. Badrick E, Kirschbaum C, Kumari M. The relationship between smoking status and cortisol secretion. *J Clin Endocrinol Metab* 2007 Mar;92(3):819-824.
18. Nielsen NR, Kristensen TS, Prescott E, Larsen KS, Schnohr P, Gronbaek M. Perceived stress and risk of ischemic heart disease: causation or bias? *Epidemiology* 2006 Jul;17(4):391-397.
19. Theorell T, Tsutsumi A, Hallquist J, Reuterwall C, Hogstedt C, Fredlund P, Emlund N, Johnson JV. Decision latitude, job strain, and myocardial infarction: a study of working men in Stockholm. The SHEEP Study Group. Stockholm Heart epidemiology Program. *Am J Public Health* 1998 Mar;88(3):382-388.
20. Alterman T, Shekelle RB, Vernon SW, Burau KD. Decision latitude, psychologic demand, job strain, and coronary heart disease in the Western Electric Study. *Am J Epidemiol* 1994 Mar 15;139(6):620-627.
21. Kivimaki M, Leino-Arjas P, Luukkonen R, Riihimaki H, Vahtera J, Kirjonen J. Work stress and risk of cardiovascular mortality: prospective cohort study of industrial employees. *BMJ* 2002 Oct 19;325(7369):857.
22. Lee S, Colditz G, Berkman L, Kawachi I. A prospective study of job strain and coronary heart disease in US women. *Int J Epidemiol* 2002 Dec;31(6):1147-1153.
23. Eaker ED, Sullivan LM, Kelly-Hayes M, D'Agostino RB, Sr., Benjamin EJ. Does job strain increase the risk for coronary heart disease or death in men and women? The Framingham Offspring Study. *Am J Epidemiol* 2004 May 15;159(10):950-958.

24. Rosengren A, Hawken S, Ounpuu S, Sliwa K, Zubaid M, Almahmeed WA, Blackett KN, Sitthi-amorn C, Sato H, Yusuf S, INTERHEART investigators. Association of psychosocial risk factors with risk of acute myocardial infarction in 11119 cases and 13648 controls from 52 countries (the INTERHEART study): case-control study. *Lancet* 2004;364:953-962.
25. Hellerstedt WL, Jeffery RW. The association of job strain and health behaviours in men and women. *Int J Epidemiol* 1997 Jun;26(3):575-583.
26. Oliver G, Wardle J, Gibson EL. Stress and food choice: a laboratory study. *Psychosom Med* 2000 Nov;62(6):853-865.
27. Moller J, Theorell T, de FU, Ahlbom A, Hallqvist J. Work related stressful life events and the risk of myocardial infarction. Case-control and case-crossover analyses within the Stockholm heart epidemiology programme (SHEEP). *J Epidemiol Community Health* 2005 Jan;59(1):23-30.
28. Malik M, Camm AJ. Components of heart rate variability--what they really mean and what we really measure. *Am J Cardiol* 1993 Oct 1;72(11):821-822.
29. Brunner EJ, Kivimaki M, Siegrist J, Theorell T, Luukkonen R, Riihimaki H, Vahtera J, Kirjonen J, Leino-Arjas P. Is the effect of work stress on cardiovascular mortality confounded by socioeconomic factors in the Valmet study? *J Epidemiol Community Health* 2004 Dec;58(12):1019-1020.
30. Everson-Rose SA, Lewis TT. Psychosocial factors and cardiovascular diseases. *Annual Review of Public Health* 2005;26:469-500.
31. Yudkin JS. Insulin resistance and the metabolic syndrome-or the pitfalls of epidemiology. *Diabetologia* 2007 Aug;50(8):1576-1586.
32. Sundstrom J, Riserus U, Byberg L, Zethelius B, Lithell H, Lind L. Clinical value of the metabolic syndrome for long term prediction of total and cardiovascular mortality: prospective, population based cohort study. *BMJ* 2006 Apr 15;332(7546):878-882.

Table 1 Hazard Ratios (95% confidence intervals) of incident CHD events (phases 2 to 7) by cumulative work stress (phases 1 to 2), age group: the Whitehall II study with an average follow up of 12 years

Case definition and sample	Work Stress			Linear trend p value
	No report	1 report	2 reports	
All CHD- All ages	1.00	1.23 (0.90, 1.68)	1.33 (1.04, 1.69)	0.01
p value <sup>1</sup>		0.19	0.02	
p value <sup>2</sup>		0.37	0.04	
Cases/N	416/6052	38/497	68/779	
CHD Death or Myocardial Infarction- All ages	1.00	1.18 (0.75, 1.87)	1.56 (1.12, 2.17)	0.01
p value <sup>1</sup>		0.47	0.01	
p value <sup>2</sup>		0.94	0.02	
Cases/N	242/6285	24/522	43/818	
Definite Angina- All ages	1.00	1.34 (0.93, 1.93)	1.43 (1.07, 1.90)	0.01
p value <sup>1</sup>		0.11	0.02	
p value <sup>2</sup>		0.23	0.03	
Cases/N	337/6276	35/523	57/819	
All CHD- Age 37-49 at baseline	1.00	1.40 (0.88, 2.22)	1.68 (1.17, 2.42)	<0.01
p value <sup>1</sup>		0.16	<0.01	
p value <sup>2</sup>		0.32	0.01	
Cases/N	174/3912	22/346	38/509	
All CHD- Age 50-60 at baseline	1.00	1.09 (0.68, 1.77)	1.13 (0.79, 1.63)	0.47
p value <sup>1</sup>		0.71	0.51	
p value <sup>2</sup>		1.00	1.00	
Cases/N	258/2314	19/170	33/300	

5 Hazard Ratios are adjusted for age, sex, employment grade, hypertension, total cholesterol and smoking history.

<sup>1</sup> p value adjusted for age, sex, employment grade, hypertension, total cholesterol and smoking.

<sup>2</sup> Bonferroni corrected p value adjusted for age, sex, employment grade, hypertension, total cholesterol and smoking.

Table 2: Odds Ratios (95% CI) of health behaviours (phase 3) and metabolic syndrome (phase 3), by cumulative work stress (phases 1 to 2): Whitehall II respondents aged under 50 at phase 2

<b>Health Behaviours</b>			
	Model 1		<b>Cases/N</b>
Less than monthly Fruit/Veg			
no report of Work Stress	1.00		42/3575
1 report	1.10 (0.43, 2.84)		5/316
2 reports	2.12 (1.07, 4.18)		11/461
No Alcohol consumption			
no report of Work Stress	1.00		558/3581
1 report	1.24 (0.92, 1.67)		66/316
2 reports	1.42 (1.11, 1.82)		101/461
No Physical Activity			
no report of Work Stress	1.00		377/3581
1 report	1.07 (0.74, 1.55)		37/316
2 reports	1.33 (1.00, 1.78)		66/460
Current Smoker			
no report of Work Stress	1.00		464/3580
1 report	1.27 (0.93, 1.73)		56/316
2 reports	1.11 (0.84, 1.47)		68/460
<b>Metabolic Syndrome</b>			
	Model 1	Model 2	<b>Cases/N</b>
High Waist			
no report of Work Stress	1.00	1.00	231/3292
1 report	1.29 (0.84, 1.99)	1.24 (0.81, 1.92)	26/283
2 reports	1.51 (1.08, 2.13)	1.46 (1.03, 2.06)	45/426
High Fasting Glucose			
no report of Work Stress	1.00	1.00	570/3201
1 report	1.02 (0.74, 1.42)	1.05 (0.76, 1.47)	48/269
2 reports	1.40 (1.08, 1.80)	1.43 (1.10, 1.85)	89/410
High Triglycerides			
no report of Work Stress	1.00	1.00	802/3308
1 report	1.18 (0.89, 1.57)	1.16 (0.87, 1.54)	78/280
2 reports	1.33 (1.06, 1.69)	1.30 (1.03, 1.65)	119/425
HDL cholesterol			
no report of Work Stress	1.00	1.00	597/3308
1 report	1.21 (0.89, 1.63)	1.17 (0.86, 1.59)	61/280
2 reports	1.32 (1.03, 1.68)	1.26 (0.98, 1.62)	95/425
Hypertension			
no report of Work Stress	1.00	1.00	1182/3332
1 report	0.87 (0.67, 1.13)	0.88 (0.67, 1.14)	93/285
2 reports	1.13 (0.91, 1.39)	1.13 (0.91, 1.40)	159/430
ATPIII Metabolic Syndrome			
no report of Work Stress	1.00	1.00	357/3308
1 report	1.33 (0.93, 1.91)	1.33 (0.93, 1.91)	39/280
2 reports	1.72 (1.30, 2.29)	1.69 (1.26, 2.25)	69/425

5 Logistic Regression Odds Ratios in Model 1 are adjusted for age, sex and employment grade. Logistic Regression Odds Ratios in Model 2 additionally adjust for health behaviours.

Table 3: Regression coefficients (95% CI) of heart rate variability (Phase 7) and morning rise in cortisol (Phase 7), by cumulative work stress (phases 1 to 2): Whitehall II respondents, all ages

	All Ages	N
<b>Log of Low Frequency Power</b>		
no report of Work Stress	0.00	2769
1 report	-0.09 (-0.23, 0.04)	211
2 reports	-0.14 (-0.25, -0.02)	310
<i>p for linear trend</i>	<0.01	
<b>Log of High Frequency Power</b>		
no report of Work Stress	0.00	2769
1 report	-0.05 (-0.21, 0.11)	211
2 reports	-0.14 (-0.27, 0.00)	310
<i>p for linear trend</i>	<0.05	
<b>Log of SD of NN intervals</b>		
no report of Work Stress	0.00	2769
1 report	-0.05 (-0.12, 0.01)	211
2 reports	-0.05 (-0.10, 0.00)	310
<i>p for linear trend</i>	<0.05	
<b>Morning Rise in cortisol</b>		
no report of Work Stress	0.00	2368
1 report	0.00 (-1.85, 1.85)	169
2 reports	-0.60 (-2.11, 0.91)	274
<i>p for linear trend</i>	0.45	

All models are adjusted for age, sex, employment grade (phase 1), total cholesterol (phase 1), hypertension (phase 1), smoking history (phase 1) and other health behaviours (phase 3). In addition, morning rise in cortisol is adjusted for waking up time.

5

Table 4 Hazard Ratios of incident all CHD events (phases 3 to 7) by cumulative work stress (phases 1 to 2) adjusted for health behaviours (phase 3) and metabolic syndrome (phase 3): Whitehall II respondents aged under 50 at phase 2

<b>Work stress</b>	Model 1	+ All Health Behaviours	
No reports	1.00	1.00	140/3408
1 report	1.52 (0.93, 2.48)	1.43 (0.87, 2.34)	18/292
2 reports	1.56 (1.02, 2.37)	1.47 (0.97, 2.25)	26/434
p for linear trend	0.02	0.04	
	Model 1	+ Metabolic Syndrome	
No reports	1.00	1.00	144/3419
1 report	1.48 (0.90, 2.41)	1.44 (0.88, 2.36)	18/294
2 reports	1.61 (1.06, 2.43)	1.51 (1.00, 2.29)	27/439
p for linear trend	0.01	0.03	
	Model 1	+ Health Behaviours&Metabolic Syndrome	
No reports	1.00	1.00	136/3265
1 report	1.41 (0.84, 2.37)	1.27 (0.75, 2.15)	16/275
2 reports	1.56 (1.02, 2.39)	1.38 (0.90, 2.13)	25/416
p for linear trend	0.03	0.11	

5

Model 1 is adjusted for age, sex and employment grade..

**Appendix Table A1: Distribution of the variables in the analysis**

			<b>Employment Grade (phase 1)</b>	
	<b>Sex</b>		High	3028
Men	3413		Middle	4943
Women	6895		Low	2337
	<b>Agegroup (phase 1)</b>		<b>Total Cholesterol (phase 1)</b>	
35-39	2811		<5.2 mmol/L	2510
40-44	2663		5-2-6.2 mmol/L	4006
45-49	2107		>6.2 mmol/L	3718
50-56	2727		Missing	74
	<b>Cigarette Smoking (phase 1)</b>		<b>Hypertension (phase 1)</b>	
Never smoker	5062		Normotensive	9461
Ex smoker	3274		Sys BP>140/Dias BP>90*	832
0-9 cigs/day	540		Missing	15
10-19 cigs/day	774			
20 or more cigs/day	418			
Missing	240			
	<b>Moderate Exercise (phase 3)</b>		<b>Iso-strain (phases 1 to 2)</b>	
3 times/week or more	1284		No report	6363
1-2 times/week	3692		1 report	529
1-3 times/month	2290		2 reports	829
Never/hardly	1042		Missing	2587
Missing	2000			
	<b>Current smoker (phase 3)</b>		<b>Alcohol consumption (Phase 3)</b>	
Non smoker	7168		Low	1625
Smoker	1145		Moderate	5399
Missing	1995		High	1288
			Missing	1996
	<b>Fruit/Veg consumption (phase 3)</b>		<b>High Triglycerides (phase 3)</b>	
Less than daily	8198		Normal	5770
Daily or more	112		≥150 mg/dL	2252
Missing	1998		Missing	2286
	<b>High Waist (phase 3)</b>		<b>Low HDL (phase 3)</b>	
Normal	7258		Normal	6477
M>102cm or F>88cm	737		M<40mg/dL, F<50mg/dL	1542
Missing	2313		Missing	2289
	<b>High Glucose (phase 3)</b>		<b>Metabolic Syndrome (phase 3)</b>	
Normal	6006		No syndrome	6897
≥110 mg/dL	1603		Metabolic Syndrome	1125
Missing	2699		Missing	2286
	<b>High Blood Pressure (phase 3)</b>		<b>Heart Rate Variability</b>	N=4095
Normal	4823		(phase 7)	
High BP*	3351		<b>Morning rise in cortisol</b>	N=3490
Missing	2134		(phase 7)	

\* includes those on antihypertensive medications

Appendix Table A2:Hazard Ratios of incident all CHD events (phases 3 to 7): Whitehall II respondents aged under 50 at phase 2.

<i>Employment Grade</i>	
High	1.00
Middle	1.14 (0.84, 1.56)
Low	1.65 (1.04, 2.60)
<i>Work Stress</i>	
No reports of work stress	1.00
1 report	1.55 (0.97, 2.46)
2 reports	1.62 (1.10, 2.40)
<i>Waist circumference</i>	
Normal	1.00
High Waist	2.04 (1.35, 3.09)
<i>Triglycerides</i>	
Normal	1.00
High Triglycerides	1.93 (1.44, 2.59)
<i>Glucose Tolerance</i>	
Normal	1.00
Glucose Intolerance	1.35 (0.96, 1.89)
<i>HDL cholesterol</i>	
Normal	1.00
Low HDL	2.03 (1.50, 2.74)
<i>Blood Pressure</i>	
Normal	1.00
High Blood Pressure/Antihypertensive medication	2.16 (1.63, 2.87)
<i>Overall Metabolic Syndrome</i>	
No syndrome	1.00
3 or more MS components	2.52 (1.82, 3.49))
<i>Reported fruit/veg consumption</i>	
Daily or more	1.00
Less than daily	2.38 (1.12, 5.06)
<i>Physical Activity</i>	
3/week or more	1.00
1-2/week	1.51 (0.93, 2.46)
1/3month	1.91 (1.15, 3.16)
Never	2.16 (1.20, 3.90)
<i>Alcohol consumption in the last week</i>	
Non drinker	1.00
Safe alcohol limits	0.62 (0.43, 0.88)
Unsafe alcohol limits	0.71 (0.46, 1.11)
<i>Cigarette smoker</i>	
Non smoker	1.00
Ex smoker	1.04 (0.75, 1.44)
1-9 cigs/day	2.15 (1.24, 3.72)
10-19 cigs/day	1.39 (0.74, 2.60)
20+ cigs/day	3.06 (1.71, 5.49)

Hazard Ratios are adjusted for age and sex.

## **Work Stress and Coronary Heart Disease- What are the Mechanisms?**

Permissions Information: No illustrations or figures were duplicated from previously published work.

**Work Stress and Coronary Heart Disease- What are the Mechanisms?**

Word count: 3910