



A tale of three cities: persisting high HIV prevalence, risk behaviour and undiagnosed infection in community samples of men who have sex with men

Julie P Dodds, Anne M Johnson, John V Parry and Danielle E Mercey

Sex. Transm. Inf. 2007;83:392-396; originally published online 1 May 2007;
doi:10.1136/sti.2006.021782

Updated information and services can be found at:
<http://sti.bmj.com/cgi/content/full/83/5/392>

These include:

References

This article cites 17 articles, 8 of which can be accessed free at:
<http://sti.bmj.com/cgi/content/full/83/5/392#BIBL>

Rapid responses

You can respond to this article at:
<http://sti.bmj.com/cgi/eletter-submit/83/5/392>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Correction

A correction has been published for this article. The contents of the [correction](#) have been appended to the original article in this reprint. The correction is also available online at:
<http://sti.bmj.com/cgi/content/full/83/6/500>

Notes

To order reprints of this article go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to *Sexually Transmitted Infections* go to:
<http://journals.bmj.com/subscriptions/>

A tale of three cities: persisting high HIV prevalence, risk behaviour and undiagnosed infection in community samples of men who have sex with men

Julie P Dodds, Anne M Johnson, John V Parry, Danielle E Mercey

Sex Transm Infect 2007;**83**:392–396. doi: 10.1136/sti.2006.021782

See end of article for authors' affiliations

Correspondence to: Ms J Dodds, Department of Primary Care and Population Science, Centre for Sexual Health & HIV Research, University College London, Mortimer Market Centre, Mortimer Market, off Capper Street, London WC1E 6JB, UK; jdodds@gum.ucl.ac.uk

Accepted 16 February 2007

Objectives: To examine the geographical variations in HIV prevalence (diagnosed and undiagnosed), use of sexual health services, sexually transmitted infections and sexual behaviour in a community sample of men who have sex with men in three cities in England, specifically London, Brighton and Manchester.

Methods: Cross-sectional surveys of men visiting gay community venues in three large cities in England. Men self-completed a questionnaire and provided an anonymous oral fluid sample for HIV antibody testing.

Results: HIV prevalence ranged from 8.6% to 13.7% in the three cities. Over one-third of HIV infection remained undiagnosed in all sites despite 69% of HIV-positive men reporting attending a genitourinary medicine clinic in the last year. Similar and high levels of risk behaviour were reported in all three cities. 18% of HIV-negative men and 37% of HIV-positive men reported unprotected anal intercourse with more than one partner in the last year. 20% of negative men and 41% of positive men reported an STI in the last year.

Conclusions: Across all cities, despite widespread availability of anti-retroviral treatment and national policy to promote HIV testing, many HIV infections remain undiagnosed. Data from this community sample demonstrate high levels of risk behaviour and STI incidence, especially among those who are HIV positive. Renewed efforts are needed to increase diagnosis and to reduce risk behaviour to stem the continuing transmission of HIV.

There is evidence of continuing transmission of HIV among men who have sex with men (MSM) in the UK, particularly in London, despite safer sex campaigns. MSM still form the largest group of HIV-infected people in the UK.¹ There has been an increase over time in the number of MSM newly diagnosed as having HIV infection.² Continuing detection of HIV infection among young MSM in the unlinked anonymous seroprevalence survey also suggests recent transmission.^{1–3}

There have been significant changes in the gay scene in Britain over the last two decades, with increased numbers of commercial venues outside of London, primarily in Brighton and Manchester. These cities have thriving gay scenes, which act as a major pull for those living in the surrounding areas. Increasing levels of sexually transmitted infections (STIs) have been reported in all three cities, most notably between 1999 and 2000, with a significant increase in new syphilis cases among MSM.^{4–5} Rates of gonorrhoea in the UK, both inside and outside London, also increased among MSM.^{2–8} STI outbreaks were found to be associated with high rates of partner change, casual sex contacts and poor contact with sexual health services.^{7–9} We have previously reported on increasing levels of sexual risk behaviour in London through our community-based programme of behavioural and HIV prevalence surveillance in London.^{10–11} More recently, we have extended this research to other cities in response to concerns about increasing incidences of STIs.

In this paper, we report on the geographical variations between three cities commonly viewed to have the largest gay communities in the UK, specifically London, Brighton and Manchester, in the prevalence of diagnosed and undiagnosed HIV infection, genitourinary medicine (GUM) clinic attendance, STIs and unprotected anal intercourse (UAI) in the last year.

METHODS

Details of the method have been reported elsewhere.^{11–12} We undertook unlinked anonymous surveys in Manchester, London and Brighton among a representative sample of MSM attending bars, clubs and saunas. The surveys were conducted in Manchester between June and August 2003, in London from November 2003 to January 2004 and in Brighton during February 2004. All men in the venue, or in a particular area of the venue in the larger sites, present during a defined time period (eg, 2 h mid-evening) were offered, by trained fieldworkers, a short self-completion questionnaire and asked to provide an oral fluid sample to be tested for anti-HIV antibodies. An OraSure device (OraSure Technologies, Bethlehem, Pennsylvania, USA; OraSure oral specimen collection pad, Epitope, Beaverton, Oregon, USA) was used to collect the oral fluid sample, linked to the respondent's questionnaire via a barcode system; no personal identifiers were recorded on either the questionnaire or OraSure. The numbers of men offered an OraSure and questionnaire were recorded to calculate a response rate.

The questionnaire collected data on demographics, sexual health service use, HIV testing, HIV status (perceived and self-reported test results), diagnosed STIs and sexual behaviour.

Laboratory testing

All oral fluid samples collected at each session were sent to the Centre for Infections, London, UK, approximately one batch every 2 weeks. Samples were tested for total IgG as a specimen quality check and then tested for anti-HIV antibody following a predetermined algorithm. All men were screened by an IgG

Abbreviations: GUM, genitourinary medicine; MSM, men who have sex with men; STI, sexually transmitted infection; UAI, unprotected anal intercourse

Table 1 Response rates, demographics and HIV antibody prevalence by city of recruitment*

	City of recruitment			Significance	
	London	Brighton	Manchester	χ^2 (2df)	p Value
Response	63 (1541/2450)	66 (387/596)	70 (383/544)	11.03	<0.01
Median (range) age, years	33 (17–78)	34 (17–73)	27 (17–77)	82.93	<0.001†
White ethnicity	85 (1222/1436)	95 (356/373)	96 (335/348)		
Unadj OR (95% CI)	1	3.67 (2.22 to 6.09)	4.51 (2.55 to 8.00)	48.25	<0.001
Educated after 16 years	88 (1265/1436)	77 (288/373)	83 (286/346)		
Unadj OR (95% CI)	1	0.46 (0.34 to 0.61)	0.64 (0.46 to 0.89)	29.53	<0.001
Employed	89 (1278/1434)	79 (219/368)	87 (304/348)		
Unadj OR (95% CI)	1	0.46 (0.34 to 0.62)	0.84 (0.59 to 1.21)	25.48	<0.001
HIV antibody positive	12.3 (177/1436)	13.7 (51/373)	8.6 (30/348)		
Unadj OR (95% CI)	1	1.13 (0.81 to 1.57)	0.67 (0.45 to 1.01)	4.85	0.09
Adj OR (95% CI)	1	0.93 (0.65 to 1.33)	0.69 (0.45 to 1.05)	3.01	0.22
HIV-positive men undiagnosed	44.1 (78/177)	33.3 (17/51)	36.7 (11/30)		
Unadj OR (95% CI)	1	0.63 (0.33 to 1.22)	0.73 (0.33 to 1.64)	2.14	0.34
Adj OR (95% CI)	1	1.04 (0.50 to 2.13)	0.72 (0.30 to 1.71)	0.62	0.73

Adj, adjusted; df, degrees of freedom; unadj, unadjusted.

Values are represented as % (n/n) unless otherwise specified.

*Showing OR and adjusted OR for outcomes by city (adjusting for age, education, ethnicity and employment status).

†Using Kruskal–Wallis test.

antibody capture ELISA HIV 1 and the sensitivity and specificity of two men were determined as 99.5% (95% CI 97.1% to 99.9%) and 99.7% (95% CI 98.9% to 99.9%).^{13 14} Reactive specimens underwent further testing by a modified indirect enzyme immunoassay test using oligopeptide antigens (Adaltis HIVDetect, BioStat, Stockport, UK). Reactive specimens whose OD/CO was <4.0 in either assay were tested by western blot (Genelabs HIVblot 2.2), using a procedure modified for application to oral fluid specimens.

Statistical analysis

Data from each city were double entered into Epi-Info and combined into a single dataset in SPSS V.12 software; these data were then transferred and analysed using STATA V.7. Single variable analysis was performed using χ^2 tests for categorical variables and the Kruskal–Wallis test for continuous variables. Multivariate analysis was used to identify differences in HIV prevalence, use of sexual health service, diagnosed STIs and sexual behaviour in London compared with Brighton and Manchester using unconditional logistic regression. Odds ratios (ORs) were used with 95% CIs to compare outcomes in London, Brighton and Manchester. ORs and CIs in this paper are unadjusted or adjusted for demographic factors associated in the univariate analysis, and p values are given for the overall significant differences between the three cities. To assess whether predictor measures for HIV infection act similarly for all three cities, we tested for factors associated with predicting HIV positivity and interactions between cities.

RESULTS

A total of 90 venues (bars, clubs and saunas) were sampled, 59 from London, 19 from Brighton and 12 from Manchester. A total of 3590 questionnaires were offered and 2640 were returned (73% (1788/2450) in London, 72% (427/596) in Brighton and 78% (425/544) in Manchester). Oral fluid samples were collected for 2311 individuals (64% response rate; table 1). Men excluded from further analysis were those who had already completed the questionnaire within the previous 6 months (70 respondents), those who specified their sexual orientation as heterosexual and did not report sex with a man in the last year (57 respondents), those whose oral fluid samples had inadequate levels of IgG (5 specimens) and those whose samples could not be matched to a questionnaire (37 specimens); similar proportions of men from each city were excluded.

Most men in all three cities were white, educated beyond the age of 16 years and currently employed; however, men recruited from Manchester were significantly younger (table 1; $p < 0.001$). Men in Brighton and Manchester were more likely to be white ($p < 0.001$) and less likely to be educated beyond the age of 16 years than the men recruited from London ($p < 0.001$). The men in Brighton were less likely to be currently employed than the men in London and Manchester ($p < 0.001$). Of the men who reported the first half of their postcode, <3% (64/2485) reported a postcode that appeared in more than one sample, suggesting that men were unlikely to have completed the survey in more than one city.

Table 1 also compares the HIV prevalence in the three cities. London and Brighton reported HIV prevalences of 12.3% and 13.7%, respectively; the prevalence was lower in Manchester at 8.6% ($p = 0.22$). The proportion of HIV-positive men whose infection was undiagnosed was high in all three cities, ranging from 33% in Brighton to 44% in London ($p = 0.73$). Of the men who reported attending a GUM clinic in the last year, 20.2% (117/578) were HIV positive in London compared with 23.8% (39/164) in Brighton and 13.2% (21/159) in Manchester ($p = 0.05$). Of the HIV-positive GUM clinic attenders, in London 30.8% (36/117) were undiagnosed compared with 28.2% (11/39) in Brighton and 19.0% (4/21) in Manchester ($p = 0.55$). Of the self-reported non-clinic attenders, 7.1% (60/849) were HIV positive in London compared with 5.4% (11/205) in Brighton and 4.8% (9/189) in Manchester ($p = 0.40$). Of the HIV-positive non-clinic attenders, in London 70% (42/60) were undiagnosed compared with 54.5% (6/11) in Brighton and 77.7% (7/9) in Manchester ($p = 0.49$).

Among all HIV-negative men, over one-third reported attending a GUM clinic in the last year (table 2). This compares with over two-thirds of the HIV-positive men, with no significant difference between cities ($p = 0.82$; table 3).

Of the HIV-negative men, the Manchester sample was significantly less likely to report HIV testing in the last 5 years (61%) than the men in London (68%; adjusted OR 0.73). The HIV-positive men reported levels of testing similar to the HIV-negative men; men in Manchester reported a higher level of testing than in the other two cities at 73%, but the difference between cities was not significant ($p = 0.48$).

Of the HIV-negative men, nearly one in five reported an STI in the last year. Among the HIV-positive men, over one-third of men recruited in London and Manchester reported an STI in the last year compared with half of the men recruited in Brighton, but the difference was not significant ($p = 0.62$).

Table 2 Use of sexual health services, sexually transmitted infections and sexual behaviour of HIV-negative men by city of recruitment*

	City of recruitment			Total	Significance	
	London	Brighton	Manchester		χ^2 (2 df)	p Value
Attended GUM clinic in the last year	37 (461/1250)	39 (125/319)	43 (138/318)	38 (724/1887)		
Unadj OR (95% CI)	1	1.10 (0.86 to 1.42)	1.31 (1.02 to 1.68)		4.65	0.10
Adj OR (95% CI)	1	1.15 (0.88 to 1.49)	1.21 (0.93 to 1.57)		2.64	0.27
Tested for HIV in the last 5 years	68 (847/1252)	63 (202/319)	61 (193/317)	66 (1242/1888)		
Unadj OR (95% CI)	1	0.83 (0.64 to 1.07)	0.74 (0.58 to 0.96)		6.16	0.05
Adj OR (95% CI)	1	0.89 (0.68 to 1.16)	0.73 (0.56 to 0.95)		5.18	0.06
STI in the last year	19 (236/1251)	23 (74/319)	22 (71/318)	20 (381/1888)		
Unadj OR (95% CI)	1	1.30 (0.97 to 1.75)	1.24 (0.92 to 1.67)		4.05	0.13
Adj OR (95% CI)	1	1.31 (0.96 to 1.77)	1.21 (0.88 to 1.64)		3.88	0.17
UAI with >1 partner in the last year	16 (197/1219)	20 (63/311)	23 (70/309)	18 (330/1839)		
Unadj OR (95% CI)	1	1.32 (0.96 to 1.81)	1.52 (1.12 to 2.07)		8.36	0.02
Adj OR (95% CI)	1	1.26 (0.91 to 1.75)	1.30 (0.94 to 1.79)		3.63	0.16
UAI with casual partners in the last year	23 (267/1145)	24 (69/284)	28 (80/288)	24 (416/1717)		
Unadj OR (95% CI)	1	1.06 (0.78 to 1.43)	1.26 (0.95 to 1.69)		2.49	0.29
Adj OR (95% CI)	1	1.05 (0.77 to 1.44)	1.13 (0.83 to 1.53)		0.65	0.72
UAI with partners of an unknown or discordant HIV status in the last year	21 (233/1132)	24 (67/280)	26 (70/274)	22 (370/1686)		
Unadj OR (95% CI)	1	1.21 (0.89 to 1.66)	1.32 (0.97 to 1.80)		3.93	0.14
Adj OR (95% CI)	1	1.21 (0.88 to 1.66)	1.13 (0.83 to 1.56)		1.62	0.44

Adj, adjusted; df, degrees of freedom; GUM, genitourinary medicine; STI, sexually transmitted infection; UAI, unprotected anal intercourse; Unadj, unadjusted. Values are represented as % (n/n) unless otherwise specified.

*OR and adjusted OR for outcomes by city (adjusting for age, education, ethnicity and employment status).

There were no significant differences between the cities in the proportion reporting UAI with more than one partner during the previous 12 months. There were also no significant differences between cities in reporting UAI with casual partners (classified as men with whom the respondent had had sex once only) and in the proportion of men who reported UAI with partners of an unknown or discordant HIV status in the last year

As table 4 shows, HIV-positive men, in all three cities, were more likely to be older and currently unemployed, and less likely to be educated beyond the age of 16 years than HIV-negative men (data on ethnic status were excluded owing to small numbers in Brighton and Manchester). HIV-positive men were more likely to have attended a GUM clinic in the last year and to have reported an STI. The HIV-positive men were also

Table 3 Use of sexual health service use, sexually transmitted infections and sexual behaviour of HIV-positive men by city of recruitment*

	City of recruitment			Total	Significance	
	London	Brighton	Manchester		χ^2 (2 df)	p Value
Attended GUM clinic in the last year	66 (117/177)	78 (39/50)	70 (21/30)	69 (177/257)		
Unadj OR (95% CI)	1	1.82 (0.87 to 3.80)	1.20 (0.52 to 2.77)		2.55	0.28
Adj OR (95% CI)	1	1.12 (0.56 to 2.71)	0.89 (0.37 to 2.16)		0.39	0.82
Tested for HIV in the last 5 years	66 (114/172)	65 (31/48)	73 (22/30)	67 (167/250)		
Unadj OR (95% CI)	1	0.93 (0.47 to 1.81)	1.40 (0.57 to 3.34)		0.70	0.71
Adj OR (95% CI)	1	1.51 (0.72 to 3.16)	1.40 (0.55 to 3.54)		1.46	0.48
STI in the last year	38 (67/176)	51 (26/51)	37 (11/30)	41 (104/257)		
Unadj OR (95% CI)	1	1.69 (0.90 to 3.16)	0.94 (0.42 to 2.10)		2.90	0.23
Adj OR (95% CI)	1	1.29 (0.67 to 2.50)	0.82 (0.36 to 1.89)		0.96	0.62
UAI with >1 partner in the last year	36 (59/165)	43 (21/49)	35 (10/29)	37 (90/243)		
Unadj OR (95% CI)	1	1.35 (0.70 to 2.58)	0.95 (0.41 to 2.17)		0.90	0.64
Adj OR (95% CI)	1	1.27 (0.64 to 2.51)	0.85 (0.36 to 2.00)		0.74	0.69
UAI with casual partners in the last year	39 (59/153)	41 (19/46)	42 (11/26)	40 (89/225)		
Unadj OR (95% CI)	1	1.12 (0.57 to 2.19)	1.16 (0.50 to 2.72)		0.20	0.90
Adj OR (95% CI)	1	0.99 (0.49 to 2.01)	1.01 (0.45 to 2.40)		0.00	0.99
UAI with partners of an unknown or discordant HIV status in the last year	28 (42/151)	34 (16/47)	31 (8/26)	30 (66/224)		
Unadj OR (95% CI)	1	1.34 (0.66 to 2.70)	1.15 (0.47 to 2.85)		0.69	0.71
Adj OR (95% CI)	1	1.44 (0.69 to 3.01)	0.99 (0.39 to 2.53)		0.97	0.62

Adj, adjusted; df, degrees of freedom; GUM, genitourinary medicine; STI, sexually transmitted infection; UAI, unprotected anal intercourse; unadj, unadjusted. Values are represented as % (n/n) unless otherwise specified.

*OR and adjusted OR for outcomes by city (adjusting for age, education, ethnicity and employment status).

more likely to report high-risk sexual behaviour: UAI with more than one partner in the last year; UAI with casual partners; and UAI with partners of an unknown or discordant HIV status. In our analysis of the interactions between cities and the above predictor measures for HIV positivity, all interactions were found not to be significant. For each predictor measure, the association with HIV positivity followed in the same direction for all three cities and these associations were generally of a similar magnitude.

DISCUSSION

This is the first report to explore geographical variations in HIV prevalence, use of sexual health services, high-risk sexual behaviour and accuracy of self-assumed HIV status from community populations of MSM in England. These results, derived from London, Brighton and Manchester, provide the means to measure the effectiveness of current HIV prevention campaigns and to inform future HIV prevention programmes. The surveys provide data that are not readily derived from other surveillance systems. Surveillance data are largely generated from GUM clinic attendances, and this is biased towards those at greater risk of STIs and HIV infection.¹ Brown *et al*¹⁵ found that 51% of MSM tested for syphilis remained unaware of their HIV infection in 2003 compared with 33–44% of men in the community survey.

The reported high levels of STIs, by both HIV-negative and HIV-positive men, which are known to facilitate the transmission of HIV, is a major public health concern.¹⁶ Regular STI screening and effective behavioural interventions are required as part of clinical care to limit further transmission.

Results from the OraSure test demonstrated a high HIV prevalence in all three cities, with a high proportion of HIV infection remaining undiagnosed. Voluntary confidential HIV testing should be promoted in all three cities to reduce the

proportion of men undiagnosed. Barriers to testing, including poor access to GUM clinics,¹⁷ and the stigma associated with HIV must be addressed.¹⁸ The reasons why men are not testing for HIV need to be investigated and the potential for further community testing to access groups who otherwise might not go to a clinic to test should be explored further. Currently, only a small number of rapid HIV and STI testing services in the UK have been developed within the community.^{19–21}

The same method was used and comparable data were collected in each city, allowing for direct comparisons to be made. Sources of bias using the above method and the validity of HIV testing using oral fluid samples have been described elsewhere.^{11 12 22} Of all the men approached, 66.3% agreed to take an oral fluid test and 73.5% agreed to complete a questionnaire, demonstrating acceptability. In all three cities, men who declined the OraSure test reported demographics and behaviour similar to those who accepted the test. An internal consistency check within the questionnaire also demonstrated high levels of reliability and validity of self-reported sexual behaviour with evidence of substantial consistency between questions (data not shown).

Our behavioural data are supported by similar findings from the national Gay Men's Sex Survey conducted by Sigma Research, in which 20.1% in London reported UAI with more than one partner (compared with 18.5% of HIV-negative and HIV-positive men from the OraSure survey reported here), 21.3% in Brighton (compared with 23.3%) and 26.5% in Manchester (compared with 23.7%).²³ However, data from convenience sampling are likely to overestimate the prevalence of sexual risk behaviour and sexual health outcomes of all MSM in Britain.^{24 25}

There is a critical need for continued surveillance in these cities to enable informed public health responses based on current HIV prevalence (diagnosed and undiagnosed), HIV

Table 4 Predictors of HIV positivity*

	City of recruitment			Interaction	
	London	Brighton	Manchester	χ^2 (2 df)	p Value
Median age (years)	36	36	34	2.66	0.27†
Unemployed					
Unadj OR (95% CI)	2.64 (1.76 to 3.96)	4.01 (2.14 to 7.47)	1.84 (0.71 to 4.80)	2.07	0.36
Adj OR (95% CI)	2.50 (1.66 to 3.79)	4.09 (2.12 to 7.87)	1.58 (0.58 to 4.37)	1.43	0.49
Educated after the age of 16 years					
Unadj OR (95% CI)	0.56 (0.37 to 0.86)	0.67 (0.35 to 1.29)	0.45 (0.20 to 1.04)	0.53	0.77
Adj OR (95% CI)	0.58 (0.37 to 0.88)	0.73 (0.37 to 1.44)	0.65 (0.26 to 1.63)	0.45	0.80
Attended GUM clinic in the last year					
Unadj OR (95% CI)	3.34 (2.40 to 4.65)	5.50 (2.72 to 11.15)	3.04 (1.35 to 6.85)	1.75	0.41
Adj OR (95% CI)	3.43 (2.44 to 4.81)	6.81 (3.12 to 14.87)	3.07 (1.33 to 7.06)	3.59	0.16
STI in the last year					
Unadj OR (95% CI)	2.64 (1.89 to 3.68)	3.44 (1.88 to 6.32)	2.01 (0.92 to 4.43)	1.16	0.56
Adj OR (95% CI)	2.64 (1.87 to 3.72)	3.54 (1.87 to 6.72)	2.25 (0.99 to 5.06)	1.39	0.50
UAI with >1 partner in the last year					
Unadj OR (95% CI)	2.89 (2.03 to 4.11)	2.95 (1.57 to 5.54)	1.80 (0.80 to 4.04)	1.18	0.56
Adj OR (95% CI)	2.89 (2.01 to 4.17)	2.71 (1.39 to 5.30)	2.02 (0.88 to 4.67)	0.75	0.69
UAI with casual partners in the last year					
Unadj OR (95% CI)	2.06 (1.45 to 2.94)	2.19 (1.15 to 4.19)	1.91 (0.84 to 4.33)	0.07	0.97
Adj OR (95% CI)	2.04 (1.42 to 2.94)	2.44 (1.22 to 4.85)	1.99 (0.86 to 4.61)	0.63	0.73
UAI with partners of an unknown or discordant HIV status in the last year					
Unadj OR (95% CI)	1.49 (1.01 to 2.18)	1.64 (0.85 to 3.18)	1.30 (0.54 to 3.11)	0.18	0.91
Adj OR (95% CI)	1.58 (1.06 to 2.34)	1.76 (0.87 to 3.58)	1.43 (0.58 to 3.49)	0.39	0.82

Adj, adjusted; df, degrees of freedom; GUM, genitourinary medicine; STI, sexually transmitted infection; UAI, unprotected anal intercourse; unadj, unadjusted.

*Adjusted OR adjusting for age, education, ethnicity and employment status.

†Using Kruskal–Wallis test.

testing and sexual behaviour in the community. These would also facilitate monitoring of the effectiveness of health prevention initiatives over time. The high levels of HIV prevalence and risk behaviour in all three cities indicate a need for more effective HIV prevention.

Our data show remarkable similarities between cities in the proportion of men reporting high-risk sexual behaviour and STIs, among both the HIV-negative and HIV-positive men. The association of predictor factors with HIV infection were often of a similar magnitude, with no significant interactions between cities; therefore, the predictors of HIV infection may be considered broadly similar for all three cities. This suggests that the same HIV risk profile for MSM is applicable to all three cities, which is useful when considering intervention designs for these cities, or the use of randomised controlled trials. However, it should be noted that although this is true for these three cities it might not be true for all UK cities. A comparison of behaviour among community samples of MSM in London and Glasgow (Scotland) showed that HIV testing behaviour and serodiscordant UAI differed by city, supporting city-specific interventions.²⁶

Data from extensive community surveillance can add considerably to the simple disease surveillance data from GUM clinics. These data demonstrate the country-wide high levels of risk behaviour, STIs and HIV prevalence, which together account for the worrying trend of continuing high HIV incidence in MSM.

ACKNOWLEDGEMENTS

We thank the Department of Health and Camden and Islington NHS Primary Care Trusts for the financial support. We thank the local coordination teams, Terrence Higgins Trust South in Brighton, the Lesbian and Gay Foundation in Manchester and Camden and Islington Health Promotion Team in London. We thank the Central Public Health Laboratory, in particular Sharon Barnett. We also thank Dr Andrew Copas for statistical advice. Finally, we thank all the venue managers, staff and respondents for their participation.

Authors' affiliations

Julie P Dodds, Anne M Johnson, Danielle E Mercey, Department of Primary Care and Population Sciences, Centre for Sexual Health & HIV Research, University College London, London, UK

John V Parry, Virus Reference Department, Centre for Infections, Health Protection Agency, London, UK

Funding: This work was supported by the Department of Health and Camden & Islington NHS Primary Care Trusts.

Competing interests: None.

JPD, DEM, JVP and AMJ initiated and designed the study. JPD managed the project. JVP coordinated the HIV testing. JPD analysed the data, wrote the first draft and coordinated subsequent revisions of the paper. All authors contributed to the final version of this paper. JPD and DEM are the guarantors.

REFERENCES

- 1 **The UK Collaborative Group for HIV and STI Surveillance.** *Mapping the issues: HIV and other sexually transmitted infections in the United Kingdom: 2005.* London: Health Protection Agency, Centre for Infections, 2005.
- 2 **The UK Collaborative Group for HIV and STI Surveillance.** *A complex picture. HIV and other sexually transmitted infections in the United Kingdom: 2006.* London: Health Protection Agency, Centre for Infections, 2006.
- 3 **Murphy G, Charlett A, Jordan LF, et al.** HIV incidence appears constant in men who have sex with men despite widespread use of effective antiretroviral therapy. *AIDS* 2004;**18**:265–72.
- 4 **Doherty L, Fenton K, Jones J, et al.** Syphilis: old problem, new strategy. *BMJ* 2002;**325**:153–6.
- 5 **Higgins SP, Sukthankar A, Mahto M, et al.** Syphilis increases in Manchester, UK. *Lancet* 2005;**355**:1466.
- 6 **Martin IMC, Ison CA.** Rise in gonorrhoea in London, UK. *Lancet* 2000;**355**:623–32.
- 7 **Fenton KA, Rogers PA, Simms I, et al.** Increasing gonorrhoea reports—not only in London. *Lancet* 2000;**355**:1907.
- 8 **Macdonald N, Dougan S, McGarrigle CA, et al.** Recent trends in diagnoses of HIV and other sexually transmitted infection in England and Wales among men who have sex with men. *Sex Transm Infect* 2004;**80**:492–7.
- 9 **Clark P, Cook PA, Syed Q, et al.** *Re-emerging syphilis in the North West: lessons from the Manchester outbreak.* Liverpool: John Moores University, 2001.
- 10 **Dodds JP, Nardone A, Mercey DE, et al.** Increase in high risk sexual behaviour among homosexual men, London 1996–98: cross sectional, questionnaire study. *BMJ* 2000;**320**:1510–11.
- 11 **Dodds JP, Mercey DE, Parry JV, et al.** Increasing risk behaviour and high levels of undiagnosed HIV infection in a community sample of homosexual men. *Sex Transm Infect* 2004;**80**:236–40.
- 12 **Nardone A, Dodds JP, Mercey DE, et al.** Active surveillance of sexual behaviour among homosexual men in London. *Commun Dis Public Health* 1998;**1**:197–201.
- 13 **Parry JV, Perry KR, Mortimer PP.** Sensitive assays for viral antibodies in saliva: an alternative to tests on serum. *Lancet* 1987;**2**:72–5.
- 14 **Connell J, Parry JV, Mortimer PP, Duncan J.** Novel assay for the detection of immunoglobulin G anti-human immunodeficiency virus in untreated saliva and urine. *J Med Virol* 1993;**41**:159–64.
- 15 **Brown AE, Tomkins SE, Logan LE, et al.** Monitoring the effectiveness of HIV and STI prevention initiatives in England, Wales, and Northern Ireland: where are we now? *Sex Transm Infect* 2006;**82**:4–10.
- 16 **Fleming DT, Wasserheit JN.** From epidemiological synergy to public health policy and practice: the contribution of other sexually transmitted diseases to sexual transmission of HIV infection. *Sex Transm Infect* 1999;**75**:3–17.
- 17 **Griffiths V, Ahmed-Jushuf I, Cassell JA.** Understanding access to genitourinary medicine services. *Int J STD AIDS* 2004;**15**:587–9.
- 18 **Hickson F, Nutland W, Weatherburn P, et al.** *Making it count—a collaborative planning framework to reduce the incidence of HIV infection during sex between men.* London: Sigma Research, 2003.
- 19 **Terrence Higgins Trust.** Fastest. <http://tht.org.uk/fastest/> (accessed 27 Feb 2007).
- 20 **Terrence Higgins Trust.** *Five steps to better sexual health.* UK: Terrence Higgins Trust, 2005.
- 21 **George House Trust, Lesbian and Gay Foundation, The Manchester Centre for Sexual Health, and The Public Health Development Service.** My HIV test. <http://www.myhivtest.org/> (accessed 27 Feb 2007).
- 22 **Nardone A, Mercey DE, Johnson AM.** Surveillance of sexual behaviour among homosexual men in a central London Health Authority. *Genitourin Med* 1997;**73**:198–202.
- 23 **Reid D, Weatherburn P, Hickson F, et al.** Vital statistics 2003—the Gay Men's Sex Survey (area sub-samples data reports). London: Sigma Research, 2004.
- 24 **Evans AR, Wiggins RD, Mercer CH, et al.** Men who have sex with men in Britain: comparison of a self-selected internet sample with a national probability sample. *Sex Transm Infect.* Published Online First: 13 December 2006. doi:10.1136/sti.2006.23283.
- 25 **Dodds JP, Mercer CH, Mercey DE, et al.** Men who have sex with men: a comparison of a probability sample survey and a community based study. *Sex Transm Infect* 2006;**82**:86–7.
- 26 **Williamson LM, Dodds JP, Mercey DE, et al.** Increase in HIV-related sexual risk behavior among community samples of gay men in London and Glasgow: how do they compare? *J Acquir Immune Defic Syndr* 2006;**42**:238–41.

conflict with UK guidelines. Their preferred regimen for the treatment of pelvic inflammatory disease is in particular surprising (doxycycline and metronidazole) since it conforms with neither UK nor US guidelines and three studies have now shown it to be inferior to alternative regimens.

Overall *Fast facts: sexually transmitted infections* is to be recommended with just a few caveats.

Jonathan Ross

CORRECTIONS

doi: 10.1136/sti.2006.021782.corr1

There was an error in the August issue of the journal (Dodds JP, Johnson AM, Parry JV, *et al.*

A tale of three cities: persisting high HIV prevalence, risk behaviour and undiagnosed infection in community samples of men who have sex with men. *Sex Transm Infect* 2007;**83**:392–6.) The last sentence on page 2 should read as follows: “All were screened by GACELISA HIV 1 and 2, whose sensitivity and specificity had been determined as 99.5% (95% CI 97.1% to 99.9%) and 99.7% (95% CI 98.9% to 99.9%), respectively.”^{13 14}

doi: 10.1136/sti.2006.023283.corr1

Several errors occurred in an article published in the July issue of the journal (Evans AR, Wiggins RD, Mercer CH, *et al.* Men who have sex with men in Great Britain: comparison of a self-selected internet sample with a national probability sample. *Sex Transm Infect* 2007;**83**:200–5.) A corrected version of the

article is available on our website at <http://sti.bmj.com/supplemental>.

doi: 10.1136/sti.2006.023531.corr1

Two articles from the August 2007 supplement were unlocked online but not in print. The articles are: Orroth KK, Freeman EE, Bakker R, *et al.* Understanding the differences between contrasting HIV epidemics in east and west Africa: results from a simulation model of the Four Cities Study. *Sex Transm Infect* 2007;**83**(Suppl 1):i5–i6 and Freeman EE, Orroth KK, White RG, *et al.* Proportion of new HIV infections attributable to herpes simplex 2 increases over time: simulations of the changing role of sexually transmitted infections in sub-Saharan African HIV epidemics. *Sex Transm Infect* 2007;**83**(Suppl 1):i17–i124. Both articles are freely available.